



4-10-1979

AGRICULTURAL BULLETIN



LANNATE® L INSECTICIDE
COTTON

1/1

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

SUPPLEMENTAL LABELING
EPA REG. NO. 352-370

LANNATE® L INSECTICIDE
FOR USE ON COTTON

ACCEPTED
APR 10 1979
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under Reg. No. 352-370

Du Pont "Lannate" L Insecticide is recommended for use on cotton to control lygus bugs.

East of Rocky Mountains

Early Season

Apply 2 pts. "Lannate" L per acre; use as an occasional spray regularly but not more often than every 10 days.

Late Season (after desired boll load set on plants)

Apply 2 to 3 pts. "Lannate" L per acre; use up to 3 applications at 3- to 5-day intervals.

West of Rocky Mountains

Apply 2 to 3 pts. "Lannate" L per acre. Apply at 5- to 7-day intervals or as needed to maintain control.

Apply in sufficient water (minimum 3 gals. per acre by air) to obtain thorough coverage.

Initiate treatment when significant populations of lygus bugs occur.

Note: "Lannate" L may be applied up to 15 days before harvest. Do not graze or feed treated cotton to livestock. Use of "Lannate" L may redden cotton foliage. If reddening is excessive, alternate with other insecticides or discontinue use.

IMPORTANT

BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

3/1/79

