

PLEASE NOTE

This image contains more than one label approved for this product on this date.

352 -370

07/26/1979




LANNATE® L INSECTICIDE
COTTON



AGRICULTURAL BULLETIN

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



SUPPLEMENTAL LABELING
EPA REG. NO. 352-370

LANNATE® L INSECTICIDE
FOR USE ON COTTON
WEST OF THE ROCKY MOUNTAINS

Du Pont Lannate® L insecticide is recommended for control of cotton bollworm and tobacco budworm on cotton west of the Rocky Mountains. When Lannate® L is applied to eggs, control is from direct kill of eggs (ovicide) and newly-hatched larvae (larvicide).

HOW TO USE

Apply 1 pt. Lannate® L per acre in sufficient water (minimum 3 gals. per acre by air) to obtain thorough coverage. Initiate schedule of sprays when significant numbers of eggs are present. Continue at 5- to 7-day intervals while eggs are present and larval control is adequate. If significant numbers of larvae survive the treatment, use higher rates of Lannate® L as registered.

Periodic scouting of fields is required to insure proper timing of spray.

Note: Lannate® L may be applied up to 15 days before harvest. Do not graze or feed treated cotton to livestock. Use of Lannate® L may redden cotton foliage. If reddening is excessive, alternate with other insecticides or discontinue use.

IMPORTANT

BEFORE USING LANNATE® L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

4/19/79

NEXT


LABEL

352-370

7-26-1979




LANNATE® L INSECTICIDE
COTTON



AGRICULTURAL BULLETIN

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



SUPPLEMENTAL LABELING

EPA REG. NO. 352-370

LANNATE® L INSECTICIDE
FOR USE ON COTTON
WEST OF THE ROCKY MOUNTAINS

Du Pont "Lannate" L insecticide is recommended for use on cotton to control tobacco budworm and cotton bollworm west of the Rocky Mountains. Control is obtained through kill of eggs (ovicide) and larvae (larvicide).

HOW TO USE

Apply 2 to 3 pts. "Lannate" L per acre in sufficient water (use a minimum of 3 gals. per acre by air) to obtain thorough coverage. Repeat at 5- to 7-day intervals or as needed.

"Lannate" L may be applied up to 15 days before harvest. Do not graze or feed treated cotton to livestock.

NOTE: Use of "Lannate" L may redden cotton foliage. If reddening is excessive, alternate with other insecticides or discontinue use of "Lannate" L.

IMPORTANT

BEFORE USING "LANNATE" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

4/20/79