

GENERAL INFORMATION

Du Pont "Velpar" Herbicide is a water-soluble powder to be mixed in water and applied as a spray on non-cropland areas. It is non-volatile, non-flammable and non-corrosive to equipment.

"Velpar" is an effective general herbicide providing both contact and residual control of many annual and biennial weeds, woody plants and, except for johnsongrass, is effective for control of most perennial weeds. It is absorbed through the roots and foliage.

Moisture is required to activate "Velpar" in the soil. Best results are obtained when weeds are young and actively growing. Soil is moist at time of application and 1/4 to 1/2 inch of rainfall occurs within 2 weeks after application. Foliar application to weeds is most effective under conditions of high temperature (above 80°F.), high humidity and good soil moisture. Application when vegetation is dormant or semi-dormant may not be effective.

Symptoms usually occur within 2 weeks after application under warm humid conditions, while 4 to 6 weeks may be required when weather is cool. The degree of control and duration of effect will vary with the amount of chemical applied, rainfall, temperature, weed species, soil texture and other conditions.

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Du Pont "Velpar" Herbicide should be used only in accordance with recommendations on this label, or in separate published Du Pont recommendations available through local dealers.

Du Pont will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by Du Pont. User assumes all risk associated with such non-recommended use.

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Warnings must include the following information: "WARNING: Area treated with 'Velpar' on (date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of contact, flush skin or eyes with plenty of water; for eyes, get medical attention."

HOW TO APPLY

Apply "Velpar" as a spray just before or soon after weed emergence. Before spraying, calibrate equipment to determine the quantity of water necessary to uniformly and thoroughly cover the vegetation and soil in a measured area to be treated. Make sure the volume of water is sufficient to completely dissolve the "Velpar" using the Water Temperature/Solubility Table as a guide. Use of concentrations greater than shown in the Water Temperature/Solubility Table may result in nozzle plugging and uneven distribution.

Water Temperature/Solubility Table	
Water Temperature Degrees F.	Solubility Max. Lbs. "Velpar" per 100 Gals.
75	23
70	20
65	18
60	16
55	14
50	13
40	12

Add the proper amount of "Velpar" into a spray tank filled with the amount of water to be used and dissolve using sufficient agitation to prevent the powder from settling. Continue agitation for approximately 10 minutes or until the powder is dissolved. Use a fixed-boom sprayer properly calibrated to a constant speed and rate of delivery. For application with a handgun sprayer, use sufficient water (50 to 200 gals. per acre) for thorough coverage. For treating small areas, a tank-type hand sprayer or sprinkling can may be used. A surfactant such as Du Pont Surfactant WK may be added at the rate of 1 quart per 100 gals. of spray to improve wetting and/or contact activity.

GENERAL WEED CONTROL—NON-CROPLAND AREAS

Railroad, Highway, Utility and Pipeline Right of Ways, Petroleum Tank Farms, Storage Areas, Industrial Plant Sites, Drainage Ditch Banks and other similar areas.

Use the higher levels of the recommended dosage ranges under the following conditions: on hard-to-kill species, on fine-textured soils (clay, clay loam); on soils containing more than 5% organic matter or carbon, and, on coarse-textured soils (sand, loamy sand, gravelly soils) where rainfall exceeds 40 inches per year. Use the lower levels of the dosage range if the growing season is short.

Continued >

4 Continued from Back Panel

Apply 6 to 12 lbs. per acre for season-long control of many annual, biennial and perennial weeds including bahiagrass*, bermudagrass*, bluegrass, broomsedge, camphorweed, Canada thistle*, chickweed, clovers, dewberry, dog fennel, fescue*, fingergrass, foxtail, guineagrass, heath aster, honeysuckle, lantana, marehail*, natagrass, plantain, prickly lettuce, ragweed, smutgrass, Spanish needle, vaseygrass, wild blackberry* and wild carrot.

For short-term control (up to 3 months) of the above weeds, apply 2 to 5 lbs. per acre. Application at 2 to 5 lbs. per acre also provides short-term control of many annual, biennial and perennial weeds including barnyardgrass, bindweed*, bouncing bet*, bromegrass, buffelgrass*, burdock, cocklebur, crabgrass, crown vetch, curly dock*, dandelion*, dogbane*, fiddleneck, filaree, fleabane, goatsbeard vine, goldenrod, lespedeza, milkweed*, mustard, nutsedge*, orchardgrass*, oxalis, paragrass, pigweed, purslane, quackgrass, ryegrass, smartweed, spurge, trumpet creeper*, wild oats*, wild parsnip and wild star thistle.

For woody plant control, apply 4 to 8 lbs. per acre as a broadcast spray for control of maple, oak, poplar, sumac, willow, hawthorn, and hickory. Use at least 40 gals. of spray per acre. Apply during the period between late winter and early summer.

Notice to Buyer: Purchase of this material does not confer any rights under patents of countries outside of the United States.

STORAGE AND DISPOSAL

STORAGE: Store product in original container only, away from other pesticides, fertilizer, food, or feed.

DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

*Considered hard to kill

© 1985 E. I. du Pont de Nemours & Co. (Inc.)

0784 04 23/85

80 9 629

SPECIMEN LABEL
73% OF ACTUAL SIZE

Handwritten initials: "J.K."