

SEP 9 1977
Federal Inspector
of Pesticides
D

ACCET



BENLATE® OD

FUNGICIDE OIL-DISPERSIBLE POWDER
For Use On Bananas

NET
50 LBS.

ACTIVE INGREDIENT
Benomyl [Methyl 1 (butylcarbamoyl) 2 benzimidazolecarbamate] 50%
INERT INGREDIENTS 50%
U.S. Pats. 3,541,211 & 3,631,176 EPA Reg. No. 352-385-AA EPA Est. 352-WV-1

Keep out of reach of children.

CAUTION! MAY IRRITATE EYES, NOSE, THROAT, AND SKIN.

Avoid breathing dust or spray mist
Avoid contact with skin, eyes, and clothing
Wash thoroughly after using Keep away from fire or sparks
In case of contact flush skin or eyes with plenty of water; for eyes, get medical attention

ENVIRONMENTAL HAZARDS This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL Never allow "Benlate" OD to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of "Benlate" OD as a fungicide. Keep container closed when not in use.

Do not contaminate water, food, or feed by storage or disposal. Do not re-use empty container, bury in a safe place away from water supplies. Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning. Open dumping is prohibited.

NOTICE TO BUYER - Purchase of this material does not confer any rights under patents of countries outside of the country in which it was purchased from Du Pont.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Du Pont "Benlate" OD Fungicide is recommended for use as a foliar spray on bananas for the control of Sigatoka disease. It may be used in spray oil alone or in oil-water emulsion, and may be applied with aerial or ground equipment. Apply at the following rates at 2- to 3 week intervals throughout the complete crop cycle; under severe disease conditions, shorten interval and use the higher rates.

Spray Oil Mixture — Use 2 to 4 oz. "Benlate" OD per acre in 1 to 2 gals. of non phytotoxic spray oil per acre.

Oil-Water Emulsion — Use 2 to 4 oz. "Benlate" OD plus 1/2 to 1 gal. of non-phytotoxic spray oil per acre (add emulsifier at rate of 1% of oil volume).

Apply a minimum of 2 1/2 gals. of spray per acre for aerial application, and at least 5 gals. per acre with mist blowers.

"Benlate" OD must be kept in suspension at all times by continuous agitation; agitate by mechanical or hydraulic means.

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

B 21447 8 77

Made in U.S.A.

Printed in U.S.A.

E. I. DU PONT DE NEMOURS & CO. (INC.), BIOCHEMICALS DEPARTMENT, WILMINGTON, DEL.