



VELPAR® L WEED KILLER
FOR CONTROL OF CERTAIN WEEDS IN ESTABLISHED ALFALFA

Add the "Velpar" I. to a water-filled tank and mix thoroughly. Apply in at least 20 gallons of water per acre using ground equipment and 5 to 10 gallons of water per acre by air.

USE RATES

Rate selection is to be based on soil texture and organic matter content as follows:

Soil Texture/Organic Matter	Quarts/Acre
Gravelly or rocky soils, exposed subsoils, hardpan, sand, poorly drained or alkali soils.	Do not use.
Sandy loam, loamy sand, loam, silt, silt loam, and sandy clay loam with less than 1% organic matter.	1 to 1 1/2
Silty clay loam, clay loam, sandy clay, silty clay, and clay with 1% organic matter or higher.	1 1/2 to 3
Soils having more than 5% organic matter. ¹	2 to 3

¹On soils high in organic matter, the effectiveness of "Velpar" L can be significantly reduced and weed control may be unsatisfactory.

WEED SPECIES CONTROLLED

"Velpar" L when applied preemergence or early postemergence will control the following winter annual and spring germinating weeds:

1 to 2 qts./acre

annual bluegrass	fleabane
blue mustard	foxtail
common chickweed	jimsonweed
common groundsel	orchardgrass
common lambsquarter	purple mustard
crabgrass	salsify
fiddleneck	shepherdspurse
field pennycress	spurry
filaree	tansy mustard
	yellow rocket

2 to 3 qts./acre

downy brome or cheatgrass (<i>Bromus tectorum</i>)	annual ryegrass
ivy leaf speedwell	sweet clover
prickly lettuce	purslane speedwell

2 to 3 qts./acre will give partial control² of:

common dandelion	seedling alfalfa
false dandelion	white cockle
Mexican tea	wild oats
perennial bluegrass	

²Partial control is observable reduction in weed competition as compared to an untreated area. Degree of partial control can vary with soil type, environmental factors, and size of weeds at the time of application.

REPLANTING

Except in low annual rainfall areas (20" or less), corn may be planted 12 months after the last treatment provided the soil is deep disked or plowed prior to planting. Do not plant treated areas to any other crop within two years after treatment or injury to crop may result.

NOTE: Because crop injury may result, do not apply to actively growing alfalfa nor to subble between cuttings, except as recommended for certain northeastern* and midwestern* states. Extremely heavy rainfall or more than 1" of irrigation soon after application may result in crop injury or non-performance. Do not use in low desert valleys in California. Do not apply to snow covered or frozen ground.

Do not graze or feed treated forage or hay to livestock within 30 days following application.

*Northeastern and midwestern states are:

Connecticut	Kentucky	Minnesota	Pennsylvania
Delaware	Maine	Missouri	Rhode Island
Illinois	Maryland	New Hampshire	Vermont
Indiana	Massachusetts	New Jersey	Virginia
Iowa	Michigan	New York	West Virginia
		Ohio	Wisconsin

IMPORTANT

BEFORE USING "VELPAR" L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

11/8/82