



AGRICULTURAL BULLETIN




LOROX® DF™ HERBICIDE
CARROTS
FLORIDA

SUPPLEMENTAL LABELING
EPA REG. NO. 352-394

LOROX® DF™ HERBICIDE
FOR PREEMERGENCE IN CARROTS
IN THE STATE OF FLORIDA

DIRECTIONS FOR USE



It is a violation of federal law to use this product
in a manner inconsistent with its labeling.

Du Pont "Lorox" DF™ Herbicide is effective for use a preemergence treatment in carrots in the state of Florida for the control of germinating weeds such as barnyardgrass (watergrass), carpetweed, chickweed, crabgrass, foxtail (including giant foxtail), goosegrass, lambsquarters, mustard, pigweed, purslane, and smartweed. Treatment will not control established perennials such as Canada thistle and purple nutsedge.

HOW TO USE

Preemergence Application (Florida) - Apply "Lorox" DF™ after planting but before carrots emerge; plant seed at least 1/2" deep. Make a single application at the rate of 1 to 2 lbs. per acre (for band treatment, use proportionately less). Use the lower rate on lighter soils (low in clay or organic matter) and the higher rate on heavier soils (high in clay or organic matter).

Note: "Lorox" DF™ is also effective for use in carrots as a postemergence application; see package label for use instructions. Carrots may be treated with multiple applications (pre- and/or postemergence) provided the total does not exceed 4 pounds "Lorox" DF™ per acre.

IMPORTANT

BEFORE USING "LOROX" DF™, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

071686