



AGRICULTURAL BULLETIN



LOROX® DF HERBICIDE
CROP ROTATION

SUPPLEMENTAL LABELING
EPA REG. NO. 352-394

LOROX® DF HERBICIDE
REPLANTING AND CROP ROTATION RECOMMENDATIONS

DIRECTIONS FOR USE

It is a violation of federal law to use this product
in a manner inconsistent with its labeling.

Du Pont Lorox® DF Herbicide is recommended for weed control in certain crops. The following "Replanting" and "Crop Rotation Recommendations" should be observed for treatments applied as directed on container labeling:

REPLANTING

If initial seeding fails to produce a stand, crops registered for the rate of Lorox® DF that has been applied may be replanted into the treated area. Do not retreat during the same crop year as injury to the crop may result.

CROP ROTATION RECOMMENDATIONS:

Unless otherwise directed, any crop may be planted after 4 months except for cereals where only barley, oats, rye, and wheat may be planted.

West of the Rocky Mountains, - carrots or celery may be planted 4 months after last application; do not plant any other crop until 1 year after the last application.

IMPORTANT

BEFORE USING LOROX® DF, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABELS.

This bulletin contains new or supplemental instructions for use of these products which may not appear on the package labels. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.