



AGRICULTURAL BULLETIN

DU PONT

BENLATE® PNW PLUS
MANZATE® 200 FLOWABLE
WHEAT
IDAHO, OREGON, UTAH
AND WASHINGTON

1 of 2

10/21

10/21

SUPPLEMENTAL LABELING
BENLATE® PNW (EPA REG. NO. 352-434)
MANZATE® 200 FLOWABLE (EPA REG. NO. 352-398)

BENLATE® PNW FUNGICIDE PLUS MANZATE® 200 FLOWABLE FUNGICIDE
AS A TANK MIXTURE FOR CONTROL OF CERTAIN DISEASES
OF WHEAT

ACCEPTED
DEC 10 1985
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under
EPA Reg. No. 352-398

DIRECTIONS FOR USE

It is a violation of federal law to use this product
in a manner inconsistent with its labeling.

Du Pont Benlate® PNW Fungicide plus Manzate® 200 Flowable Fungicide are recommended for use in a tank mixture foliar spray for control of Septoria Leaf and Glume Blotch, Helminthosporium Leaf Blight, Leaf Rust and Powdery Mildew on wheat.

HOW TO USE

Leaf Rust and Powdery Mildew

Apply 1/4 to 1/2 lb. of Benlate® PNW plus 1 to 2 qts. Manzate® 200 Flowable per acre. Begin applications when disease first appears or at flag leaf emergence. Repeat at 14-day intervals. Do not make more than three applications. Applications should be timed to keep the flag leaf free of disease. Treatments made after the disease is severe or after the flag leaf becomes infected may not provide economical disease control.

Helminthosporium Leaf Blight, Septoria Leaf and Glume Blotch

Apply 1/4 to 1/2 lb. of Benlate® PNW plus 1 to 2 qts. Manzate® 200 Flowable per acre. Begin applications when disease first appears or at flag leaf emergence. Repeat at 14-day intervals. Applications should be timed to keep the flag leaf and emerging seed head free of disease.

Note: Do not make more than a total of three applications per year.

011085
080985

A nonphytotoxic, superior-type spray oil (60 to 70 second viscosity) at the rate of 1 quart per acre, or a "spreader-sticker" spray adjuvant may be added to improve spray coverage; add oil or adjuvant as last ingredient to spray tank.

Note: Do not apply within 26 days of harvest.
Do not allow livestock to forage in treated fields. Do not tank mix "Benlate" with lime or alkaline pesticides such as Bordeaux mixture or lime sulfur.

IMPORTANT

BEFORE USING "BENLATE" PNW PLUS "MANZATE" 200 FLOWABLE, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABELS.

This bulletin contains new or supplemental instructions for use of these products which may not appear on the package labels. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

011085
080985




BULLETIN

WHEAT
IDAHO, OREGON, UTAH
AND WASHINGTON

SUPPLEMENTAL LABELING
BENLATE[®] PNW (EPA REG. NO. 352-434)
MANZATE[®] 200 FLOWABLE (EPA REG. NO. 352-398)

BENLATE[®] PNW FUNGICIDE PLUS MANZATE[®] 200 FLOWABLE FUNGICIDE
AS A TANK MIXTURE FOR CONTROL OF CERTAIN DISEASES
OF WHEAT



DIRECTIONS FOR USE

It is a violation of federal law to use this product
in a manner inconsistent with its labeling.

Du Pont Benlate[®] PNW Fungicide plus Manzate[®] 200 Flowable Fungicide are recommended for use in a tank mixture foliar spray for control of Septoria Leaf and Glume Blotch, Helminthosporium Leaf Blight, Leaf Rust and Powdery Mildew on wheat.

HOW TO USE

Leaf Rust and Powdery Mildew

Apply 1/4 to 1/2 lb. of Benlate[®] PNW plus 1 to 2 qts. Manzate[®] 200 Flowable per acre. Begin applications when disease first appears or at flag leaf emergence. Repeat at 14-day intervals. Do not make more than three applications. Applications should be timed to keep the flag leaf free of disease. Treatments made after the disease is severe or after the flag leaf becomes infected may not provide economical disease control.

Helminthosporium Leaf Blight, Septoria Leaf and Glume Blotch

Apply 1/4 to 1/2 lb. of Benlate[®] PNW plus 1 to 2 qts. Manzate[®] 200 Flowable per acre. Begin applications when disease first appears or at flag leaf emergence. Repeat at 14-day intervals. Applications should be timed to keep the flag leaf and emerging seed head free of disease.

Note: Do not make more than a total of three applications per year.

011085
080985

A nonphytotoxic, superior-type spray oil (60 to 70 second viscosity) at the rate of 1 quart per acre, or a "spreader-sticker" spray adjuvant may be added to improve spray coverage; add oil or adjuvant as last ingredient to spray tank.

Note: Do not apply within 26 days of harvest. Do not allow livestock to forage in treated fields. Do not tank mix "Benlate" with lime or alkaline pesticides such as Bordeaux mixture or lime sulfur.

IMPORTANT

BEFORE USING "BENLATE" PNW PLUS "MANZATE" 200 FLOWABLE, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABELS.

This bulletin contains new or supplemental instructions for use of these products which may not appear on the package labels. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

011085
080985



AGRICULTURAL BULLETIN

SUTOLIN

1067

BENLATE® PNW PLUS
MANZATE® 200 FLOWABLE
WHEAT
IDAHO, OREGON, UTAH
AND WASHINGTON

SUPPLEMENTAL LABELING
BENLATE® PNW (EPA REG. NO. 352-434)
MANZATE® 200 FLOWABLE (EPA REG. NO. 352-398)

BENLATE® PNW FUNGICIDE PLUS MANZATE® 200 FLOWABLE FUNGICIDE
AS A TANK MIXTURE FOR CONTROL OF CERTAIN DISEASES
OF WHEAT

ACCEPTED
DEC 10 1985
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticides
registered under
EPA Reg. No. 352-398

DIRECTIONS FOR USE

It is a violation of federal law to use this product
in a manner inconsistent with its labeling.

Du Pont Benlate® PNW Fungicide plus Manzate® 200 Flowable Fungicide are
recommended for use in a tank mixture foliar spray for control of Septoria
Leaf and Glume Blotch, Helminthosporium Leaf Blight, Leaf Rust and Powdery
Mildew on wheat.

HOW TO USE

Leaf Rust and Powdery Mildew

Apply 1/4 to 1/2 lb. of Benlate® PNW plus 1 to 2 qts. Manzate® 200
Flowable per acre. Begin applications when disease first appears or at flag
leaf emergence. Repeat at 14-day intervals. Do not make more than three
applications. Applications should be timed to keep the flag leaf free of
disease. Treatments made after the disease is severe or after the flag leaf
becomes infected may not provide economical disease control.

Helminthosporium Leaf Blight, Septoria Leaf and Glume Blotch

Apply 1/4 to 1/2 lb. of Benlate® PNW plus 1 to 2 qts. Manzate® 200
Flowable per acre. Begin applications when disease first appears or at flag
leaf emergence. Repeat at 14-day intervals. Applications should be timed to
keep the flag leaf and emerging seed head free of disease.

Note: Do not make more than a total of three applications per year.

011085
080985

A nonphytotoxic, superior-type spray oil (60 to 70 second viscosity) at the rate of 1 quart per acre, or a "spreader-sticker" spray adjuvant may be added to improve spray coverage; add oil or adjuvant as last ingredient to spray tank.

Note: Do not apply within 26 days of harvest. Do not allow livestock to forage in treated fields. Do not tank mix "Benlate" with lime or alkaline pesticides such as Bordeaux mixture or lime sulfur.

IMPORTANT

BEFORE USING "BENLATE" PNW PLUS "MANZATE" 200 FLOWABLE, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABELS.

This bulletin contains new or supplemental instructions for use of these products which may not appear on the package labels. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

011085
080985