


"OUST" HERBICIDE
EPA REG. NO. 352-401

SUPPLEMENTAL LABELING
"OUST" HERBICIDE
FOR NONCROP WEED CONTROL IN CALIFORNIA



GENERAL

Du Pont "Oust" herbicide is recommended for preemergence and postemergence control of many annual and perennial grasses and broadleaf weeds. "Oust" must be applied in ground equipment as a selective or nonselective spray treatment of noncropland areas such as airports, fencerows, highways, industrial turf, lumberyards, petroleum tank farms, pipeline rights-of-way, plant sites, railroads, roadside turf, storage areas and utility rights-of-way. See the container and/or specimen label for specific recommendations.

Most crops, particularly irrigated crops, are very sensitive to "Oust".

Therefore, to reduce off-target risk, it is essential that great care be taken to apply "Oust" only:

- (a) where it will not physically drift during application,
- (b) where it will not be washed from treated areas, or
- (c) where "Oust" treated soil will not be blown or otherwise be moved into irrigated cropland.

DIRECTIONS FOR USE

It is a violation of federal law to use this product
in a manner inconsistent with its labeling.

Du Pont "Oust" herbicide is recommended for noncrop weed control in California. However, there are several precautions that must be observed to avoid injury to crops, especially irrigated crops which are widespread in California:

- o Do not use in areas where irrigated cropland is next to a potential treatment site, nor where the herbicide can be moved or washed, or where treated soil can be blown, into irrigated cropland as injury to crops is likely to occur.
- o Do not allow spray to drift, during application, onto irrigated cropland where crops are being grown or will be grown.

BEST AVAILABLE COPY

- o Do not apply through any type of irrigation system.
- o Do not apply to impervious substrates such as paved or highly compacted surfaces, nor to frozen ground.
- o Do not apply to soils saturated with water.

USE RATES

Economical treatment programs and less off-target risk can be achieved by surveying the weeds present and using "Oust" rates appropriate for these weeds in California, followed by a spot treatment, or retreatment as necessary to achieve full season control. Also several combination treatments are registered to afford broader spectrum weed control programs.

Based on California data, apply "Oust" as a preemergence treatment or early postemergence treatment during the rainy season at 1 1/3 to 2 ounces* per acre to control:

Annual bluegrass	Italian ryegrass
Annual sowthistle	Prickly coontail
Barnyardgrass	Prickly lettuce
Black mustard	Puncturevine
Buckhorn plantain	Red brome
Burclover	Ripgut brome
Canarygrass	Seashore saltgrass
Cheat	Seaside heliotrope
Chickweed	Signalgrass
Common mallow	Spreading orach
Common speedwell	Sunflower
Common yarrow	Western ragweed
Curly dock	Whitestern filaree
Fiddleneck	Wild oats
Foxtail barley	Yellow foxtail
Foxtail fescue	

Apply at 2 to 3 ounces* per acre to control:

Russian thistle
Smooth brome

* Add surfactant at 0.25% by volume to improve control of emerged weeds.

For other weeds, refer to container and/or specimen label for recommendations.

For broader spectrum weed control than offered by "Oust" alone, tank mixes are recommended with Du Pont "Krovar"[1] 1 DF, "Karmex"[1] DF, "Hyvar"[1] X or "Hyvar" X-L, "Velpar"[1] or "Velpar" L or "Telar"[1] Herbicides; or "Roundup"[2] or 2,4-D.

For season long weed control, "Oust" or "Oust" combinations applied early followed by a contact herbicide applied later for clean-up, may be required.

APPLICATION

Application by an experienced, and/or state certified or licensed applicator is suggested.

Calibrate sprayer to spray coarse droplets and use a drift control agent at the recommended rate to help minimize spray drift.

IMPORTANT

BEFORE USING "OUST", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on the package label. Follow instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Do not apply this product through any type of irrigation equipment.

[1] Registered trademark of E. I. du Pont de Nemours & Co. (Inc.)

[2] Registered trademark of Monsanto Company

E. I. du Pont de Nemours & Co. (Inc.)
Agricultural Products Department
Wilmington, Delaware 19898


101088



"OUST" HERBICIDE
EPA REG. NO. 352-401

SUPPLEMENTAL LABELING
"OUST" HERBICIDE
FOR NONCROP WEED CONTROL IN CALIFORNIA

GENERAL



Du Pont "Oust" herbicide is recommended for preemergence and postemergence control of many annual and perennial grasses and broadleaf weeds. "Oust" must be applied in ground equipment as a selective or nonselective spray treatment of noncropland areas such as airports, fencerows, highways, industrial turf, lumberyards, petroleum tank farms, pipeline rights-of-way, plant sites, railroads, roadside turf, storage areas and utility rights-of-way. See the container and/or specimen label for specific recommendations.

Most crops, particularly irrigated crops, are very sensitive to "Oust".

Therefore, to reduce off-target risk, it is essential that great care be taken to apply "Oust" only:

- (a) where it will not physically drift during application.
- (b) where it will not be washed from treated areas, or
- (c) where "Oust" treated soil will not be blown or otherwise be moved into irrigated cropland.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Du Pont "Oust" herbicide is recommended for noncrop weed control in California. However, there are several precautions that must be observed to avoid injury to crops, especially irrigated crops which are widespread in California:

- o Do not use in areas where irrigated cropland is next to a potential treatment site, nor where the herbicide can be moved or washed, or where treated soil can be blown, into irrigated cropland as injury to crops is likely to occur.
- o Do not allow spray to drift, during application, onto irrigated cropland where crops are being grown or will be grown.

- o Do not apply through any type of irrigation system.
- o Do not apply to impervious substrates such as paved or highly compacted surfaces, nor to frozen ground
- o Do not apply to soils saturated with water.

USE RATES

Economical treatment programs and less off-target risk can be achieved by surveying the weeds present and using "Oust" rates appropriate for these weeds in California, followed by a spot treatment, or retreatment as necessary to achieve full season control. Also several combination treatments are registered to afford broader spectrum weed control programs.

Based on California data, apply "Oust" as a preemergence treatment or early postemergence treatment during the rainy season at 1 1/3 to 2 ounces* per acre to control.

Annual bluegrass	Italian ryegrass
Annual sowthistle	Prickly coontail
Barnyardgrass	Prickly lettuce
Black mustard	Puncturevine
Buckhorn plantain	Red brome
Burclover	Ripgut brome
Canarygrass	Seashore saltgrass
Cheat	Seaside heliotrope
Chickweed	Signalgrass
Common mallow	Spreading orach
Common speedwell	Sunflower
Common yarrow	Western ragweed
Curly dock	Whitestern filaree
Fiddleneck	Wild oats
Foxtail barley	Yellow foxtail
Foxtail fescue	

Apply at 2 to 3 ounces* per acre to control:

Russian thistle
Smooth brome

* Add surfactant at 0.25% by volume to improve control of emerged weeds.

For other weeds, refer to container and/or specimen label for recommendations.

For broader spectrum weed control than offered by "Oust" alone, tank mixes are recommended with Du Pont "Krovar"[1] I DF, "Karmex"[1] DF, "Hyvar"[1] X or "Hyvar" X-L, "Velpar"[1] or "Velpar" L or "Telar"[1] Herbicides; or "Roundup"[2] or 2,4-D.

For season long weed control, "Oust" or "Oust" combinations applied early followed by a contact herbicide applied later for clean-up, may be required.

APPLICATION

Application by an experienced, and/or state certified or licensed applicator is suggested.

Calibrate sprayer to spray coarse droplets and use a drift control agent at the recommended rate to help minimize spray drift.

IMPORTANT

BEFORE USING "OUST", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on the package label. Follow instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Do not apply this product through any type of irrigation equipment.

[1] Registered trademark of E. I. du Pont de Nemours & Co. (Inc.)

[2] Registered trademark of Monsanto Company

E. I. du Pont de Nemours & Co. (Inc.)
Agricultural Products Department
Wilmington, Delaware 19896

101088

