

XXXXXX

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

2 - Bromacil Liquid should be used only in accordance with recommendations on this label, or in separate published manufacturer's recommendations available through local dealers.

Seller will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by seller. User assumes all risk associated with such nonrecommended use.

Apply 2 - Bromacil Liquid Weed Killer just before or during the period of active growth of plants to be controlled when rainfall can be expected for soil activation. If dense growth is present, results will be improved if vegetation is removed before treatment. Do not apply when ground is frozen.

Before spraying, calibrate equipment to uniformly apply the product. Use a fixed boom power sprayer properly calibrated to a constant speed and rate of delivery. For application with a hand gun sprayer, be careful to obtain even distribution of the herbicide to the soil. For treating small areas, a tank type hand sprayer may be used.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

2 - Bromacil Liquid should be used only in accordance with recommendations on this label, or in separate published manufacturer's recommendations available through local dealers.

Seller will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by seller. User assumes all risk associated with such nonrecommended use.

Apply 2 - Bromacil Liquid Weed Killer just before or during the period of active growth of plants to be controlled when rainfall can be expected for soil activation. If dense growth is present, results will be improved if vegetation is removed before treatment. Do not apply when ground is frozen.

Before spraying, calibrate equipment to uniformly apply the product. Use a fixed boom power sprayer properly calibrated to a constant speed and rate of delivery. For application with a hand gun sprayer, be careful to obtain even distribution of the herbicide to the soil. For treating small areas, a tank type hand sprayer may be used.

WEED CONTROL: To control most weeds for an extended period of time on noncropland areas such as RAILROAD, HIGHWAY AND PIPE LINE RIGHT OF WAYS, PETROLEUM TANK FARMS, LUMBERYARDS, STORAGE AREAS, AND INDUSTRIAL PLANT SITES.

Apply 15 to 35 gals. 2 - Bromacil Liquid Weed Killer per acre to control ANNUAL WEEDS and GRASSES such as bromegrass, cheatgrass, crab grass, foxtail, lambsquarters, or hardgrass, plus foxtail, ragweed, rye grass, turkey mullein, and wild oats. When applied just prior to or after emergence of annuals, rates as low as 10 gals. per acre control many annual weeds and grasses in low rainfall areas and give short term control in higher rainfall areas.

Apply 35 to 60 gals. 2 - Bromacil Liquid Weed Killer per acre to control PERENNIAL WEEDS and GRASSES such as aster, bahiagrass, bluegrass, broomsedge, timothy, and alfalfa, plus foxtail, plantain, purpletop, blackgrass, reed can, or bromegrass, and wild carrot. In areas with low or seasonal rainfall, rates as low as 30 gals. per acre control many perennial weeds and grasses.

ATTENTION: IN RIGHT PANEL

2% BROMACIL LIQUID WEED KILLER

WATER SOLUBLE LIQUID

Active Ingredient
2% Bromacil
Inert Ingredients
98%

Keep out of reach of children
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS

WARNING! HARMFUL OR FATAL IF SWALLOWED. CAUSES EYE IRRITATION. MAY IRRITATE NOSE, THROAT AND SKIN.
Do not get in eyes. Avoid contact with skin and clothing. Avoid breathing vapors. Wash thoroughly with soap and water after handling and before eating or smoking.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas that contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL AND CHEMICAL HAZARDS

Keep container closed when not in use. Product will freeze at temperatures below 32°F.

IMPORTANT — Injury to or loss of desirable trees or other plants may result from failure to observe the following: Do not apply, drain, or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants. Do not contaminate domestic waters. Keep from contact with fertilizers, insecticides, fungicides and seeds. Thoroughly clean all traces of 2% Bromacil Liquid Weed Killer from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens. Clean these parts separately.

GENERAL INFORMATION — 2% Bromacil Liquid Weed Killer is a water-soluble liquid to be sprayed undiluted on non-riparian areas. It is an effective, general herbicide for control of brush and many annual and perennial weeds and is particularly useful for control of perennial grasses. Effects are slow to appear and may not be noticeable until the chemical has been carried into the root zone of the weed by rain or irrigation. Control and duration of effect will vary with the amount of chemical applied, soil type, rainfall, and other conditions.

2% Bromacil Liquid Weed Killer is non-volatile, non-flammable, and non-corrosive to metals except aluminum. Do not use with a water mist spray nozzle.

NET 5 GALS.

JOHN DOE CHEMICAL CO. • ANYPLACE IN THE U.S.A.

CONTINUED FROM LEFT PANEL

Apply 5 to 130 gals. 2- Bromacil Liquid Weed Killer per acre to control JOHNSONGRASS. Use at the same rate for OTHER HARD TO KILL PERENNIAL WEEDS and GRASSES such as bermudagrass, boundingbriar, bracken fern, dallisgrass, daphne, forsetia, nutsedge, saltgrass, and wiseegg. Where limited rainfall (usually less than 4 inches) occurs during the active growth period, such as some areas of the West, 2- Bromacil Liquid Weed Killer will not provide satisfactory control of hard to kill, deep rooted perennial weeds such as johnsongrass.

Use the higher levels of the dosage ranges on adsorptive soils (usually those high in organic matter or carbon).

Retreatment: Apply 10 to 35 gals. 2- Bromacil Liquid Weed Killer per acre when annual weeds and grasses reappear in sites where weed growth has been controlled.

For small areas: 15 qts. of 2- Bromacil Liquid Weed Killer per 1,000 sq. ft. is approximately 54 gals. per acre.

BRUSH CONTROL: To control undesirable woody plants on noncrop land areas such as RAILROAD RIGHT OF WAYS, STORAGE AREAS, INDUSTRIAL PLANT SITES, AND DRAINAGE DITCHES.

Apply in spring or summer as a broadcast or basal (soil) treatment. For use on drainage ditches, apply as a basal (soil) treatment only.

Note: For effective brush control and prevention of damage to desirable vegetation, do not apply to brush standing in water, do not use water from treated ditches for irrigation, do not use in irrigation ditches nor on right of ways or other sites where marketable timber or other desirable trees or shrubs are immediately adjacent to the treated area.

Broadcast Treatment: Apply 25 to 55 gals. per acre to control oak, pine, sweet gum, and willow; apply 65 to 130 gals. per acre to control brush such as American elm, cottonwood, hackberry, sumac, and winged elm. Use the higher levels of the dosage ranges on adsorptive soils (those high in organic matter or carbon).

Basal (Soil) Treatment: Apply at the rate of 1 to 4 fl. oz. per stem 2" to 4" in basal diameter, wet base of stem to run off. Treatment controls woody plants such as cottonwood, hackberry, maple, oak, poplar, red bud, sweet gum, wild cherry, willow, and winged elm.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

STORAGE AND DISPOSAL

- 1 Prohibitions:** Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited. Avoid contamination of feed and food stuff.
- 2 Pesticide Disposal:** Pesticide or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.
- 3 Container Disposal:** Triple rinse container. Then offer for recycling, reconditioning, or dispose of in a sanitary landfill in accordance with all laws by state and local authorities.