


SPECIMEN LABEL

ACTUAL SIZE



Pesticide,
water reactive NA 2210

ORM-C

Keep dry

MANEB 80

FUNGICIDE

ACTIVE INGREDIENT:
 Maneb (Manganese ethylenebisdithiocarbamate) 80.0%
 (Manganese equivalent as metallic 16.5%)
INERT INGREDIENTS 20.0%

GENERAL INFORMATION: "Maneb 80 Fungicide" may be used as a spray or dust for the control of many important plant diseases. "Maneb 80 Fungicide" has good adhesive properties on plants, high fungicidal efficiency and is easy to use. It is particularly effective against many of the fungus diseases of certain vegetable, fruit and ornamental crops.

"Maneb 80 Fungicide" may be used undiluted for treatment of certain crop seeds.

Keep out of reach of children.
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS

CAUTION! MAY IRRITATE EYES, NOSE, THROAT, AND SKIN
 MAY BE HARMFUL IF INHALED OR SWALLOWED.

Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. Keep away from fire or sparks.

Wear protective clothing including long pants, long sleeve shirt, gloves, hat and boots during mixing and loading operations.

Do not apply this product in such a manner as to directly or through drift expose workers, other persons, or animals.

First Aid: In case of contact, flush skin or eyes with plenty of water. For eyes, get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply where run-off is likely. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of waste.

IMPORTANT—Never allow "Maneb 80 Fungicide" to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of "Maneb 80 Fungicide" as a fungicide and create vapors which may be flammable. Keep container closed when not in use.

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only, when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

DIRECTIONS FOR USE

CARROTS: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

WATERMELONS: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

CUCUMBERS: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

CANTALOUPE and SQUASH: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

LETTUCE and ENDIVE: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

LIMA BEANS: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

ONIONS: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

SPINACH: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

PEACHES and NECTARINES: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

APRICOTS: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

GRAPES: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

ROSES: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

SEED TREATMENT — WHEAT, BARLEY, RYE, FLAX: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

APPLES: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

CABBAGE: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

CARROTS: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

WATERMELONS: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

CUCUMBERS: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

CANTALOUPE and SQUASH: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

LETTUCE and ENDIVE: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

LIMA BEANS: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

ONIONS: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

SPINACH: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

PEACHES and NECTARINES: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

APRICOTS: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

GRAPES: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

ROSES: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

SEED TREATMENT — WHEAT, BARLEY, RYE, FLAX: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

APPLES: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

CABBAGE: Apply when the plants are young. Apply when the plants are young. Apply when the plants are young.

LABEL TREATED SEED

STORAGE AND DISPOSAL

**PRECAUTION STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

CAUTION! MAY IRRITATE EYES, NOSE, THROAT, AND SKIN.
MAY BE HARMFUL IF INHALED OR SWALLOWED.

Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. Keep away from fire or sparks.

Wear protective clothing including long pants, long sleeve shirt, gloves, hat and boots during mixing and loading operations.

Do not apply this product in such a manner as to directly or through drift expose workers, other persons, or animals.

First Aid: In case of contact, flush skin or eyes with plenty of water, for eyes, get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

IMPORTANT— Never allow "Maneb 80 Fungicide" to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of "Maneb 80 Fungicide" as a fungicide and create vapors which may be flammable. Keep container closed when not in use.

"Maneb 80 Fungicide" may be used undiluted for treatment of certain crop seeds.

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

DIRECTIONS FOR USE

This is a list of Federal law (FIFRA) registered uses of Maneb 80 Fungicide. It should be used only in accordance with the label on separate packages of Du Pont Maneb 80 Fungicide. Du Pont will not be responsible for loss of effectiveness of Maneb 80 Fungicide in any manner if specifically recommended by the label of another pesticide registered with the Federal Government.

Do not use on food crops until the crop has been thoroughly washed and has settled. Because of its status as a general fungicide, it is not intended for various crops treated with this product. Consult the State Department of Agriculture for further information.

Written or oral warnings must be given to the operator of the equipment used in an area about to be treated with this fungicide. Written or oral warnings must be given if there is reason to believe that written warnings are not being observed. Warnings must include the following information: All crops are to be treated with Maneb 80 Fungicide on (date of application). Do not enter with or apply to the protected crop until sprays have dried. Dusts should be applied to the crop with plenty of water for eyes, get medical attention.

SPRAYS: Before there is a disease problem, spray the crop with Maneb 80 Fungicide in a sufficient water to provide 100 to 200 gallons per acre. For most crops, application should be 10 to 15 gallons per acre. For severe disease problems when weather conditions favor disease, spray at 15 to 20 gallons per acre. Wash sprayer thoroughly after each use. Shorten intervals between applications. Wash sprayer thoroughly after the filter screen while tanks are empty with water. After sprayer is clean, drain and dry it.

DUSTS: Maneb 80 Fungicide may be used as a dust. It is not intended for use as an active ingredient otherwise. It is not intended for use in conjunction with pesticides or some other neutral dust. Use dusts in accordance with the label on separate packages. Commonly used insecticides may be used in conjunction with Maneb 80 Fungicide in the dust mixture.

ALMONDS: Do not use on almonds until the crop is thoroughly washed and has settled. Use 10 to 15 gallons per acre for most crops. For severe disease problems, spray at 15 to 20 gallons per acre. Wash sprayer thoroughly after each use. Shorten intervals between applications.

TOMATOES: Do not use on tomatoes until the crop is thoroughly washed and has settled. Apply when the crop is in the early stages of growth. Repeat applications at 7 to 10 day intervals. Shorten intervals between applications when weather conditions favor disease. Do not use on tomatoes when plants are in the late stages of growth. Do not use on tomatoes when plants are in the late stages of growth. Do not use on tomatoes when plants are in the late stages of growth.

POTATOES: Do not use on potatoes until the crop is thoroughly washed and has settled. Apply when the crop is in the early stages of growth. Repeat applications at 7 to 10 day intervals. Shorten intervals between applications when weather conditions favor disease.

APPLES: Do not use on apples until the crop is thoroughly washed and has settled. Apply when the crop is in the early stages of growth. Repeat applications at 7 to 10 day intervals. Shorten intervals between applications when weather conditions favor disease.

CABBAGE: Do not use on cabbage until the crop is thoroughly washed and has settled. Apply when the crop is in the early stages of growth. Repeat applications at 7 to 10 day intervals. Shorten intervals between applications when weather conditions favor disease.

CELERY: Do not use on celery until the crop is thoroughly washed and has settled. Apply when the crop is in the early stages of growth. Repeat applications at 7 to 10 day intervals. Shorten intervals between applications when weather conditions favor disease.

CARROTS: Do not use on carrots until the crop is thoroughly washed and has settled. Apply when plants are 6 weeks old or when disease first appears. Repeat every 7 to 10 days.

WATERMELONS: Do not use on watermelons until the crop is thoroughly washed and has settled. Apply when plants are 6 weeks old or when disease first appears. Repeat every 7 to 10 days. Under severe conditions, repeat every 5 to 7 days.

CANTALOUPE: Do not use on cantaloupes until the crop is thoroughly washed and has settled. Apply when plants are 6 weeks old or when disease first appears. Repeat every 7 to 10 days.

LETTUCE: Do not use on lettuce until the crop is thoroughly washed and has settled. Apply when plants are 6 weeks old or when disease first appears. Repeat every 7 to 10 days. Under severe conditions, repeat every 5 to 7 days.

LIMA BEANS: Do not use on lima beans until the crop is thoroughly washed and has settled. Apply when plants are 6 weeks old or when disease first appears. Repeat every 7 to 10 days.

ONIONS: Do not use on onions until the crop is thoroughly washed and has settled. Apply when plants are 6 weeks old or when disease first appears. Repeat every 7 to 10 days.

SPINACH: Do not use on spinach until the crop is thoroughly washed and has settled. Apply when plants are 6 weeks old or when disease first appears. Repeat every 7 to 10 days.

PEACHES: Do not use on peaches until the crop is thoroughly washed and has settled. Apply when plants are 6 weeks old or when disease first appears. Repeat every 7 to 10 days. Under severe conditions, repeat every 5 to 7 days.

APRICOTS: Do not use on apricots until the crop is thoroughly washed and has settled. Apply when plants are 6 weeks old or when disease first appears. Repeat every 7 to 10 days.

GRAPES: Do not use on grapes until the crop is thoroughly washed and has settled. Apply when plants are 6 weeks old or when disease first appears. Repeat every 7 to 10 days.

ROSES: Do not use on roses until the crop is thoroughly washed and has settled. Apply when plants are 6 weeks old or when disease first appears. Repeat every 7 to 10 days.

SEED TREATMENT — WHEAT, BARLEY, RYE, FLAX: Do not use on wheat, barley, rye, or flax until the crop is thoroughly washed and has settled. Apply when plants are 6 weeks old or when disease first appears. Repeat every 7 to 10 days.

LABEL TREATED SEED: Do not use on label treated seed until the crop is thoroughly washed and has settled. Apply when plants are 6 weeks old or when disease first appears. Repeat every 7 to 10 days.

STORAGE AND DISPOSAL

PESTICIDE DISPOSAL: Do not use on pesticides until the crop is thoroughly washed and has settled. Apply when plants are 6 weeks old or when disease first appears. Repeat every 7 to 10 days.

CONTAINER DISPOSAL: Do not use on containers until the crop is thoroughly washed and has settled. Apply when plants are 6 weeks old or when disease first appears. Repeat every 7 to 10 days.

NET 50 LBS.

Made By Du Pont in Colombia

E. I. DU PONT DE NEMOURS & CO. (INC.), AGRICULTURAL CHEMICALS DEPT., WILMINGTON, DE