


SPECIMEN LABEL



Classic[®]

HERBICIDE



MAR 1 2 1957

Handwritten signature

DISPERSIBLE GRANULES

Classic[®]
 HERBICIDE
 DISPERSIBLE GRANULES

ACTIVE INGREDIENT:

Chlorimuron Ethyl: 2-[[[[(4-chloro-6-methoxypyrimidin-2-yl)amino]-
carbonyl]amino]sulfonyl]benzoate 25%

INERT INGREDIENTS 75%

Net Wt. 4.0 lb (1.8 kg)

EPA Reg. No. 352-41

EPA EST. 352-WV-1

KEEP OUT OF REACH OF CHILDREN
PRECAUTIONARY STATEMENTS — HAZARDS TO HUMANS

CAUTION! MAY IRRITATE EYES, NOSE, THROAT AND SKIN.

May be harmful if absorbed through skin. Avoid breathing dust or spray mist.

Avoid contact with skin, eyes and clothing. Get medical attention if irritation persists. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

For medical emergencies involving this product, call toll free 1-800-441-3637.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

STORAGE: Store product in original container only, away from other pesticides, fertilizer, food or feed.

DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and crush into small pieces, and dispose of in accordance with state and local authorities, by burning. If burned, do not breathe fumes.

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated on such label only when used in accordance with the directions under normal use conditions. It is not a guarantee of performance, and any losses or damages resulting from the use of this product are not covered by this warranty. Crop injury, pest resistance, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or other manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential or special damages resulting from the use or handling of this product. A full and complete description of the product and its use is on the label. DUPONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, AND THERE ARE NO EXPRESS OR IMPLIED WARRANTIES OF A STATE OF ORIGIN.

E. I. du Pont de Nemours & Co. (Inc.), Agricultural Products Dept., Wilmington, DE

IMPORTANT

Injury to or loss of desirable trees or vegetation may result from failure to observe the following. Do not apply or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water. Keep from contact with fertilizers, insecticides, fungicides, and seeds during storage.

Thoroughly clean all traces of "Classic" Herbicide from application equipment immediately after use and prior to spraying crops other than soybeans.

GENERAL INFORMATION

Du Pont "Classic" Herbicide is a dispersible granule formulation to be mixed with water and sprayed for selective postemergence weed control in soybeans. When applied according to the instructions on this label, it will control many broadleaf weeds and yellow nutsedge.

IMPORTANT: Prior to using "Classic", consideration should be given to crop rotation plans. Crops other than soybeans can be extremely sensitive to low concentrations of "Classic" remaining in the soil the next planting season. Choice of rotation crops is restricted following application of "Classic". Crop rotation may be further restricted if "Classic" is applied over Du Pont Canopy*, Gemini*, Preview* or Lorox Plus* Herbicides (See "CLASSIC RECROP TABLE"). Do not apply to soil with a pH greater than 8 except in those states as directed in the "CLASSIC RECROP TABLE", as extended soil residual activity could adversely affect crop rotation plans beyond indicated intervals.

"Classic" will provide best postemergence results when applied to young, actively growing weeds. Degree of control, and timing of effect depend on rate used, weed spectrum, weed size, growing conditions at and following time of treatment, soil moisture, treatment and spray adjuvants. **Some weeds, such as pigweed, are more affected by stress than others. Delay application until stress passes and weeds start to grow again. If weeds get large, use higher rate and spray volume. Treating weeds under stress or large weeds may result in only partial control.**

Because most crops other than soybeans are highly sensitive to "Classic", all direct or indirect contact (such as spray drift) to crop other and soil known to be planted during the current growing season to crops other than soybeans should be avoided.

BIOLOGICAL ACTIVITY

"Classic" rapidly inhibits the growth of susceptible weeds. Following postemergence application, leaves of susceptible plants appear yellow in 3-5 days followed by death of the growing point. Susceptible plants are controlled in 7-21 days. Death of leaf tissue and growing point will follow in some species while others will remain green but stunted and noncompetitive.

DIRECTIONS FOR USE

"Classic" is to be used only in the manner indicated on this label. Do not use in a manner inconsistent with its labeling.

SPRAY TANK PREPARATION

Fill spray tank with water, add "Classic" in the spray tank before adding any other material (e.g., surfactant). Add surfactant to spray tank in the order listed in Appendix "Classic" spray preparation within 24 hours of product mixing. Do not add surfactant until after "Classic" has been added. Do not agitate before using. Allow to settle.

APPLICATION RECOMMENDATION

Applications should be made by ground application only to actively growing weeds. Applications of "Classic" must include a surfactant. Use a surfactant that is compatible with "Classic" and that does not cause phytotoxicity or injury to the crop. Do not use a surfactant that causes phytotoxicity or injury to the crop.

Do not spray "Classic" on a wet surface. Do not spray "Classic" on a surface that is wet with rain or irrigation water. Do not spray "Classic" on a surface that is wet with dew or frost. Do not spray "Classic" on a surface that is wet with snow or ice. Do not spray "Classic" on a surface that is wet with oil or grease. Do not spray "Classic" on a surface that is wet with any other liquid.

- APPLICATION**
- minimum 10 gallons water per acre
 - use flat fan nozzle at 25-40 psi or hollow cone nozzle at 40 to 60 psi
 - use proper boom height for good weed coverage

Do not use "Classic" on a surface that is wet with rain or irrigation water. Do not use "Classic" on a surface that is wet with dew or frost. Do not use "Classic" on a surface that is wet with snow or ice. Do not use "Classic" on a surface that is wet with oil or grease. Do not use "Classic" on a surface that is wet with any other liquid.

Do not use "Classic" on a surface that is wet with rain or irrigation water. Do not use "Classic" on a surface that is wet with dew or frost. Do not use "Classic" on a surface that is wet with snow or ice. Do not use "Classic" on a surface that is wet with oil or grease. Do not use "Classic" on a surface that is wet with any other liquid.

Do not use "Classic" on a surface that is wet with rain or irrigation water. Do not use "Classic" on a surface that is wet with dew or frost. Do not use "Classic" on a surface that is wet with snow or ice. Do not use "Classic" on a surface that is wet with oil or grease. Do not use "Classic" on a surface that is wet with any other liquid.

For band application use proportionately less. Carefully calibrate band applicator to not exceed labeled rates or injury may occur. Multiple nozzles directed at soybean foliage may increase soybean phytotoxicity.

Do not use "Classic" on a surface that is wet with rain or irrigation water. Do not use "Classic" on a surface that is wet with dew or frost. Do not use "Classic" on a surface that is wet with snow or ice. Do not use "Classic" on a surface that is wet with oil or grease. Do not use "Classic" on a surface that is wet with any other liquid.

POSTEMERGENCE APPLICATION GUIDE

"Classic" can be applied after first trifoliate until 60 days before soybean harvest

Weeds Controlled	Application rate			
	1/2 oz/acre		3/4 oz/acre	
	Leaves†	Inches	Leaves†	Inches
Bristly starbur	2	2	4	4
Cocklebur	6	6	10	12
Florida beggarweed	3	4	6	6
Hemp sesbania	3	4	4	6
Jimsonweed	4	4	6	6
*Morningglory (annual)				
Entireleaf	2	2	3	4
Ivyleaf	2	2	3	4
Pitted	2	2	3	4
Smallflower	2	2	3	4
Tail	2	2	3	4
Mustard	4	4 (Diameter)	6	6 (Diameter)
Pigweed				
Palmer	4	2	6	4
Redroot	4	2	6	4
Smooth	—	—	3	2
Tail waterhemp	—	—	3	2
Ragweed				
Common	4	2	6	3
*Giant	4	4	6	6
*Sicklepod	1	2	2	3
Smartweed				
Ladysthumb	4	2	6	3
Pennsylvania	4	2	6	3
Sunflower	6	5	8	8
*Wild poinsettia	2	2	4	4
Yellow nutsedge	4	3	6	4
Weeds Suppressed				
*Burdock	2	2	4	6
*Cowpea	2	4	4	6
Marestail	4	3	6	6
Purple nutsedge	4	3	6	4
Pigweed				
Smooth	3	2	6	4
Tail waterhemp	3	2	6	4
*Velvetleaf††	4	2	6	4

*See "CLASSIC" SPLIT APPLICATION section

† When counting leaves for correct application time, do not count the cotyledons. Count the first true leaves to determine correct stage to spray

†† The addition of ammonium nitrogen fertilizers such as 1 gallon 28% nitrogen A or 1 to 2 qts 10-34-0 A improves "Classic" activity on velvetleaf. A surfactant must also be used with the fertilizer. A cultivation at approximately 14 days after application will improve velvetleaf control.

Spraying weeds in cotyledon stage is not recommended. "Classic" sprayed at stage indicated will either control or suppress (stunt) the weeds, delaying growth for 2 to 3 weeks. Weeds larger than on the label will be suppressed but may recover after 2-3 weeks. A cultivation approximately 14 days after application but before regrowth will greatly aid control.

Weed suppression is a visual reduction in weed competition (reduced population, size, and/or vigor) as compared to an untreated area. Degree of control can be increased by treating weeds when actively growing (not under stress) and by using maximum use rate.

Temporary yellowing and/or retardation of soybean growth may occur following application of "Classic". These effects will generally be most evident 5-7 days after application to soybeans under stress. Under favorable soybean growing conditions, the crop will quickly recover.

Cultivation before, during or within 3 days after the application is not recommended. Cultivation may put weeds under stress by pruning roots, thus making control more difficult to obtain. The best time to cultivate is approximately 14 days after application.

Under stress conditions, delay application until stress passes and weeds resume growth.

"CLASSIC" SPLIT APPLICATIONS

Some weeds with multiple germination times and suppressed (stunted) weeds may require two applications of "Classic". Applications should be timed 14 to 21 days apart or when sufficient regrowth has occurred to require a second treatment. A cultivation approximately 14 days after treatment may replace the need for the second application. Weeds such as burdock, cocklebur, cowpea, giant ragweed, morningglory, pigweed and sicklepod may require a second application.

"CLASSIC" RECROP TABLE

TIME INTERVAL BEFORE REPLANTING IN MONTHS

(from last "Classic" application)

SECTION I: "CLASSIC"

Group A

- All states
Soil pH 6.8 or less
—OR—
■ States of AL, AR, FL, GA, LA, MS, TX with soil
pH 6.9-7.5

Group B

- States of DE, MD, NJ, PA, NC, SC, WV, VA,
KY, TN and MO Bootheel with soil pH 6.9-7.5

Crops

Minimum Time in Months Before Planting Recrop*

Group A

Group B

	Anytime	Anytime
Soybeans	3	3
Cereal grains	3	3
Ryegrass	3	3
Field Corn	9	18
Cotton	9	18
Aufala	9	18
Clover	9	18
Dry beans	9	18
Sorghum	9	18
Peanut	9	18
Rice	9	18

All other crops require a successful test broadcast.

Suggested interval before initiating broadcast is 9 months from Group A and 18 months from Group B.

In states of AL, AR, FL, GA, LA, MS and TX with soil pH 6.8 or less, field corn can be recropped after 8 months.

SECTION II: "CLASSIC" FOLLOWING "CANOPY" OR "GEMINI" HERBICIDE

Group C

Group D

- States of AL, AR, FL, GA, LA, MS, TX with soil
pH 6.8 or less —OR—
■ States of DE, MD, NJ, NC, PA, SC, VA, WV,
KY, TN, MO Bootheel with soil pH 6.8 or less
AND

- All other states with soil pH 6.8 or less
—OR—
■ States listed in Section II Group C not meet-
ing pH, rate, or rainfall requirement or receiv-
ing more than one "Classic" application

- "Gemini" use rates 16 oz A or less, or
- "Gemini" use rate 16-22 oz A and normal
rainfall, or
- "Canopy" use rate 7 oz A or less, or
- "Canopy" use rate 7-10 oz A and normal
rainfall

AND

- Only one "Classic" application per season or
recrop injury may occur

Crops*

Minimum Time in Months Before Planting Recrop*

Group C

Group D

	Anytime	Anytime
Soybeans	3	3
Cereal grains	3	3
Ryegrass	3	3
Field Corn	9	18
Cotton	9	18
Aufala	9	18
Clover	9	18
Dry beans	9	18
Sorghum	9	18
Peanut	9	18
Rice	9	18

All other crops require a successful test broadcast.

Suggested interval before initiating broadcast is 9 months for Group C and 18 months for Group D.

*Consult "Canopy" and "Gemini" labels on recrop intervals. Some crop rotations may be longer than indicated here for one or the other products. Use the longest interval.

SECTION III: "CLASSIC" FOLLOWING "PREVIEW" OR "LOROX PLUS" HERBICIDE

Group E

Group F

- "Preview" or "Lorox Plus" rates greater than
16 oz A or less
and
— Normal rainfall
and
— One "Classic" application per season

- "Preview" or "Lorox Plus" rates greater than
16 oz A or less
or
■ Not meeting normal rainfall requirements
or
■ More than one "Classic" application

Crops*

Minimum Time in Months Before Planting Recrop*

Group E

Group F

	Anytime	Anytime
Soybeans	3	3
Cereal grains	3	3
Ryegrass	3	3
Field Corn	9	18
Cotton	9	18
Aufala	9	18
Clover	9	18
Dry beans	9	18
Sorghum	9	18
Peanut	9	18
Rice	9	18

All other crops require a successful test broadcast.

Suggested interval before initiating broadcast is 9 months for Group E and 18 months for Group F.

*Consult "Preview" and "Lorox Plus" labels on recrop intervals. Some crop rotations may be longer than indicated here for one or the other products. Use the longest interval.

- ⁴ If "Classic" is applied after August 1, extend recrop interval on corn, cotton, rice and sorghum 2 months or recrop may be injured.
- ⁵ Field Bioassay--The season before planting any crop not listed above, a successful field bioassay must be completed. The field bioassay will detect small quantities of "Classic" which can remain in the soil and injure rotational crops. A successful field bioassay means growing to maturity a test strip of the crop(s) intended for production the following year. **The test strip should cross the whole field including knolls and low areas. Suggested interval before planting bioassay crop is listed on "Recrop Table"**
- ⁶ Normal rainfall is the 30 year rainfall average for the field location for the months of April through October. **This 7 month period (April to October) must receive a total rainfall 85% of normal or greater with no two consecutive months receiving less than 50% of normal.**

SPRAYER CLEANUP

IMPORTANT: To avoid subsequent injury to crops other than soybeans, immediately after spraying and prior to spraying other crops, thoroughly remove all traces of "Classic" from mixing and spray equipment. For more information, consult supplement booklet on sprayer cleanup following use of Du Pont "Classic", "Canopy", "Preview", "Gemini" or "Lorox Plus" Herbicides.

PRECAUTIONS

- Do not apply "Classic" before first trifoliolate opens or later than 60 days before soybean maturity.
- Do not apply "Classic" on soil with a pH greater than 6.8 except in those states as directed in the "CLASSIC" RECROP TABLE" section.
- Do not graze treated fields or harvest for forage or hay.
- Do not apply "Classic" to plants under stress from abnormal weather (hot or cold) or growing conditions, drought, water saturated soil, disease, insect or prior herbicide injury, as crop injury or poor weed control may result. Severe stress, drought, disease, insect damage or nutrient deficiency following application may also result in crop injury. **Delay application until stress passes and weeds and soybeans resume growth.**
- Do not apply "Classic" to Black Belt Soils of Alabama and Mississippi with a soil pH greater than 7.0 or history of nutrient deficiency such as iron chlorosis or injury may occur.
- Do not apply "Classic" if rain is expected within 4 hours of application as control may be unsatisfactory.
- Follow extended crop rotation interval as directed on this label when "Classic" is applied over "Canopy", "Gemini", "Preview" or "Lorox Plus" Herbicides.
- If "Scepter" Herbicide is used the same year as "Classic" extend recrop interval one month past the most restrictive interval for crop to be grown on the two labels or recrop injury may occur.
- "Classic" may cause antagonism to post grass herbicides such as "Poast",² "Fusilade"³ or "Assure"⁴ when tank mixed or sprayed within 7 days prior to or 1 day after the grass herbicide. Antagonism is more apt to occur under stress conditions.
- Do not apply in areas north of the Platte River in NE or west of Highway 10 in NE and highway 81 in KS or west of Interstate 35 in OK and TX as the rotational crop guidelines may not be applicable.
- NOTE:** It is a violation of federal law to use this product at rates above those recommended on this label. Crop injury will also result if rates higher than recommended are used. Applications later than 60 days before soybean maturity will be uneconomical and are not recommended as they may be ineffective.
- NOTICE TO BUYER:** Purchase of this material does not confer any rights under patents of countries outside of the United States.
- ¹Trademark of American Cyanamid Corp., Princeton, NJ.
- ²Trademark of BASF Wyandotte Corporation.
- ³Trademark of ICI Americas, Inc.
- ⁴Trademark of E. I. du Pont de Nemours & Co., Inc.

BEST AVAILABLE COPY