


"CLASSIC"
HERBICIDE
BROADLEAF WEED CONTROL
IN SOYBEANS

"DRAFT, NOT APPROVED"

"CLASSIC" HERBICIDE
EPA REG. 352-436

SUPPLEMENTAL LABELING
"CLASSIC" HERBICIDE
FOR PREEMERGENCE OR PREPLANT INCORPORATED
BROADLEAF WEED CONTROL IN SOYBEANS



DIRECTIONS FOR USE

It is a violation of federal law to use this product
in a manner inconsistent with its labeling

GENERAL INFORMATION

DuPont "Classic" Herbicide is a dispersible granule formulation to be mixed with water and sprayed for selective preemergence and preplant incorporated weed control in soybeans. When applied according to this label, it will control many broadleaf weeds and nutsedge.

EQUIPMENT-SPRAY VOLUMES

Apply uniformly by ground equipment with a properly calibrated low pressure(20-40 psi) fixed-boom sprayer equipped with fan-type nozzles and screens no finer than 50 mesh. Use 10-40 gallons of water per acre. Continuous agitation in the spray tank is required to keep the material in suspension. Avoid overlap and shut off spray booms while starting, turning, slowing or stopping, as injury to the crop may result.

PREPLANT OR PREPLANT INCORPORATED

"Classic" can be applied at planting or up to 30 days prior to planting either preemergence or preplant incorporated. If incorporated, uniformly incorporate the treatment no deeper than the top 1-2" of soil prior to planting soybeans. Use equipment suited to proper incorporation into the top 1-2" of soil, eg: Do-all, field cultivator, rotterra. Rainfall (~1") is required to activate both preemergence and preplant incorporated treatments.

*Rainfall required to activate pre or ppi treatments depends on soil moisture when activating rainfall or irrigation is received. Several small rainfalls of less than 1/4" are not as beneficial as one large rain of 1/2" or more for activation.

BEST AVAILABLE COPY

RATE CHART

Apply as follows:

<u>SOIL TEXTURE</u>	<u>BROADCAST RATE(Oz.product/acre)</u>
<u>Course:</u> loamy sand, sandy loam	2 ounces
<u>Medium:</u> loam, silt loam, silt, sandy clay loam	4 ounces
<u>HEAVY:</u> Silty clay loam, clay loam, clay	6 ounces

For band applications, use proportionately less than the rates listed above.


Plant seed 1 1/2" to 2" deep on a flat or raised seedbed only or crop injury may occur.

Soybean stunting may occur if excessive rainfall occurs after application but before soybeans germinate. Injury is more prevalent under poor drainage or compacted conditions or when soil is saturated for long periods of time. Soybeans outgrow stunting once favorable growing conditions return.

DO NOT APPLY TO SOILS OF pH GREATER THAN 6.8 EXCEPT AS DIRECTED IN ROTATIONAL CROP GUIDELINES

WEEDS CONTROLLED

Cocklebur	Morningglory sp.	Yellow nutsedge
Beggarticks	Entireleaf	
Florida beggarweed	Ivyleaf	
Pigweed sp	Pitted	
Palmer	Smallflower	
Redroot	Tall	
Smooth	Sicklepod	
Spiny amaranth	Smartweed	
Tall Waterhemp	Ladysthumb	
Mustard, wild	Pennsylvania	
Hemp sesbania	Ragweed
Jimsonweed	Common
	Giant



ROTATIONAL CROP GUIDELINES

The minimum length of time in months before " Classic" ppi or pre treated soils can be planted with major crops is given below. Choice of rotational crop is determined by geographic location (state), rate, soil pH and time interval elapsed since last application. Guidelines are divided into three groups based on these criteria.

GROUP 1

* States of AL, AR, GA, LA, MS, TN and MO Bootheel soil pH less than 6.8..

GROUP 2

* States of DE, KY, MD, NJ, PA, SC, VA and WV with soil pH of less than 6.8 and use rate of 2-4 oz per acre.

GROUP 3


- * All states not listed in Group 1 and 2 with soil pH less than 6.8.
- * States of AL, AR, FL, GA, LA, MS, TN, TX and MO Bootheel with soil pH 7.1-7.5
- * States of DE, KY, MD, NJ, NC, PA, SC, VA or WV
 - with soil pH 6.-7.5
 - fields not meeting rate requirements in Group 2

Minimum time in months before planting recrop:

<u>CROPS</u>	<u>GROUP1</u>	<u>GROUP2</u>	<u>GROUP3</u>
soybeans	anytime	anytime	anytime
barley	4	4	4
ryegrass	4	4	4
wheat	4	4	4
alfalfa	10	10	18
clover	9	10	18
field corn	9	10	18
cotton	10	10	18
peanuts	12	12	18
rice	10	10	18
sorghum	10	10	18
tobacco	10	10	18
(transplant)			
Tomato	10	10	18
(transplant)			

All other crops require a minimum of 18 month rotational interval and a successful field bioassay.**

The suggested interval before starting a field bioassay is 10 months for Group 1 and 15 months for Group 2 and 3.



** The season before planting any crop not listed above, a successful field bioassay must be completed. The field bioassay will detect small quantities of "Classic" which can remain in the soil and injure rotational crops. A successful field bioassay means growing to maturity a test strip of the crop(s) intended for production the following year. The test strip should cross the whole field including knolls and low areas.

SPRAYER CLEANUP

To avoid subsequent injury to crops other than soybeans, immediately after spraying and prior to spraying other crops, thoroughly remove all traces of "Classic" from mixing and spraying equipment. For more information, consult supplemental bulletin on sprayer cleanout following the use of Du Pont "Classic", "Canopy", "Preview", "Lorox" Plus or "Gemini" Herbicide.

IMPORTANT


BEFORE USING "CLASSIC", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABELS.

This bulletin contains new or supplemental instructions for use of this product which may not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.


[1] Registered trademark of E. I. du Pont de Nemours & Co. (Inc.)

E. I. du Pont de Nemours and Company
Agricultural Products Department
Wilmington, Delaware 19880-0038



"CLASSIC" HERBICIDE
EPA REG. NO. 352-436

SUPPLEMENTAL LABELING
"CLASSIC" HERBICIDE FOR
NONCROP USAGE



Directions for Use

It is a violation of federal law to use this product
in a manner inconsistent with its labeling.

Du Pont "Classic" Herbicide is recommended for postemergence usage to control certain annual weeds on noncrop sites such as fence rows, roadsides, equipment storage areas, and other similar areas.

HOW TO USE

Velvetleaf, Pigweeds (and other annuals)

Apply 1 to 2 ounces "Classic" per acre plus surfactant to seedling weeds.

TANK MIXTURES

"Classic" may be tank mixed with suitable registered herbicides to control weeds other than those listed. Read and follow all manufacturer's label recommendations for the companion herbicide.

APPLICATION

GROUND APPLICATION

For optimum spray distribution and thorough coverage, use flat fan nozzles. Do not use less than 10 gallon spray volume per acre (GPA).

SURFACTANT

Add a nonionic surfactant of at least 80% active ingredient strength at 0.25% vol/vol (1 quart per 100 gallons of water) to the spray solution.

923 77 200 COPY

injury to plants other than soybeans will likely occur. When spraying close to adjacent crops or plants, do everything possible to reduce the occurrence of off-target spray movement such as:

- Increase spray volume.
- Use lower sprayer pressure (but not less than 20 psi).
- Add an appropriate drift control agent.
- Stop spraying if wind speed becomes excessive.

Use the spray preparation of "Classic" within 24 hours as product degradation may occur. If spray preparation is left standing, thoroughly agitate before reusing.

Do not graze or feed forage or hay from treated areas to livestock.

IMPORTANT

BEFORE USING "CLASSIC", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on the package label. Follow instructions carefully.

This labeling must be in the possession of the user at the time of application.

.....
: [1] Registered trademark of E. I. du Pont de Nemours & Co. (Inc.)

.....
: E. I. Du Pont de Nemours & Co., Inc.

.....
: Agricultural Products Department

.....
: Wilmington, Delaware 19880-0038

