Unacreased Act, as Americal for the penticular and an american and american and an american and american and an american and an american and american an

"CLASSIC" HERBICIDE EPA REG. NO. 352-436

SUPPLEMENTAL LABELING
"CLASSIC" HERBICIDE
FOR CONTROL OF FLORIDA BEGGARWEED
IN PEANUTS

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Du Pont "Classic" is recommended for use for control of Florida Beggarweed in peanuts.

HOW TO USE

Make a single postemergence application of 1/2 ounce "Classic" per acre by ground or air for the control of actively growing Florida Beggarweed. Apply before Florida Beggarweed has begun to bloom and before it has reached 10 inches in height.

Florida Beggarweed exceeding 10 inches in height, or blooming will be only partially controlled (suppressed).

"Classic" can be applied from 60 days after rop emergence to within 45 days of harvest. Where peanut stands are erratic, or have been replanted, do not apply "Classic" until 60 days after the youngest peanuts have emerged. Do not make more than one application per season.

A surfactant mus* be included in the spray solution at 1 quart per 100 gallons of water (.25% volume per volume). Use only EPA approved surfactants of at least 80% active ingredient. Do not use a crop oil concentrate, or vegetable oil as crop injury will result.

Prior to using "Classic" Herbicide, consideration should be given to crop rotation plans. Consult Section I of the Rotational Crop Guidelines on the "Classic" product label.

Note: "Classic" may cause stunting of peanut growth. Test data have shown no adverse affects on yields from this interruption of peahut plant growth.



Do not graze treated fields or harvest for forage or hay.

Do not apply to peanuts under stress resulting from weather (drought), insects, previous herbicide injury, or disease (fungi or nematodes).

Research on peanut varieties indicates varietal tolerance to "Classic" applications may vary. The following runner type and Virginia type varieties have shown adequate tolerance:

Runher varieties: Florunner, Sunrunner and GK-7. Do not apply to Southern Runner as excessive crop injury may result.

Virginia type varieties: Florigiant, GK-3 and NC-7. Do not apply to early bunch varieties as excessive crop injury may result.

Florida Beggarweed more than 10 inches in height, blooming or regrowing as a result of mechanical injury (mowing or cultivation) may be only partially controlled (suppressed).

IMPORTANT

BEFORE USING "CLASSIC" HERBICIDE READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABELS.

This bulletin contains new or supplemental instructions for use of this product which may not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Do not apply this product through any type of irrigation system.

- [1] Registered trademark of E. I. du Pont de Nemours & Co. (Inc.)
- E. I. du Pont de Nemours and Company Agricultural Products Department Wilmington, Delaware 19898

060589