




Reg # 352-436

PM-25

192

SUPPLEMENTAL LABELING

AGRICULTURAL PRODUCTS

"..... A Growing Partnership With Nature"

**CLASSIC® HERBICIDE
FOR CONTROL OF FLORIDA
BEGGARWEED ON SOUTHERN
RUNNER VARIETY PEANUTS**

CLASSIC® HERBICIDE

EPA Reg. No. 352-436

FOR CONTROL OF FLORIDA BEGGARWEED ON SOUTHERN RUNNER VARIETY PEANUTS**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with the labeling.

Do not apply this product through any type of irrigation system.

HOW TO USE

Du Pont "Classic" Herbicide is recommended for use on "Southern Runner" variety of peanuts at the rate of 1/2 ounce per acre for the control of Florida beggarweed.

Make a single postemergence application of 1/2 ounce per acre of "Classic" by air or ground for control of actively growing Florida beggarweed. Apply before Florida beggarweed has begun to bloom and prior to exceeding 10 inches in height. Applications made to Florida beggarweed in bloom and/or taller than 10 inches may result in only partial control (suppression).

"Classic" can be applied from 60 days after crop emergence to within 45 days of harvest. Where peanut emergence is erratic, or stands have been replanted, do not apply "Classic" until 60 days after the youngest peanuts have emerged. Do not make more than 1 application per season.

A surfactant must be included in the spray solution at 1 quart per 100 gallons of water (0.25% v/v) of at least 80% active ingredient. Do not use a crop oil concentrate, or vegetable oil as crop injury will result. Do not use "Classic" with the surfactant "LI-700" as poor control of Florida beggarweed will result.

Refer to the "Classic" label for use directions, use restrictions, rotational considerations, and precautions.

NOTE: "Classic" may cause reduction in peanut growth. Test data on Southern Runner variety have shown no adverse effects on yields from this temporary reduction in vine length.

RESTRICTIONS

Do not apply within 45 days of harvest.

Do not graze treated fields or harvest for hay.

Do not apply "Classic" to Southern Runner variety peanuts growing under stress resulting from weather (drought), insects, nematodes, previous herbicide injury, or disease.

DO NOT APPLY "CLASSIC" TO SOUTHERN RUNNER VARIETY PEANUTS IN A TANK MIX WITH 2,4-DB

DO NOT APPLY "CLASSIC" TO SOUTHERN RUNNER VARIETY PEANUTS IN TANK MIXES WITH SULFUR OR PRODUCTS CONTAINING SULFUR AS EXCESSIVE CROP INJURY CAN RESULT.

Florida beggarweed more than 10 inches tall in height, blooming, or regrowing as a result of mechanical injury (mowing or cultivation) may be only partially controlled (suppressed).

IMPORTANT - BEFORE USING "CLASSIC", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of application.

1 Registered trademark of Loveland Industries, Inc.

D-021392

BEST AVAILABLE COPY

NOTIFICATION
REVISED
10-4

MAP
1992



AGRICULTURAL PRODUCTS

"..... A Growing Partnership With Nature"



SUPPLEMENTAL LABELING

TRI-PACK

**LEXONE® DF HERBICIDE PLUS
"TREFLAN" E.C. HERBICIDE
FOLLOWED BY CLASSIC®
HERBICIDE FOR WEED CONTROL
IN SOYBEANS**

LEXONE® DF HERBICIDE

EPA Reg. No. 352-390

CLASSIC® HERBICIDE

EPA Reg. No. 352-436

"LEXONE" DF HERBICIDE PLUS "TREFLAN" E.C. HERBICIDE FOLLOWED BY "CLASSIC" HERBICIDE FOR WEED CONTROL IN SOYBEANS

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Du Pont "Lexone" DF Herbicide is recommended for preplant incorporated application in a tank mix with "Treflan" E.C. or as a preemergence overlay application following a preplant incorporated application of "Treflan" E.C. Herbicide. Apply Du Pont "Classic" Herbicide as a postemergence application for additional control of broadleaf weeds.

HOW TO USE

PREPLANT INCORPORATED OR PREEMERGENCE APPLICATIONS

Apply "Lexone" DF plus "Treflan" E.C. as a preplant incorporated tank mix application OR apply "Lexone" DF as a preemergence treatment following a preplant incorporated application of "Treflan" E.C. Select rates, according to soil type, from the chart below.

Broadcast Rates Per Acre		
Soil Texture	"Lexone" DF	"Treflan" E.C.
Coarse	1 1/4 to 1 3/4 lb*	1 pt
Medium	1 3/4 to 2 lb*	1 1/2 pt
Fine	2 to 2 3/4 lb*	2 pt
*Select the higher rate for soils with organic matter greater than 3.0% or for fields with heavy broadleaf weed pressure.		

Refer to the "Lexone" DF and "Treflan" E.C. labels for further information on use instructions, weeds controlled, restrictions and precautions. The most restrictive labeling will apply.

Postemergence applications

For additional control of broadleaf weeds, apply "Classic" Herbicide at a rate of 1/2 to 3/4 ounces per acre as a postemergence when weeds are small and actively growing (usually 14-28 days after preplant incorporated or preemergence application of "Lexone" DF). Always include a nonionic surfactant or crop oil concentrate as a spray adjuvant. Refer to the "Classic" Herbicide label for information on use instructions, rates, weeds controlled, weed size, restrictions, precautions, and rotational cropping intervals.

IMPORTANT

BEFORE USING "LEXONE" DF, "TREFLAN" E.C., AND "CLASSIC" HERBICIDES, READ AND CAREFULLY NOTE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABELS.

This bulletin contains new or supplemental instructions for use of these products which may not appear on the package labels. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Registered trademark of DowElanco, Indianapolis, IN

120491

02-13-90

BEST AVAILABLE COPY

NOTIFICATION
DATE 1/13/93
1993