



**DuPont Agricultural  
Products**

RECD L.A./OPP/DPD

'97 MAR 10



"..... A Growing Partnership With Nature"

PM 25 352-436 **SUPPLEMENTAL LABELING** 10/2

**CLASSIC® HERBICIDE  
PINNACLE® HERBICIDE  
TANK MIXES WITH PURSUIT® OR  
PURSUIT ECO PAK® HERBICIDE  
FOR CONTROL OF NIGHTSHADE  
IN SOYBEANS IN CERTAIN STATES**

**CLASSIC® HERBICIDE**

Reg. No. 352-436

**PINNACLE® HERBICIDE**

EPA Reg. No. 352-525

**"CLASSIC", "PINNACLE" AND "CLASSIC" PLUS "PINNACLE" HERBICIDES TANK MIX APPLICATION  
WITH REDUCED RATES OF PURSUIT OR PURSUIT ECO PAK HERBICIDE  
FOR CONTROL OF NIGHTSHADE IN SOYBEANS IN THE STATES OF INDIANA, IOWA, MICHIGAN,  
MINNESOTA, OHIO, PENNSYLVANIA, SOUTH DAKOTA AND WISCONSIN**

Either DuPont "Classic", "Pinnacle" or "Classic" plus "Pinnacle" Herbicides may be applied in a tank mix with a reduced rate of "Pursuit" or "Pursuit Eco Pak" for the control of nightshade in addition to those weeds listed on the "Classic" or "Pinnacle" labels in the states of Indiana, Iowa, Michigan, Minnesota, Ohio, Pennsylvania, South Dakota and Wisconsin.

**ADJUVANTS:** Postemergence applications of either "Classic", "Pinnacle" or "Classic" plus "Pinnacle" tank mixed with either "Pursuit" or "Pursuit Eco Pak" must include the addition of a nonionic surfactant and ammonium nitrogen fertilizer.

- Use a nonionic surfactant at the rate of 1 pint per 100 gal of solution (0.125% v/v). Under dry, cool (generally 70° F or less) conditions the rate of nonionic surfactant may be increased to 2 pints per 100 gal. of solution (0.25% v/v).
- Use a high quality nitrogen fertilizer product such as 28-0-0 at a rate of 4 - 8 pints per acre, or 10-34-0 at a rate of 2 - 4 pints per acre. Alternately, a high-quality, sprayable grade of ammonium sulfate (21-0-0) may be used at a rate of 2 - 4 pounds per acre. Use the lower rate for spray volumes less than 15 gal/ac.
- Do not use "Dash", "Dash" HC<sup>2</sup>, crop oil concentrates or methylated seed oil products such as "Sun-It" II<sup>3</sup> when tank mixing either "Classic", "Pinnacle", or "Classic" plus "Pinnacle" with either "Pursuit" or "Pursuit Eco Pak" as excessive crop injury may occur.

For additional information refer to the DuPont bulletin "Approved Adjuvants for Use With DuPont Row Crop and Cereal Herbicides" for a list of approved adjuvants and suggested use rates.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**HOW TO USE**

"Classic" at 1/4-3/4 oz/ac, "Pinnacle" at 1/4 oz/ac, or "Classic" at 1/4-1/3 oz/ac plus "Pinnacle" at 1/4 oz/ac may be tank mixed with either "Pursuit" at 2 oz/ac or with "Pursuit Eco Pak" at a rate of one soluble bag per 10 acres for postemergence control of weeds listed on the "Classic" and/or "Pinnacle" labels, and for the control of eastern black nightshade less than 2 inches tall. Refer to the "Classic" and "Pinnacle" labels for other weeds controlled and maximum heights.

Best results are obtained when either "Classic", "Pinnacle" or "Classic" plus "Pinnacle" are tank mixed with either "Pursuit" or "Pursuit Eco Pak" and applied to weeds that are young (after the first true leaves have expanded, but before they exceed the size indicated on this label) and actively growing. This is generally 21-30 days after planting of soybeans. Applications made to weeds that are in the cotyledon stage, or to weeds larger than the sizes indicated, or to weeds under stress (weather, herbicide, or other) may result in unsatisfactory control.

This program is recommended for the control of broadleaf weeds only. Other measures should be used to control grassy weeds.

**APPLICATION INFORMATION**

**Broadcast Application:** Use flat fan nozzles at 25 - 40 psi. Do not use flood, hollow cone, rain drop, whirl chamber or controlled droplet applicator (CDA) type nozzles as unacceptable crop injury, excessive spray drift, or poor weed control may result. Use 10 - 25 gallons of water per acre. For proper spray coverage, adjust the boom and nozzle height according to the specifications listed by the nozzle manufacturer.

© 1997 E. I. du Pont de Nemours and Company, DuPont Agricultural Products, Wilmington, Delaware 19898

20/2

**Band Application:** For band application, use proportionately less spray mixture. To avoid crop injury, carefully calibrate the band applicator so as not to exceed the desired use rate. Carefully follow the manufacturer's instructions for nozzle type, (flat fan preferred), nozzle orientation, distance of the nozzles from the crop and weeds, spray volumes, calibration, and spray pressure. For additional information on row banders, see DuPont bulletin "Application Accuracy - Row Banders."

**Aerial Application:** Use nozzle types and arrangements that will provide for optimum spray distribution and maximum coverage at 5 to 10 gallons per acre. Do not apply during a temperature inversion, when winds are gusty, or when other conditions will favor poor coverage and/or off target spray movement. Use a minimum of 5 gallons of water per acre. Consult the respective product labels for special directions for aerial application.

**IMPORTANT PRECAUTIONS**

- Soybeans should be free from stress and actively growing at the time of application. Stress may be caused by abnormally hot or cold weather, growing conditions such as drought or water-saturated soil, disease, soil nutrient deficiencies such as iron chlorosis, or injury from nematodes, insects, or prior herbicide applications.
- Applications of either "Classic", "Pinnacle", or "Classic" plus "Pinnacle" when tank mixed with either "Pursuit" or "Pursuit Eco Pak" may shorten stem internodal length and cause temporary crop injury. Crop response may be increased when applications are made to soybeans that are under stress. Soybeans will recover quickly under normal growing conditions.
- Cultivation may put weeds under stress by pruning roots, thus reducing weed control. Avoid cultivation 7-10 days prior to or following application of the herbicide treatment. For maximum weed control, cultivate 7-10 days after application.
- Apply this treatment after the first trifoliolate of the soybean has fully expanded and the plants are actively growing, but before soybeans begin to flower.
- Refer to the "Classic", "Pinnacle", "Pursuit" or "Pursuit Eco Pak" Herbicide labels for additional use directions, use restrictions, rotational crop intervals, and precautions. The most restrictive provision on either label will apply.
- Applications within 1 hour of rain may reduce weed control.

**IMPORTANT**

**BEFORE USING "CLASSIC", "PINNACLE", "PURSUIT" OR "PURSUIT ECO PAK" HERBICIDES, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA-REGISTERED LABELS.**


This bulletin contains new or supplemental instructions for use of these products which do not appear on the EPA-registered package labels. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

<sup>1</sup> Registered trademark of American Cyanamid Company

<sup>2</sup> Registered trademark of BASF AG.

<sup>3</sup> Registered trademark of Agsco, Inc.



102296