



Velpar[®] RP

HERBICIDE
Dispersible Liquid

Contains 2 lbs Active Ingredient per Gallon

ACTIVE INGREDIENT

Hexazinone [3-cyclohexyl-6-(dimethylamino)-1-methyl-1,3,5-triazine-2,4(1H,3H)-dione]..... 26%

INERT INGREDIENTS..... 74%

U.S. Pats 3,902,887 & 3,983,116

EPA Reg. No. 352-437

RECEIVED
DEC 1 1987
352-437

352-437
EPA Est. 4805-AR-1 42781-MS-1 34704-MS-1
Super used is first letter of batch number

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Du Pont will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by Du Pont. User assumes all risk associated with such non-recommended use.

Make application just before or during period of active growth of brush species to be controlled. Apply "Velpar" RP with an exact delivery handgun applicator. This equipment delivers a thin stream of predetermined volume when triggered. Direct the application to the soil within 3' of the root collar at the rate of 2 to 4 milliliters for each 1" of stem diameter of brush to be controlled, but not to exceed 1/3 gallon of "Velpar" RP per acre per season. When treating large stems, and when more than one shot of "Velpar" RP is needed per stem, make application on opposite sides of the stem. Do not treat more than 300 one-inch stem equivalents (approximately 70 bush plants) per acre per season.

NOTE: For effective brush control and prevention of damage to desirable vegetation; do not apply to brush standing in water; do not use water from treated ditches for irrigation; do not use where desirable trees or shrubs are immediately adjacent to the treated area.

Do not use on marshy or poorly drained sites nor on soils classified as clays. Do not cut hay from treated grasses.

Do not apply to snow covered or frozen ground.

Do not apply "Velpar" RP within three times the height or canopy diameter (whichever is greater) of desirable trees or other plants. Injury or death of desirable trees or plants may result if this buffer is not maintained.

Do not apply this product through any type of irrigation system.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

STORAGE AND DISPOSAL

STORAGE: Do not subject to temperatures below 32°F. Store product in original container only, away from other pesticides, fertilizer, food, or feed.

DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent) the container. Then offer for recycling, or reconditioning, or puncture and dispose of in a sanitary landfill, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Keep out of reach of children
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER! CAUSES EYE DAMAGE.

Do not get in eyes. Wear goggles or face shield and rubber gloves when handling. Avoid contact with skin and clothing. Avoid breathing spray mist. Harmful if swallowed.

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Get medical attention if skin irritation persists. Remove and wash contaminated clothing before re-use.

For medical emergencies involving this product, call toll free 1-800-441-3637.

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

Refer to supplemental labeling for use instructions to protect ENDANGERED SPECIES when used on rangeland and pastures.

PHYSICAL AND CHEMICAL HAZARDS

Flammable. Keep away from heat, sparks, and open flame. Keep container closed.

IMPORTANT

Injury or loss of desirable trees or other plants may result from failure to observe the following: Do not apply except as recommended or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift or spray to desirable plants. Keep from contact with fertilizers, insecticides, fungicides and seeds.

Thoroughly clean all traces of "Velpar" RP from application equipment immediately after use.

GENERAL INFORMATION

Du Pont "Velpar" RP Herbicide is a liquid product to be applied undiluted with an exact delivery handgun applicator as an individual stem treatment for control of undesirable woody plants on rangeland and pastures. It is non-corrosive to equipment.

At recommended rates, "Velpar" RP provides control of undesirable woody plants such as elm, hackberry, huisache, juniper spp. (red-berry, eastern red cedar), mesquite, myrtle, oaks, osage orange (bois d'arc), tallow, whitebrush and willow. On woody plants, symptoms usually appear within 3 weeks after sufficient rainfall has carried the herbicide into the root zone where it is absorbed during periods of active plant growth. Defoliation and subsequent refoliation may occur, but susceptible plants are killed. The effect of treatment will vary with the amount applied, soil type, rainfall, timing and uniformity of application and other conditions.

Net 1896 ml

©1987 E. I. DU PONT DE NEMOURS & CO. (INC.), AGRICULTURAL PRODUCTS DEPARTMENT, WILMINGTON, DELAWARE

AG-2467 1184/8016/8107

Made in U.S.A.

Printed in U.S.A.

205-
SPECIMEN LABEL


100% OF ACTUAL SIZE

SECOND PROOF



Velpar[®] RP

HERBICIDE



TRADEMARK

DISPERSIBLE
LIQUID

Net
1896 ml