



ESCORT® HERBICIDE

REGISTRATION CODE xxxx-000

PACKAGE SIZE
8 - 8 OZ. BOTTLES/CARTON

DRY FLOWABLE

ACTIVE INGREDIENT

Metsulfuron methyl

(Methyl 2-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]-
carbonyl]-amino]sulfonyl]benzoate).....

INERT INGREDIENTS.....

BY WEIGHT

80%

40%

TOTAL 100%

EPA Reg. No. 352-439

U.S. Patent 4,383,113

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! Harmful if absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT

In case of contact with eyes, immediately flush with plenty of water. If on skin, wash with plenty of soap and water. Get medical attention if irritation persists. For medical emergencies involving this product, call toll free 1-800-441-3637.

ENVIRONMENTAL HAZARDS

Do not apply to open water nor to swamps, bogs, marshes or potholes while water is present. Do not contaminate water when disposing of equipment washwaters.

This herbicide is injurious to plants at extremely low concentrations. Nontarget plants may be adversely effected from drift and run-off.

IMPORTANT

DO NOT USE ON FOOD OR FEED CROPS. Injury to or loss of desirable trees or other plants may result from failure to observe the following: Do not apply "Escort" Herbicide (except as recommended), or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water, including irrigation water. Keep from contact with fertilizers, insecticides, fungicides and seeds.

Following an "Escort" application, do not use sprayer for application to crops. This is extremely important, as low rates of "Escort" can kill or severely injure most crops (except small grains).

GENERAL INFORMATION

Do not use this product in California nor in the following counties of Colorado: Saguache, Rio Grande, Alamosa, Costilla and Conejos.

Du Pont "Escort" Herbicide is a dispersible granule to be mixed in water and applied as a foliar spray for control of unwanted yearly plants and annual and perennial broadleaf weeds, basal soil treatment of multiflora rose, for pre-emergence and postemergence industrial weed control, and for the growth suppression of certain desirable grasses in noncrop areas such as airports, highways, roadsides, utility and pipeline rights-of-ways, petroleum tank farms, storage areas, pumping installations, plant sites, industrial turf, roadside turf, railroad rights-of-way and fence rows.

Do not apply to open water (such as lakes, reservoirs, rivers, streams, creeks, salt water bays or estuaries) nor while water is present in fresh water wetlands (such as marshes, swamps, bogs or potholes) nor to salt water marshes within tidal areas nor to areas where the herbicide is likely to move into water (such as insides of ditches, steep banks along waterways or impervious substrates) nor to areas near desirable plants where roots of these plants may extend. It is permissible to treat the berm of ditches, seasonally dry flood plains, deltas, marshes, swamps, bogs and transitional areas between upland and lowland sites.

Warm, moist conditions following treatment promote the activity of "Escort", while cold, dry conditions may reduce or delay activity. Weeds and brush hardened off by cold weather or drought stress may not be controlled. Degree of control and duration of effect are dependent on the rate used, sensitivity and size of target species, as well as soil moisture and soil temperature.

Following use of "Escort" on some naturally-occurring biotypes, may not be effectively controlled. Biotypes are naturally-occurring individuals of the species which have a slightly different genetic makeup. Resistant biotypes may look exactly the same as susceptible biotypes but are able to survive a use rate several times higher than needed to control susceptible biotypes. To delay the occurrence of resistant biotypes, use "Escort" in tank mixes and/or sequential treatments with other registered herbicides effective on the same broadleaf weed species. Do not let weed escapes go to seed; time postemergence treatments before seed formation. Respray problem areas in a timely and effective manner using a herbicide with a different mode of action.

"Escort" is noncorrosive, nonflammable, nonvolatile and does not freeze.

Mode of Action

"Escort" rapidly stops growth of susceptible plant species; however, typical symptoms (discoloration) may not be noticeable for several weeks after application, depending on growing conditions and susceptibility. Weeds absorb "Escort" through both the foliage and roots resulting in visual effects progressing from growth inhibition to reddish-purple coloration, chlorosis, necrosis, vein discoloration and death of terminal. Initial effects are usually seen 2 or 3 weeks following application; however, the final effects on weeds are evident at about 4 to 6 weeks after application. On brush, ultimate effects occur during the season after treatment.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

"Escort" Herbicide should be used only in accordance with recommendations on this label or in separate published Du Pont recommendations available through local suppliers.

Du Pont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by Du Pont. User assumes all risks associated with such nonrecommended use.

Du Pont "Escort" Herbicide is recommended as a postemergence application for control of brush and certain perennial broadleaf weeds and preemergence and postemergence for weed control on noncropland. "Escort" is also recommended for weed control on established, unimproved, industrial turf and native perennial grasses. On fescue/bluegrass turf, "Escort" is recommended for growth suppression and seedhead inhibition.

For tank mixes, use the most restrictive limitations from labeling of both products.

Consult your chemical supplier, applicator, consultant, appropriate state agricultural extension service representative or your local Du Pont representatives for specific recommendations.

Non-crop (Industrial) Weed Control

"Escort" is recommended for use for general weed control on non-crop, industrial sites such as airports, military installations, fence rows, roadsides and associated rights-of-ways, lumberyards, petroleum tank farms, pipeline and utility rights-of-ways, pumping installations, railroads, storage areas, plant sites and other similar areas including governmental and private lands.

When applied at lower dosage rates (as listed below), "Escort" provides short term control of weeds listed; usage at higher rates increases length and degree of weed control. Combinations with other herbicides such as 2,4-D, dicamba, Karmex® DF, Hyvar® X and Krovar® I DF increases the spectrum of weeds controlled. Bare ground objectives can be achieved with higher rates of "Escort" plus residual-type companion herbicides.

In states of Montana, North Dakota and South Dakota, "Escort" is for use only on railroad rights-of-ways.

"Escort" may be applied preemergence or postemergence to the weeds, but for best results, apply postemergence to young, actively growing weeds (less than 2" tall or 2" across) at any time of the year, except when ground is frozen. For postemergence treatment add surfactant.

"Escort" should be used during seasons when rainfall normally occurs as moisture is required to move "Escort" into the root zone of weeds before they develop an established root system. Preemergence treatments control or suppress weeds through root uptake, while postemergence works through root uptake and foliar uptake. Under limited rainfall conditions, "Escort" may not provide satisfactory weed control.

Treated soils should be left undisturbed to reduce the potential for "Escort" movement by soil erosion due to wind or water. Injury to crops may occur when treated soil is blown or moved onto land used to produce crops.

Rate selection is based on weed species, weed size and soil texture. Use the higher rates on established plants and on fine textured soils, and lower rates on smaller weeds and coarse textured soils.

Since the degree of control and duration of effect will vary with the amount of chemical applied, soil texture, soil pH, soil organic matter, weed size, rainfall and other conditions, it is suggested that users limit their first use to small areas.

For broadcast spray, apply "Escort" in a minimum of 10 gallons of water per acre. Spray volumes of 20 to 40 gallons per acre and pressures of 25 to 35 psi are preferred. Use coarse spray nozzles.

Application may also be made with a hand gun using 100 to 300 gals water per acre (as needed for coverage of weed growth).

Use the following rates of "Escort" (including surfactant at 0.25% to 0.5% by volume) to control weeds listed:

1/3 to 1/2 ounce "Escort"/acre

Annual broomweed	Corn cockle	Plantain
Annual sowthistle	Cow cockle	Redroot pigweed
Aster	Crownvetch	Redstem filaree
Bahiagrass	Dandelion	Rough fleabane
Beebalm	Dogfennel	Shepherd's purse
Bittercress	False chamomile	Smallseed falseflax
Bitter sneezeweed	Fiddleneck tarweed	Smooth pigweed
Blackeyed-susan	Field pennycress	Sweet clover
Blue mustard	Flaxweed	Tansymustard
Bur buttercup	Goldenrod	Treacle mustard
Chicory	Gumweed	Tumble mustard
Clover	Henbit	Wild carrot
Cocklebur	Lambsquarters	Wild lettuce
Goldenrod	Marestail	Wild mustard
Common chickweed	Maximillion	Woolly croton
Common groundsel	sunflower	Wood sorrel
Common mullein	Miners lettuce	Yankee weed
Common purslane	Pennsylvania	
Common yarrow	smartweed	
Conical catchfly	Plains coreopsis	

Brush Control

Foliar applications should be made after brush is fully leafed until the beginning of fall leaf coloration. Complete coverage of all foliage and stems is required for brush control. Effectiveness may be reduced if rainfall occurs within 24 hours after application.

High Volume Foliar Ground Applications: Apply as a full coverage spray to foliage, stems and limbs using up to 400 gallons of total spray per acre depending on plant species, height and density of growth and on the type of spray equipment used. On tall, dense brush, it is often necessary to spray from both sides to obtain adequate coverage. Use the following amounts of "Escort" per 100 gallons of water (include surfactant at 1 quart/100 gallons) and spray to runoff for control of:

1/2 to 1 Ounce*

Ash	Elm	Snowberry
Aspen	Hawthorn	Thimbleberry
Blackberry	Multiflora rose	Wild roses
Cherry	Salmonberry	Willow

to 2 Ounces*

Ylder	Ocean spray	Red maple
Oaks	(Holodiscus)	Cottonwood
	Osage orange	

*Where mixed stands occur, use the higher rate for the most difficult-to-control species.

For broader spectrum of control, "Escort" may be tank mixed with Du Pont Krenite® S, "Arsenal" [2] triglopyr or glyphosate using labeled use rates for each product.

"Escort" at 1/2 to 1 oz plus "Krenite" S at 1 1/2 to 3 gals (plus surfactant) per 100 gals water provides improved control of black cherry and red maple; do not exceed 6 gals "Krenite" S per acre.

Low Volume Foliar Ground Applications: Apply 1 to 3 ounces "Escort" per acre in 25 to 50 gals of water plus 0.25% surfactant by volume for control of ash, cherry, elm, winged elm, black locust, red oak, tuliptree and blackberry.

For improved control of black cherry and red maple, apply 1/2 to 2 ozs "Escort" plus 1-1/2 to 3 gals "Krenite" S (plus surfactant) per acre.

Spotgun Basal Soil Treatment: For control of multiflora rose, prepare a spray suspension of "Escort" using 1 ounce product per gallon of water. Mix vigorously until "Escort" is dispersed and agitate periodically while applying the spray suspension.

Apply the spray preparation with an exact-delivery hand gun applicator. This equipment delivers a thin stream of a predetermined volume when triggered. Apply at the rate of 4 milliliters for each 2 feet of rose canopy diameter. Direct the treatment to the soil within 2 feet of the stem union of the rose to be controlled. When treating large plants and more than one delivery of "Escort" dispersion is required, make applications on opposite sides of the plant.

Applications should be made from early spring to summer, when the ground is not frozen.

-Weed and Brush Control

CONIFER PLANTATIONS

Site Preparation

Loblolly Pine and Slash Pine Only

For control of blackberry and other broadleaf weeds (see listing under Non-crop Weed Control), apply 1/2 to 1-1/2 ozs "Escort" per acre (plus surfactant) by air or ground equipment. Treat blackberry plants after they have reached full leaf, but before leaf tissue has hardened in the fall. Use sufficient spray volume for complete coverage of these plants. For control of diffuse knapweed, Japanese honeysuckle, palmetto, black cherry and black locust, apply 3-1/3 ozs "Escort" per acre (plus surfactant).

For broader spectrum control, tank mix with:

- o "Arsenal" (as registered) for dewberry, greenbrier, honeysuckle, morning-glory, poison ivy, redvine, trumpet creeper, wild grape, sumac, beech, aspen, black gum, box elder, chinaberry, dogwood, hickory, persimmon, red maple, sassafras, sourwood, sweetgum, yellow poplar and others (see container label).
- o "Velpar" L (as registered) for bracken fern, ash, aspen, balsam, blackgum, cherry, poplar, birch, box elder, brambles, cherry (black and pin), elm, winged elm, hawthorne, hazel, honeysuckle, oaks (red, black and white), and willow (see container label).
- o "Accord" (or "Roundup" [3] both as registered) for blackgum, bracken fern, cherry, Virginia creeper, dewberry, honeysuckle, black locust, persimmon, poison ivy, sassafras, sourwood, sumac, sweet gum, hawthorn, trumpet creeper, wax myrtle and others (see container label).
- o "Oust" at 2 to 3 ozs per acre (loblolly pine only) for many herbaceous annual broadleaves and grasses, such as, fireweed, horseweed, common ragweed, annual sowthistle, black mustard, bur clover, sweet clover, curly dock, barnyardgrass, panicums, reed canarygrass, pokeweed, signalgrass and others (see container label). Apply to seedling weeds.

Douglas Fir

For control of blackberry and other broadleaf weeds, apply 1/2 to 1 oz "Escort" per acre (plus surfactant) in the spring. Douglas fir may be transplanted in treated areas the next season.

For broader spectrum control, "Escort" should be tank-mixed with "Velpar" L Herbicide. Use "Velpar" L as directed on the package for ground equipment.

RELEASE

Loblolly Pine (established plantations)

For control of blackberry, apply 1/2 to 1-1/2 ozs "Escort" per acre in a minimum of 10 gals water per acre by air or ground equipment, plus surfactant. Treat plants when they have reached full leaf but before leaf tissue has hardened in the fall. Treatment will also control many annual broadleaf weeds (see Weed Control); best results occur when annuals are treated in the seedling stage.

For broader spectrum control, "Escort" should be tank-mixed as follows:

"Arsenal" - Apply 1/2 to 1-1/2 ozs "Escort" plus 1 pt "Arsenal" (plus surfactant) per acre. Apply during summer to late fall (preferred) as described on the "Arsenal" container label. The combination controls or

improves control of blackberry, black gum, elm, winged elm, and cherry and controls many weeds as shown on product labels.

"Velpar" L (or "Velpar") - For brush control and herbaceous weed control (see container labels for species controlled), apply 1/2 to 1-1/2 ozs "Escort" plus 1-1/2 to 6 qts "Velpar" L (or equivalent "Velpar") per acre. Do not use surfactant.

"Oust" at 2 to 3 ozs per acre for herbaceous weed control. Do not use surfactant. Apply late winter through spring after the soil has settled around the pines.

Do not apply where conifers are suffering from loss of vigor caused by insects, disease, drought, winter damage, or other stresses, as injury may result. Excessive injury may result to conifers less than 4 years of age from transplanting on coarse-textured soils, and less than 3 years of age from transplanting on medium and fine-textured soils. Use on gravelly or rocky soils and exposed subsoils may cause injury or mortality to conifers. Poor weed and brush control may occur from application made when the soil is saturated with water and rain is imminent within 24 hours. Do not use on poorly drained or marshy sites but it may be used where pines are planted on beds. A foam-reducing agent may be added at the recommended rate, if needed.

Apply as a full coverage spray to foliage and stems. Total spray volume per acre will depend upon plant height and density of growth and the type of equipment used. See "Recommended Minimum Spray Volumes."

Loblolly pine may be transplanted into treated areas 4 months after application; slash pine the following season, at least 6 months after application; Douglas fir the following season after treatment of "Escort".

For application of "Escort" alone or in combination on loblolly pine and slash pine, follow directions on the package label of the companion product in addition to "Escort" directions above; see labels for additional plants controlled. Follow all restrictions on labeling of both products in the tank mixes.

Livestock may graze treated areas 0 days after application of up to and including 3/4 oz "Escort" per acre. For combinations, use the most restrictive recommendation of products used.

RECOMMENDED MINIMUM SPRAY VOLUMES			
PRODUCT SITE PREPARATION FOR LOBLOLLY AND SLASH PINES	SPRAY GROUND	VOLUME - GALS/ACRE AERIAL - HELICOPTER ONLY*	RATE NONIONIC SURFACTANT QTS/100 GALS
"Escort"	10	10	1
"Escort" + "Arsenal"	15	10	1
"Escort" + "Velpar" L	25	10	1
"Escort" + "Accord" or Roundup	10	10	2
"Escort" + "Oust"	15	10	1
PRODUCT SITE PREPARATION FOR DOUGLAS FIR	SPRAY GROUND	VOLUME - GALS/ACRE AERIAL - HELICOPTER ONLY*	RATE NONIONIC SURFACTANT QTS/100 GALS
"Escort"	10	10	1
"Escort" + "Velpar" L	25	10	1
PRODUCT RELEASE FOR LOBLOLLY PINE	SPRAY GROUND	VOLUME - GALS/ACRE AERIAL - HELICOPTER ONLY*	RATE NONIONIC SURFACTANT QTS/100 GALS
"Escort"	10	10 Pines established 1 yr.	1
"Escort" + "Arsenal"	15	10 Pines established 2 yrs.	1
"Escort" + "Oust"	10	10 Pines established 1 yr.	0
"Escort" + "Velpar" L	25	10 Pines established 1 yr.	0
* Do not apply by air (helicopter) within 200 feet of any homestead, agricultural land or other desirable plantings. Extreme care must be taken to prevent drift or runoff to homesteads, desirable plantings, agricultural land or any body of water. Do not apply when weather conditions favor drift from treated areas.			

USE PRECAUTIONS

Do not apply to any body of water, including streams, irrigation water or wells. Do not apply where runoff water may flow onto agricultural land, as injury to crops may result.

Do not apply "Escort" during periods of intense rainfall or to water-saturated soils as off-target movement may occur.

Do not apply to impervious substrates such as paved or highly compacted surfaces nor to frozen ground as off-target movement will occur.

Do not allow spray to drift onto adjacent crops or other desirable plants or trees as injury may occur.

Do not use on lawns, golf courses, athletic fields, commercial sod operations, or other high-maintenance, fine turfgrass areas.

Do not apply this product through any type of irrigation system.

In areas where sensitive crops are grown, make applications to soils whose surface has been settled by rain. Do not treat powdery dry soil or light sandy soils, when there is little likelihood of rainfall soon after treatment, as wind may cause off-target movement.

Following an "Escort" application, do not use sprayer for application to crops. This is extremely important, as low rates of "Escort" can kill or severely injure most crops except small grains).

SPRAY PREPARATION

Fill the spray tank half full of water and, with agitator running, add the proper amount of "Escort". Finish adding the required amount of water. Continuous agitation is required to keep the product in suspension. For postemergence applications, a nonionic surfactant of at least 80% active ingredient should be added (unless specified otherwise) to improve wetting and/or contact activity. To minimize drift, a drift control agent may be added at the recommended label rate. "Escort" spray preparations may degrade in acidic solutions below pH 7 if not used in 24 hours; they are stable in alkaline solutions. Thoroughly reagituate before using. Tank mixes with other registered herbicides should be tested for compatibility before full scale mixing.

Sprayer Agitation: Use mechanical or bypass agitation to thoroughly mix the spray solution. After initial mixing do not use excessive agitation when using a surfactant as foaming problems may result. To avoid foaming problems, use antifoam agents.

SPRAYER CLEANUP

Immediately after spraying, thoroughly remove all traces of "Escort" from mixing and spray equipment as follows:

1. Drain tank, rinse interior surfaces of tank; then flush tank, boom and hoses with clean water from a minimum of 5 minutes.
2. Partially fill the tank with clean water, then add cleaning solution*. Complete filling of the tank with clean water. Flush solution through boom, hoses, and nozzles, then allow to sit for 15 minutes with agitation running; then drain.
3. Repeat Step 2.
4. Repeat Step 1.
5. Nozzles, screens, and strainers should then be removed and cleaned separately.
6. To remove traces of cleaning solution, rinse the tank thoroughly with clean water and flush through hoses and boom.

7. Flush boom and hoses with clean water for 5 minutes just prior to using the sprayer for the first time after the "Escort" application.

* Use any of the following cleaning solutions:

1. One gallon ammonia (containing 3% active ingredient) per 100 gallons of water.
2. Nutra-sol[4] (carefully read and follow Nutra-sol label directions).
3. Loveland Spray Tank Cleaner[5] (carefully read and follow Loveland Spray Tank Cleaner label directions).
4. Tank-Aid[6] (carefully read and follow Tank-Aid label directions).

CAUTION: Do not use chlorine bleach with ammonia. All traces of liquid fertilizer containing ammonia, ammonium nitrate or ammonium sulphate must be rinsed with water from the mixing and application equipment before adding chlorine bleach solution. Failure to do so will release a gas with a musty chlorine odor which can cause eye, nose, throat and lung irritation. Do not clean equipment in an enclosed area.

STORAGE AND DISPOSAL

STORAGE: Store product in original container, away from other pesticides, fertilizer, food or feed.

PRODUCT DISPOSAL: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent) the container and then offer for recycling or reconditioning, or puncture and dispose of in a sanitary land fill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Injury to adjacent crops, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other material, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

- [1] Registered trademark of 3M Agricultural Products
- [2] Registered trademark of American Cyanamid Company.
- [3] Registered trademark of Monsanto Company
- [4] Sold by Thomas G. Kifol
- [5] Registered trademark of Loveland Tank Cleaner
- [6] Manufactured for Cornbelt Chemical Company


D - 071991

© 1991 E.I. DU PONT DE NEMOURS AND COMPANY, AGRICULTURAL PRODUCTS, WILMINGTON, DELAWARE 19096



AGRICULTURAL PRODUCTS

'..... A Growing Partnership With Nature'



SUPPLEMENTAL LABELING

ESCORT® HERBICIDE FOR PINE PLANTATIONS

ESCORT® HERBICIDE

EPA Reg. No. 352-439

SELECTIVE WEED CONTROL (RELEASE) IN LOBLOLLY PINE PLANTATIONS

DIRECTIONS

It is a violation of federal law to use this product in a manner inconsistent with the labeling.

Du Pont "Escort" Herbicide is recommended for control of competitive plants in loblolly pine plantations to improve tree growth (release).

HOW TO USE

For control of blackberry, apply 1/2 to 1-1/2 ozs. "Escort" per acre in a minimum of 10 gals. water per acre by air or ground equipment. Do not use surfactant on trees in the field for 1 year or less. For older trees, add 1 qt. surfactant (80% active) per 100 gals. water. Treat when blackberry has reached full leaf but before leaf tissue has hardened in the fall. Treatment will also control many annual broadleaf weeds; best results occur when annuals are treated in the seedling stage.

For broader spectrum control, "Escort" should be tank-mixed as follows:

"Arsenal" - Apply 3/4 to 1-1/2 ozs. "Escort" plus "Arsenal" Applicators Concentrate (at rates on the container label). Do not use surfactant on trees in the field for 1 year or less; on older trees, add surfactant as above. Apply during summer to late fall (preferred) as described on the "Arsenal" container label. This combination controls or improves control of blackberry, black gum, elm, winged elm, and cherry and controls many weeds as shown on product labels.

Du Pont Velpar® L (or "Velpar") - For brush control and herbaceous weed control (see container labels for species controlled), apply 1/2 to 1-1/2 ozs. "Escort" plus 1-1/2 to 6 qts. "Velpar" L (or equivalent "Velpar") per acre. Do not use surfactant. Excessive injury may result to conifers less than 4 years of age from transplanting on coarse-textured soils, and less than 3 years of age from transplanting on medium and fine-textured soils. Use on gravelly or rocky soils and exposed subsoils may cause injury or mortality to conifers. Poor weed and brush control may occur from application made when the soil is saturated with water and rain is imminent within 24 hours. Do not use on poorly drained or marshy sites, but it may be used where pines are planted on beds. A foam-reducing agent may be added at the recommended rate, if needed.

NOTE: For application of "Escort" alone or in combination on loblolly pine, follow directions on the package label of the companion product in addition to "Escort" directions above; see labels for additional plants controlled. Follow all restrictions on labeling of both products in the tank mixes.

Apply as a full coverage spray to foliage and stems. Total spray volume per acre will depend upon plant height and density of growth and the type of equipment used.

Do not apply where conifers are suffering from loss of vigor caused by insects, disease, drought, winter damage, or other stresses, as injury may result.

Livestock may graze treated areas 0 days after application of up to and including 3/4 oz. "Escort" per acre. For combinations, use the most restrictive recommendation of products used.

USE PRECAUTIONS

Do not apply to any body of water, including streams, irrigation water or wells.

Do not apply where runoff water may flow onto agricultural land, as injury to crops may result.

Do not apply "Escort" during periods of intense rainfall or to water-saturated soils as off-target movement may occur.

Do not apply to impervious substrates such as paved or highly compacted surfaces nor to frozen ground as off-target movement will occur.

Do not allow spray to drift onto adjacent crops or other desirable plants or trees as injury may occur.

Do not apply this product through any type of irrigation system.

In areas where sensitive crops are grown, make applications to soils whose surface has been settled by rain. Do not treat powdery dry soil or light sandy soils, when there is little likelihood of rain, all soon after treatment, as wind may cause off-target movement.

Following an "Escort" application, do not use sprayer for application to crops. This is extremely important, as low rates of "Escort" can kill or severely injure most crops except small grains).

BEST AVAILABLE COPY

828

IMPORTANT

BEFORE USING "ESCORT", "ARSENAL" OR "VELPAR" L, READ AND CAREFULLY NOTE THE CAUTIONARY STATEMENTS AND OTHER INFORMATION APPEARING ON THE PRODUCT CONTAINER LABELS.

This bulletin contains new or supplemental instructions for use of these products in combination which does not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

*Registered trademark of American Cyanamid

D-1029/91

