

herbicide
ESCORT

ACCEPTED

DEC 15 1999

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 352-439

PM 25

352-439

12-15-99

1 of 2

DUPONT Escort[®] herbicide

ACTIVE INGREDIENT: BY WEIGHT

Metsulfuron methyl
{Methyl 2-[[(4-methoxy-6-methyl-
1,3,5-triazin-2-yl)amino]carbonyl]-
amino]sulfonyl}benzoate} 60%

INERT INGREDIENTS 40%

EPA Reg. No. 352-439 **TOTAL 100%**

EPA Est. 39926-PR-1

**KEEP OUT OF
REACH OF CHILDREN
CAUTION**

See Additional Precautionary Statements
and Directions for Use in Supplemental
Labeling Enclosed.

Agricultural Use Requirements


Use this product only in accordance with
its labeling and with the Worker Protec-
tion Standard, 40 CFR part 170. Refer to
supplemental labeling under "Agricultural
Use Requirements" in the Directions for
Use section for information about this
standard.

Notice to Buyer: Purchase of this material
does not confer any rights under patents of
countries outside of the United States.

Notice: Please read the entire label, includ-
ing the supplemental labeling enclosed.
Before buying or using this product, read
the Limitation of Warranty and Liability in
the supplemental labeling. If the terms are
not acceptable, return the product at once,
unopened, for a refund of the purchase
price.

Produced by
DuPont Agricultural Caribe Industries, LTD for
E. I. du Pont de Nemours and Company,
Agricultural Products, Wilmington, DE 19888
A00444200 Made in Puerto Rico Printed in U.S.A.

herbicide



ACTIVE INGREDIENT: BY WEIGHT

Metsulfuron methyl
{Methyl 2-[[(4-methoxy-6-methyl-
1,3,5-triazin-2-yl)amino]carbonyl]-
amino]sulfonyl}benzoate} 60%

INERT INGREDIENTS 40%

TOTAL 100%

**KEEP OUT OF
REACH OF CHILDREN
CAUTION**

Dry Flowable

Net 1 oz


Refer to back panel for additional
information.

2af2

SUPPLEMENTAL LABELING

**DuPont Agricultural
Products**

"..... A Growing Partnership With Nature"



**ESCORT® HERBICIDE
SPOT TREATMENT
APPLICATION IN RANGE
AND NONCROP AREAS**

ESCORT® HERBICIDE

EPA Reg. No. 352-439

APPLICATION AS A SPOT TREATMENT TECHNIQUE IN RANGE AND NONCROP AREAS

DIRECTIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont ESCORT® herbicide is recommended for use as spot treatment to control noxious and troublesome weeds on range-land, noncrop areas such as roadsides and industrial sites including government and private lands.

APPLICATION INFORMATION

ESCORT® is recommended to control many species of weeds, including noxious weeds, in certain established grasses growing on noncrop areas and forage grasses growing on range land. Refer to the "Weeds Controlled" section of the package label or supplemental labeling for a listing of susceptible weed species.

If the sprayer is calibrated, consult the package label or other supplemental labeling to select the application rate per acre of ESCORT® appropriate for the target weeds.

OR

Use the measuring guide enclosed with the ESCORT® 1 ounce container to mix one gram of ESCORT® per one gallon of water along with a suitable surfactant. Spray to the point of wetting the entire surface of the target weeds, approximately 40 gallons of solution per acre.

When applied in this manner there is no grazing restriction following the use of ESCORT®.

APPLICATION TIMING

Applications may be made at anytime of the year, except when the soil is frozen.

USE PRECAUTIONS

- Varieties and species of grasses differ in their tolerance to herbicides. When using ESCORT® on a particular grass for the first time, limit use to a single 1 ounce container. If no injury occurs throughout the season larger acreage may be treated.
- Applications made to some established grasses may cause temporary stunting, yellowing or seedhead suppression (i.e. fescue, timothy).
- Applications made to newly established grasses less than 2 years from seeding may result in injury or loss.
- Do not apply to forage grasses known to be sensitive to ESCORT® such as ryegrass (Italian and perennial), bahia or Garrison's creeping foxtail.
- Broadleaf forage species, such as alfalfa and clover, are highly sensitive to ESCORT® and will be severely injured or killed.
- When used as directed, there is no grazing restriction for use rates of 1 2/3 ounce per acre and less. At use rates of 1 2/3 to 3 1/3 ounce per acre forage grasses may be cut for hay, fodder or green forage and fed to livestock, including lactating animals, 3 days after treatment.

IMPORTANT

BEFORE USING ESCORT® READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

D-111599