



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

APR - 2 2003

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Mr. Jacob J. Vukich
E.I. DuPont de Nemours & Company
Stine-Haskell Research Center
P.O. Box 30
Newark, DE 19714

Dear Mr. Vukich:

Subject: DuPont Escort XP Herbicide (Add Statement to Non-Agricultural Use Requirements
Box and Other Changes)
EPA Registration No. 352-439
Your Application Dated January 16, 2003

The labeling referred to above submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended is acceptable provided you make the following change before you release the product for shipment.

-Delete the "Integrated Pest Management" section pending resolution of issues concerning this section.--

Please submit three (3) copies of your final printed labeling incorporating the above change before you release the product for shipment. Amended labeling supercedes all previously accepted ones. A stamped copy of labeling is enclosed for your records.

Sincerely,


James A. Tompkins
James A. Tompkins *for*
Product Manager 25
Herbicide Branch
Registration Division (7505C)



DuPontTM Escort[®] XP

herbicide

DRAFT LABEL



“..... A Growing Partnership With Nature”



DuPontTM Escort[®] XP

herbicide

Dry Flowable

<i>Active Ingredient</i>	<i>By Weight</i>
Metsulfuron methyl Methyl 2-[[[(4-methoxy-6-methyl- 1,3,5-triazin-2-yl)amino]- carbonyl]amino]sulfonyl]benzoate	60%
<i>Inert Ingredients</i>	40%
TOTAL	100%

EPA Reg. No. 352-439

ACCEPTED
with COMMENTS
In EPA Letter Dated:
APR 2 2003

Product of DuPont Company
as registered under EPA Reg. No.
352-439

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! Causes eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

Long-sleeved shirt and long pants.

Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This herbicide is injurious to plants at extremely low concentrations. Nontarget plants may be adversely effected from drift and run-off.

IMPORTANT

DO NOT USE ON FOOD OR FEED CROPS EXCEPT AS RECOMMENDED BY THIS LABEL OR SUPPLEMENTAL LABELING. Injury to or loss of desirable trees or other plants may result from failure to observe the following: Do not apply DuPont™ ESCORT® XP herbicide (except as recommended), or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water, including irrigation water. Keep from contact with fertilizers, insecticides, fungicides and seeds.

Following an ESCORT® XP application, do not use sprayer for application to crops. This is extremely important, as low rates of ESCORT® XP can kill or severely injure most crops (except small grains).

GENERAL INFORMATION

ESCORT® XP herbicide is a dispersible granule that is mixed in water and applied as a spray. ESCORT® XP controls many annual and perennial weeds and woody plants in non-crop areas and conifer plantations. ESCORT® XP may be used for general weed and brush control on industrial non-crop sites and for selective weed control in certain types of unimproved turf grasses on industrial sites and in native grasses. It can also be used for controlling and suppressing undesirable weeds and hardwoods in conifer plantations.

ESCORT® XP controls weeds and woody plants primarily by postemergent activity. Although ESCORT® XP has preemergence activity, best results are generally obtained when ESCORT® XP is applied to foliage after emergence or dormancy break. Except where noted, ESCORT® XP provides the best results when applied to young, actively growing weeds. The use rate depends upon the weed species and size at the time of application.

The degree and duration of control may depend on the following:

- weed spectrum and infestation intensity
- weed size at application
- environmental conditions at and following treatment
- soil pH, soil moisture, and soil organic matter

It is permissible to apply ESCORT® XP to floodplains where surface water is not present, terrestrial areas of deltas and low lying areas where water is drained but may be isolated in pockets due to uneven or unlevel conditions. ESCORT® XP is noncorrosive, nonflammable, nonvolatile and does not freeze.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

ESCORT® XP is absorbed primarily through the foliage of plants, and by the roots to a lesser degree. Plant cell division is generally inhibited in sensitive plants within a few hours following uptake. Two to 4 weeks after application, leaf growth slows followed by discoloration and tissue death. The final effect on annual weeds are evident about 4 to 6 weeks after application. The ultimate effect on perennial weeds and woody plants occurs in the growing

season following application.

Warm, moist conditions following treatment promote the activity of ESCORT® XP, while cold, dry conditions may reduce or delay activity. Weeds and brush hardened off by cold weather or drought stress may not be controlled.

The use of a surfactant is recommended to enhance the control of susceptible plants, except where noted. Apply at a minimum rate (concentration) of 1/4% volume/volume (1 qt. per 100 gal. of spray solution), or at the manufacturer's recommended rate. Use only EPA approved surfactants containing at least 80% active ingredient. Certain types of surfactants, such as those incorporating acetic acid (i.e. LI-700), may not be compatible with ESCORT® XP and may result in decreased performance. Certain surfactants may not be suitable for use on desirable plants, such as turf and conifers, listed on this label. Consult the surfactant manufacturer's label for appropriate uses.

Weed and brush control may be reduced if rainfall occurs soon after application.

RESISTANCE

Biotypes of certain weeds listed on this label are resistant to ESCORT® XP and other herbicides with the same mode of action, even at exaggerated application rates. Biotypes are naturally occurring individuals of a species that are identical in appearance but have slightly different genetic compositions; the mode of action of a herbicide is the chemical interaction that interrupts a biological process necessary for plant growth and development.

If weed control is unsatisfactory, it may be necessary to retreat problem areas using a product with a different mode of action, such as postemergence broadleaf and/or grass herbicides.

If resistant weed biotypes such as kochia, prickly lettuce, and Russian thistle are suspected or known to be present use a tank-mix partner with ESCORT® XP to help control these biotypes, or use a planned herbicide rotation program where other residual broadleaf herbicides having different modes of action are used.

INTEGRATED PEST MANAGEMENT

To better manage weed resistance when using ESCORT® XP, use a combination of tillage, and tank-mix partners or sequential herbicide applications that have a different mode of action than ESCORT® XP, to control escaped weeds. Do not let weed escapes go to seed.

Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative herbicide recommendations available in your area.

It is advisable to keep accurate records of pesticides applied to treated areas to help obtain information on the spread and dispersal of resistant biotypes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ESCORT® XP should be used only in accordance with recommendations on this label or in separately published DuPont recommendations.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not

specifically recommended by DuPont. User assumes all risks associated with such nonrecommended use. For tank mixes, use the most restrictive limitations from the labeling of the products being mixed. Use only those tank mix partners which are labeled for the appropriate use site. Do not apply more than 4 ounces of DuPont™ ESCORT® XP per acre per year.

Do not use on food or feed crops except as recommended by this label or supplemental labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency in your State responsible for pesticide regulation.

AGRICULTURAL USES

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to use of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls.

Shoes plus socks.

CONIFER PLANTATIONS

Application Information

ESCORT® XP is recommended to control many species of weeds and deciduous trees on sites where conifers are growing or are to be planted. Apply by ground equipment or by air (helicopter only). Refer to the "Weeds Controlled" and "Brush Species Controlled" for a listing of susceptible species.

Application Timing

Apply ESCORT® XP after weeds have emerged or after undesirable hardwoods have broken winter dormancy and have reached the point of full leaf expansion.

Conifer Site Preparation

—Application Before Transplanting

After consulting the "Weeds Controlled" and "Brush Species Controlled" tables apply the rates of ESCORT® XP recommended for the most difficult to control species on the site.

Southeast—Apply up to 4 oz per acre for loblolly and slash. Transplant the following planting season.

Northeast and Lake States—Apply up to 2 oz per acre for red pine. Transplant the following planting season.

West—Apply up to 2 oz per acre prior to planting Douglas Fir, Sitka Spruce, Western Red Cedar, Western Hemlock, Ponderosa Pine, and Grand Fir in the Coast Range and western slope of the Cascades in Oregon and Washington. These conifer species listed can be planted anytime after application. Other conifer species can be planted providing the user has prior experience indicating acceptable tolerance to ESCORT® XP soil residues. Without prior experience, it is recommended that other species be planted on a small scale to determine selectivity before large scale plantings are made as unacceptable injury may occur. DuPont will not assume responsibility for injury to any conifer species not listed on this label.

Tank Mix Combinations—

For broader spectrum control the following products are recommended in combination with ESCORT® XP.

"Accord"

Tank mix 1 to 2 ounces of ESCORT® XP with 2 to 10 quarts of "Accord" per acre. Refer to the product container for a list of species controlled.

"Arsenal" Applicator's Concentrate

Tank mix 1 to 2 ounces of ESCORT® XP with 10 to 24 fluid ounces of "Arsenal" Applicator's Concentrate per acre. Loblolly and slash pines may be transplanted the planting season following application. The combination controls ash, black gum, cherry, hawthorn, honeysuckle, hophornbeam, persimmon, oaks (red, white and water), sassafras, sweetgum, Vaccinium species, and suppresses blackberry, dogwood, elms, myrtle dahoon, hickories, and red maple.

"Accord" + "Arsenal" Applicators Concentrate

Tank mix 1/2 to 1 ounce of ESCORT® XP with 16 to 64 fluid ounces of "Accord" and 10 to 12 fluid ounces of "Arsenal" Applicator's Concentrate per acre. Slash and loblolly pines may be transplanted the planting season following application. The combination controls cherry, dogwood, elms, oaks (red and water), persimmon, sassafras, sweetgum and suppresses hickory.

DuPont™ VELPAR® L or DuPont™ VELPAR® DF

Tank mix 1 to 2 ounces of ESCORT® XP per acre with VELPAR® L or VELPAR® DF at the rates recommended on the container for various soil textures. Loblolly and slash pines may be transplanted the planting season following application. Refer to the product container for a list of species controlled.

DuPont™ OUST®

Tank mix 1/2 to 1 1/2 ounces of ESCORT® XP with 2 to 3 ounces of OUST® per acre for herbaceous weed control. Refer to the product container and the "Weeds Controlled" section of this label for a listing of the weeds controlled. Loblolly and slash pines may be transplanted the planting season following application.

Tank mix 2 ounces of ESCORT® XP with 3 ounces of OUST® per acre for herbaceous weed control and early spring suppression of bull thistle and Canada thistle in the Coast Range and western slope of the Cascade Mountains. Douglas fir may be transplanted at least 90 days following application.

Release—Hardwood Control and Suppression

DuPont™ ESCORT® XP is recommended for application over the top of established slash and loblolly pine to control the species listed in "Weeds Controlled" and "Brush Species Controlled" section of this label. Apply 1 to 4 ounces per acre to control the species indicated, including kudzu.

Tank Mix Combinations—

For broader spectrum control the following products are recommended in combination with ESCORT® XP.

"Arsenal" Applicator's Concentrate

Tank mix 1 to 2 ounces of ESCORT® XP with 8 to 16 fluid ounces of "Arsenal" Applicator's Concentrate per acre may be applied to loblolly pine. Refer to the "Arsenal" Applicator's Concentrate label regarding the use of surfactants and the appropriate application timing with respect the age and development stage of the pines. The combination controls ash, black gum, cherry, hawthorn, honeysuckle, hophornbeam, oaks (red, white and water), sassafras, sweetgum, Vaccinium species, and suppresses blackberry, dogwood, elms, myrtle dahoon, hickories, persimmon, and red maple.

DuPont™ VELPAR® L or DuPont™ VELPAR® DF

Tank mix 1 to 2 ounces of ESCORT® XP with VELPAR® L or VELPAR® DF at the rates recommended on the container for various soil textures. The combination may be applied to loblolly and slash pines.

Release—Herbaceous Weed Control

ESCORT® XP may be applied to transplanted loblolly and slash pine for the control of herbaceous competition. Consult the "Weeds Controlled" for a listing of the susceptible species and recommended application rates. Best results are obtained when ESCORT® XP is applied just before weed emergence until shortly after weed emergence.

Tank Mix Combinations—

For broader spectrum control the following products are recommended in combination with ESCORT® XP.

"Arsenal" Applicators Concentrate

Tank mix 1/2 to 1 ounce of ESCORT® XP with 4 fluid ounces of "Arsenal" Applicators Concentrate per acre. The tank mix may be used on loblolly pine.

DuPont™ OUST®

Tank mix 1/2 to 1 1/2 ounces of ESCORT® XP with 2 to 3 ounces of OUST® per acre. Best results are obtained when ESCORT® XP is applied just before weed emergence until shortly after weed emergence. The tank mix may be used on loblolly and slash pine.

VELPAR® L or VELPAR® DF

Tank mix 1/2 to 1 ounce of ESCORT® XP with VELPAR® L or VELPAR® DF at the rates recommended on the container for various soil textures. The combination may be applied to loblolly and slash pines.

IMPORTANT PRECAUTIONS

—CONIFER PLANTATIONS ONLY

- Applications of ESCORT® XP made to conifers that are suffering from loss of vigor caused by insects, diseases, drought, winter damage, animal damage, excessive soil moisture, planting shock, or other stresses may injure or kill the trees.

- Applications of ESCORT® XP made for herbaceous release should only be made after adequate rainfall has closed the planting slit and settled the soil around the roots following transplanting.
- Do not apply ESCORT® XP to conifers grown as ornamentals.
- ESCORT® XP applications may result in damage and mortality to other species of conifers when they are present on sites with those listed in the preceding recommendations for conifer plantations.

HARDWOOD PLANTATIONS

Application Information

ESCORT® XP is recommended at rates up to 2 ounces per acre for the control of many weed species on sites where yellow poplar is growing or is to be planted and on sites where red alder is to be planted. Apply by ground equipment or by air (helicopter only). Refer to the "Weeds Controlled" sections of this label for a listing of susceptible species.

Application Timing

ESCORT® XP may be applied as a site preparation treatment prior to planting red alder or yellow poplar. As a prior to planting, site preparation treatment for red alder, ESCORT® XP may be tank mixed with other herbicides labeled for this use.

ESCORT® XP may also be applied over-the-top of planted yellow poplar seedlings after the soil has settled around the root system but before the seedlings have broken dormancy (prior to bud break).

Release—Herbaceous Weed Control

ESCORT® XP may be applied to yellow poplar for the control of herbaceous competition. Consult the "Weeds Controlled" for a listing of the susceptible species and recommended application rates. Best results are obtained when ESCORT® XP is applied just before weed emergence until shortly after weed emergence.

Tank Mix Combinations—

Tank mix 1/2 ounce of ESCORT® XP with 4 to 6 pints of VELPAR® L as recommended on the package label for "RELEASE--HERBACEOUS WEED CONTROL" in pine plantations in the eastern U.S. Follow the VELPAR® L label recommendations regarding altering the application rate by soil texture.

IMPORTANT PRECAUTIONS

—HARDWOOD PLANTATIONS ONLY

- Application of VELPAR® L and ESCORT® XP made to yellow poplar that are suffering from loss of vigor caused by insects, disease, drought, winter damage, animal damage, excessive soil moisture, planting shock or other stresses may injure or kill the seedlings.
- Applications of ESCORT® XP made for release should only be made after adequate rainfall has closed the planting slit and settled the soil around the roots following transplanting.
- The use of surfactant is not recommended for applications made over the tops of trees.
- Careful consideration must be given by an experienced and knowledgeable forester to match the requirements of yellow poplar and/or red alder to the conditions of

the site. Treatment of yellow poplar and/or red alder planted on a site inadequate to meet its requirements may injure or kill the seedlings.

NON-AGRICULTURAL USES

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

Non-crop industrial weed control and selective weed control in turf (industrial, unimproved only) are not within the scope of the Worker Protection Standard.

WEEDS CONTROLLED

1/3 to 1/2 ounce per acre

Annual sowthistle	Goldenrod
Aster	Lambsquarters
Bahiagrass	Marestail
Beebalm	Maximillion sunflower
Bittercress	Miners lettuce
Bitter sneezeweed	Pennsylvania smartweed
Blackeyed-susan	Plains coreopsis
Blue mustard	Plantain
Bur buttercup	Redroot pigweed
Chicory	Redstem filaree
Clover	Rough fleabane
Cocklebur	Shepherd's purse
Common chickweed	Silky crazyweed (locoweed)
Common groundsel	Smallseed falseflax
Common purslane	Smooth pigweed
Common yarrow	Sweet clover
Conical catchfly	Tansymustard
Corn cockle	Treacle mustard
Cow cockle	Tumble mustard
Crown vetch	Wild carrot
Dandelion	Wild garlic
Dogfennel	Wild lettuce
False chamomile	Wild mustard
Fiddleneck tarweed	Woolly croton
Field pennycress	Wood sorrel
Flixweed	Yankeweed

1/2 to 1 ounce per acre

Blackberry	Honeysuckle
Black henbane	Multiflora rose and other wild roses
Broom snakeweed*	Musk thistle***
Buckhorn plantain	Plumeless thistle
Common crupina	Prostrate knotweed
Common sunflower	Rosering gaillardia
Curly dock	Seaside arrowgrass
Dewberry	Sericea lespedeza
Dyer's woad	Teasel
Gorse	Wild caraway
Halogeton	
Henbit	

1 to 2 ounces per acre

Bull thistle	Purple loosestrife
Common mullein	Scotch thistle
Common tansy	Scouringrush
Field bindweed**	Salsify
Gumweed	Snowberry
Houndstongue	St. Johnswort
Perennial pepperweed	Western salsify
Poison hemlock	Whitetop (hoary cress)

1 1/2 to 2 ounces per acre

Canada thistle**	Russian knapweed**
Dalmation toadflax**	Tall larkspur
Duncecap larkspur	Yellow toadflax**

3 to 4 ounces per acre

Kudzu

* Apply fall through spring.

** Suppression, which is a visual reduction in weed competition (reduced population or vigor) as compared to untreated areas. Apply as a full coverage spray for best performance.

*** Certain biotypes of musk thistle are more sensitive to DuPont™ ESCORT® XP and may be controlled with rates of 1/4 to 1/2 ounce per acre. Treatments of ESCORT® XP may be applied from rosette through bloom stages of development.

Tank Mix Combination

For broader spectrum control and for use on certain biotypes of broadleaf weeds which may be resistant to ESCORT® XP and herbicides with the same mode of action, the following tank mixes are recommended.

Dicamba + 2,4-D

Combine 1/2 to 1 ounce of ESCORT® XP with 8 fluid ounces of dicamba and 16 fluid ounces of 2,4-D for the control of kochia.

Combine 1/2 ounce of ESCORT® XP with 8 fluid ounces of dicamba and 16 fluid ounces of 2,4-D for the control of spotted knapweed.

Combine 1 ounce of ESCORT® XP with 8 fluid ounces of dicamba and 16 fluid ounces of 2,4-D for the suppression of rush skeletonweed.

NONCROP (INDUSTRIAL) SITES

Application Information

ESCORT® XP is recommended for use for general weed and brush control on non-crop, industrial sites such as airports, military installations, fence rows, roadsides and associated rights-of-way, petroleum tank farms, pipeline and utility rights-of-way, pumping stations, railroads, storage areas, plant sites and other similar areas including governmental and private lands. It is also recommended for the control of certain noxious and troublesome weeds. Consult the "Weeds Controlled" and "Brush Species Controlled" tables to determine the appropriate application rate.

ESCORT® XP may be applied in tank mixture with other herbicides labeled for use on non-crop sites. Fully read the labels and follow all directions and restrictions on each label.

Application Timing

For best results, DuPont™ ESCORT® XP should be applied postemergence to young, actively growing weeds.

Applications may be made at any time of the year, except when the ground is frozen.

GRASS REPLANT INTERVALS

Following an application of ESCORT® XP to non-crop areas, the treated sites may be replanted with various species of grasses at the intervals recommended below.

For soils with a pH of 7.5 or less observe the following replant intervals:

Species	ESCORT® XP Rate oz/a	Replant Interval (months)
Brome, Meadow	1/2--1	2
	1--2	3
Brome, Smooth	1/2--1	2
	1--2	4
Fescue, Alta	1/2--1	2
	1--2	4
Fescue, Red	1/2--1	2
	1--2	4
Fescue, Sheep	1/2--1	1
	1--2	4
Foxtail, Meadow	1/2--1	2
	1--2	4
Green Needlegrass	1/2--2	1
Orchardgrass	1/2--1	2
	1--2	4
Russian wildrye	1/2	1
	1	2
	2	3
Switchgrass	1/2--1	1
	1--2	3
Timothy	1/2--1	2
	1--2	4
Wheatgrass, Western	1/2--1	2
	1--2	3

For soils with a pH of 7.5 or greater observe the following replant intervals:

Species	ESCORT® XP Rate oz/a	Replant Interval (months)
Alkali Sacaton	1/2--1	1
	1--2	3
Bluestem, Big	1/2--2	3
Brome, Mountain	1/2--1	1
	1--2	2
Gamma, Blue	1/2--2	1
Gamma, Sideoats	1/2	2
	>1/2	>3
Switchgrass	1/2	2
	>1/2	>3
Wheatgrass, Thickspike	1/2--2	1
Wheatgrass, Western	1/2--1	2
	1--2	3

The recommended intervals are for applications made in the Spring to early Summer. Because ESCORT® XP degradation is slowed by cold or frozen soils, applications made the late Summer or Fall should consider the intervals as beginning in the Spring following treatment.

Testing has indicated that there is considerable variation in response among the species of grasses when seeded into areas treated with ESCORT® XP. If species other than those listed above are to be planted into areas treated with ESCORT® XP a field bioassay should be performed, or previous experience may be used, to determine the feasibility of replanting treated sites.

TURF, INDUSTRIAL (UNIMPROVED ONLY)**Application Information**

ESCORT® XP is recommended for selective weed control in unimproved industrial turf where certain grasses are well established and desired as ground cover. ESCORT® XP is also recommended for the control of certain noxious and troublesome weeds in turf.

In addition to conventional spray equipment, ESCORT® XP may also be applied with invert emulsion equipment. When using an invert emulsion, mix the prescribed rate of ESCORT® XP in the water phase.

Consult the "Weeds Controlled" table to determine which weeds will be controlled by the following recommendations.

Fescue and Bluegrass--

Apply 1/4 to 1/2 ounce of ESCORT® XP per acre.

Crested Wheatgrass and Smooth Brome--

Apply 1/4 to 1 ounce of ESCORT® XP per acre.

Bermudagrass--

Apply 1/4 to 2 ounces of ESCORT® XP per acre.

Application Timing

Applications may be made at anytime of the year, except when the soil is frozen.

When a spring application is made on fescue or bluegrass, a second application may be made during the summer after full seedhead maturation.

**Growth Suppression and Seedhead Inhibition
(Chemical Mowing)****Application Information**

ESCORT® XP is recommended for growth suppression and seedhead inhibition in well established fescue and bluegrass turf at the use rate of 1/4 to 1/2 ounce per acre.

Tank Mix Combination

ESCORT® XP may be tank mixed with "Embark" for improved performance in the regulation of growth and seedhead suppression. Tank mix 1/4 to 1/2 ounce of ESCORT® XP with 1/8 to 1/4 pint of "Embark".

Application Timing

Application may be made after at least 2 to 3 inches of new growth has emerged until the appearance of the seed stalk.

IMPORTANT PRECAUTIONS**—INDUSTRIAL TURF ONLY**

- An application of ESCORT® XP may cause temporary discoloration (chlorosis) of the grasses. Use the lower recommended rates for minimum discoloration.
- With fescue and bluegrass, sequential applications made during the same or consecutive growth periods (i.e. spring and fall) may result in excessive injury to turf.

- Excessive injury may result when DuPont™ ESCORT® XP is applied to turf that is under stress from drought, insects, disease, cold temperatures (winter injury) or poor fertility.
- ESCORT® XP is not recommended for use on bahiagrass.

NATIVE GRASSES

ESCORT® XP is recommended for weed control and suppression in the establishment and maintenance of native grasses. It may be used where blue grama, bluestems (big, little, plains, sand, ww spar) bromegrasses (meadow), buffalograss, green sprangletop, indiangrass, kleingrass, lovegrasses (atherstone, sand, weeping, wilman), orchardgrass, sideoats grama, switchgrass (blackwell), wheatgrass (bluebunch, intermediate, pubescent siberian, slender, streamband, tall, thickspike, western), and russian wildrye are established. It may also be applied over these species in the seedling stage, except for orchardgrass and russian wildrye.

Application Information

Apply ESCORT® XP at the rate of 1/10 ounce per acre for the control and suppression* of bur buttercup (testiculate), common purslane, common sunflower*, cutleaf eveningprimrose*, flixweed*, lambsquarters* (common and slimleaf), maretail*, pigweed (redroot and tumble), snow speedwell, tansymustard* and tumble mustard (Jim Hill mustard).

- * Suppression is a visual reduction in weed competition (reduced population or vigor) as compared to untreated areas. Degree of suppression will vary with the size of weed and environmental conditions following treatment.

Application Timing

For established grasses, apply when weeds are in the seedling stage.

For grasses in the seedling stage, apply preplant or preemergence where the soil (seed bed) has been cultivated.

BRUSH CONTROL

Application Information

ESCORT® XP is recommended for the control of undesirable brush growing in non-crop areas. Applications may be made by air, high volume ground application, low volume ground application and ultra-low volume ground application. Except as noted for multiflora rose, ESCORT® XP should be applied as a spray to the foliage.

The application volume required will vary with the height and density of the brush and the application equipment used. Generally, aerial applications will require 15 to 25 gallons of water per acre; high volume ground application will require 100 to 400 gallons of water per acre; low volume ground application will require 20 to 50 gallons of water per acre; and ultra-low volume ground application will require 10 to 20 gallons of water per acre.

Regardless of the application volume and equipment used, thorough coverage of the foliage is necessary to optimize results.

BRUSH SPECIES CONTROLLED

Species	High Volume	Broadcast
	ESCORT® XP Rate oz/100 gal	ESCORT® XP Rate oz/a
Ash	1--2	1--3
Aspen	1--2	1--3
Black locust	1--2	1--3
Blackberry	1--2	1--3
Camelthorn	1--2	1--3
Cherry	1--2	1--3
Cottonwood	1--2	2--3
Eastern red cedar	1--2	2--3
Elder	1--2	2--3
Elm	1--2	1--3
Firs	3	1--2
Hawthorn	1--2	1--3
Honeysuckle	1--2	1/2--1
Mulberry	1--2	2--3
Multiflora rose	1--2	1--3
Muscadine (wild grape)	1--2	2--3
Oaks	1--2	1--3
Ocean spray (Holodiscus)	1--2	2--3
Osage orange	1--2	2--3
Red maple	1--2	2--3
Salmonberry	1/2--1	1--3
Snowberry	1/2--1	1--3
Spruce (black and white)	3	2--3
Thimbleberry	1/2--1	1--3
Tulip tree	1/2--1	1--3
Wild roses	1/2--1	1--3
Willow	1/2--1	1--3

For low volume and ultra-low volume ground applications, mix 4 to 8 ounces of ESCORT® XP per 100 gallons of spray solution.

Application Timing

Make a foliar application of the recommended rate of ESCORT® XP during the period from full leaf expansion in the spring until the development of full fall coloration on deciduous species to be controlled. Coniferous species may be treated at anytime during the growing season.

Tank Mix Combinations—

“Accord”

After consulting the “Brush Species Controlled” table, tank mix the prescribed rate of ESCORT® XP with the rate of “Accord” indicated for the various application methods on the “Accord” label. Refer to the “Accord” label for list of species controlled.

“Arsenal” herbicide

Combine 1 to 2 ounces of ESCORT® XP with 1 to 4 pints of “Arsenal” herbicide per acre and apply as a broadcast spray. Aerial applications should use a minimum of 15 gallons per acre spray volume. In addition to species listed above controlled by ESCORT® XP, this combination controls black gum, hophornbeam, sassafras, sweetgum, Vaccinium species, dogwood, myrtle dahoon, hickories, and persimmon.

"Garlon" 3A or "Garlon" 4

After consulting the "Brush Species Controlled" table, tank mix the prescribed rate of DuPont™ ESCORT® XP with the rate of "Garlon" indicated for the various application methods on the "Garlon" label. Refer to the "Garlon" label for list of species controlled.

DuPont™ KRENITE® S

After consulting the "Brush Species Controlled" table, tank mix the prescribed rate of ESCORT® XP with the rate of KRENITE® S indicated for the various application methods on the KRENITE® S label. Refer to the KRENITE® S label for list of species controlled.

"Tordon" K

After consulting the "Brush Species Controlled" table, tank mix the prescribed rate of ESCORT® XP with the rate of "Tordon" K indicated for the various application methods on the "Tordon" K label. Refer to the "Tordon" K label for list of species controlled.

"Tordon" K + "Arsenal" herbicide

Combine 1 to 1 1/2 ounce of ESCORT® XP with 2 to 8 fluid ounces of "Arsenal" and 1 to 2 pints of "Tordon" K per 100 gallons of water. Apply as a high volume spray. The tank mix controls cherry, elms, box elder, maples, hackberry, redbud, ash, oaks (including shingle oak), black locust and sassafras.

*"Tordon" K is a restricted use pesticide.

Spotgun Basal Soil Treatment

For control of multiflora rose, prepare a spray suspension of ESCORT® XP by mixing 1 ounce per gallon of water. Mix vigorously until the ESCORT® XP is dispersed and agitate periodically while applying the spray suspension.

Apply the spray preparation with an exact delivery handgun applicator. Apply at the rate of 4 milliliters for each 2 feet of rose canopy diameter. Direct the treatment to the soil within 2 feet of the stem union. When treating large plants and more than one delivery is required, make applications on opposite sides of the plant.

Applications should be made from early spring to summer.

IMPORTANT PRECAUTIONS

—NON-CROP BRUSH ONLY

- When using tank mixtures of ESCORT® XP with companion herbicides, read and follow all use instructions, application rates, warnings and precautions appearing on the labels. Follow the most restrictive label instructions for each of the herbicides used.

SPRAY EQUIPMENT

Following an ESCORT® XP application, do not use the sprayer or mixing equipment for application to agricultural crops, except that it may be used to treat pasture, range and wheat. This is extremely important as low rates of ESCORT® XP can kill or severely injure most agricultural crops.

The selected sprayer should be equipped with an agitation system to keep ESCORT® XP suspended in the spray tank. Use a sufficient volume of water to thoroughly cover the foliage of undesirable weeds, generally 10 to 40 gallons per acre. Select a spray volume and delivery system that will deliver a uniform spray pattern. Be sure the sprayer is calibrated before use. Avoid overlapping and shut off spray

booms while starting, turning, slowing or stopping to avoid injury to desired plants.

Refer to the brush control section of this label for information unique to that particular use.

MIXING INSTRUCTIONS

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of ESCORT® XP.
3. Continue agitation until the ESCORT® XP is fully dispersed, at least 5 minutes.
4. Once the ESCORT® XP is fully dispersed, maintain agitation and continue filling tank with water. ESCORT® XP should be thoroughly mixed with water before adding any other material.
5. As the tank is filling, add tank mix partners (if desired) then add the necessary volume of nonionic surfactant. Always add surfactant last.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. ESCORT® XP spray preparations are stable if they are pH neutral or alkaline and stored at or below 100° F.
8. If ESCORT® XP and a tank mix partner are to be applied in multiple loads, pre-slurry the ESCORT® XP in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the ESCORT® XP.

SPRAYER CLEANUP

Spray equipment must be cleaned before ESCORT® XP is sprayed. Follow the cleanup procedures specified on the labels of previously applied products. If no directions are provided, follow the six steps outlined below.

At the End of the Day

When multiple loads of ESCORT® XP herbicide are applied, it is recommended that at the end of each day of spraying, the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits that can accumulate in the application equipment.

1. Drain tank; thoroughly rinse spray tanks, boom, and hoses with clean water. Loosen and physically remove any visible deposits.
2. Fill the tank with clean water and 1 gal of household ammonia* (contains 3% active) for every 100 gal of water. Flush the hoses, boom, and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 min. Flush the hoses, boom, and nozzles again with the cleaning solution, and then drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat step 2.
5. Rinse the tank, boom, and hoses with clean water.
6. If only Ammonia is used as a cleaner, the rinsate solution may be applied back to the crop(s) recommended on this label. Do not exceed the maximum labeled use rate. If other cleaners are used, consult the cleaner label for rinsate

disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

- * Equivalent amounts of an alternate-strength ammonia solution or a DuPont-approved cleaner can be used in the cleanout procedure. Carefully read and follow the individual cleaner instructions. Consult your agricultural dealer, applicator, or DuPont representative for a listing of approved cleaners.

Notes:

1. **Attention:** Do not use chlorine bleach with ammonia, as dangerous gases will form. Do not clean equipment in an enclosed area.
2. Steam-cleaning aerial spray tanks is recommended prior to performing the above cleanout procedure to facilitate the removal of any caked deposits.
3. When DuPont™ ESCORT® XP is tank mixed with other pesticides, all required cleanout procedures should be examined and the most rigorous procedure should be followed.
4. In addition to this cleanout procedure, all precleanout guidelines on subsequently applied products should be followed as per the individual labels.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150 - 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS! See **Wind**, **Temperature and Humidity**, and **Temperature Inversions** sections of this label.

Controlling Droplet Size - General Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.

- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - The boom length should not exceed 3/4 of the wing or rotor length - longer booms increase drift potential.
- **Application Height** - Application more than 10 ft above the canopy increases the potential for spray drift.

BOOM HEIGHT

Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. AVOID GUSTY OR WINDLESS CONDITIONS.

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

USE PRECAUTIONS

Injury to or loss of desirable tree or other plants may result from failure to observe the following.

- If equipment is drained or flushed on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be wash or moved into contact with their roots.
- Treatment of powdery, dry soil or light, sandy soil when there is little likelihood of rainfall soon treatment may result in off target movement and possible damage to susceptible crops when soil particles are moved by wind or water. Injury to crops may result if treated soil is washed, blown, or moved onto land used to produce crops. Exposure to DuPont™ ESCORT® XP may injure or kill most crops. Injury may be more severe when the crops are irrigated.
- Applications made where runoff water flows onto agricultural land may injure crops. Applications made during periods of intense rainfall, to soils saturated with water, surfaces paved with materials such as asphalt or concrete, or soils through which rainfall will not readily penetrate may result in runoff and movement of ESCORT® XP. Do not treat frozen soil. Treated soil should be left undisturbed to reduce the potential for ESCORT® XP movement by soil erosion due to wind or water.
- Do not use on lawns, walks, driveways, tennis courts or similar areas.
- Do not apply through any type of irrigation system.
- Do not use the equipment used to mix or apply ESCORT® XP on crops (except pasture, range and wheat). The mixing and application equipment may be used for noncrop areas and conifer plantations only.
- When used as directed, there is no grazing restriction for use rates of 1 2/3 ounce per acre and less. At use rates of 1 2/3 to 3 1/3 ounce per acre forage grasses may be cut for hay, fodder or green forage and fed to livestock, including lactating animals, 3 days after treatment.
- Do not use this product in the following counties of Colorado: Saguache, Rio Grande, Alamosa, Costilla and Conejos.
- Do not use this product in California.

STORAGE AND DISPOSAL

Storage: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a cool, dry place.

Product Disposal: Do not contaminate water, food, or feed by disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: For Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **For Fiber Sacks:** Completely empty fiber sack by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into manufacturing or application equipment. Then dispose of sack in a sanitary landfill or by incineration if allowed by State and local authorities. **For Fiber Drums With Liners:** Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner. **For Bags Containing Water Soluble Packets:** Do not reuse the outer box or the resealable plastic bag. When all water-soluble packets are used, the outer packaging should be clean and may be disposed of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by open burning. If burned, stay out of smoke. If the resealable plastic bag contacts the formulated product in any way, the bag must be triple-rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer wrap as described above. **For Metal Containers (non aerosol):** Triple rinse (or equivalent) the container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. **For Paper and Plastic Bags:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

13/13

The DuPont Oval Logo, DuPont™, ESCORT®, OUST® and VELPAR® are trademarks or registered trademarks of E.I. duPont de Nemours & Company.

"Arsenal" is a registered trademark of BASF Specialty Products.

"Accord" is a registered trademark of Monsanto Company.

"Embark" is a registered trademark of PBI Gordon Corporation.

"Garlon" and "Tordon" are registered trademarks of Dow Agrosciences.

D - 883 011603

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product; crop injury, or, injury to non-target crops or plants.

DuPont does not agree to be an insurer of these risks. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT

Internet address: <http://cropprotection.dupont.com/>

© 2001-2003 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898.

All rights reserved.