



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

March 26, 2015

Rebecca Ashley
Product Registration Manager
DuPont Crop Protection
Stine-Haskell Research Center
P.O. Box 30
Newark, DE 19714-0030

Subject: Supplemental Label Amendment – Revise expiration date
Product Name: DuPont Escort XP Herbicide
EPA Registration Number: 352-439
Application Date: 3-13-15
Decision Number: 502631

Dear Ms. Ashley:


The supplemental label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. The supplemental label must be incorporated into the master label within 18 months of the date of this notice.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Erik Kraft by phone at 703-308-9358, or via email at kraft.erik@epa.gov.

Sincerely,


A handwritten signature in black ink, appearing to read "Shaja Joyner", with a stylized flourish at the end.

Shaja Joyner, Product Manager 20
Fungicide and Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure



**DuPont Crop
Protection**



| | | |
|-------|---|-----------|
| GROUP | 2 | HERBICIDE |
|-------|---|-----------|

SUPPLEMENTAL LABELING

**DUPONT™ ESCORT® XP
HERBICIDE
FOR PASTURE, RANGELAND AND
CONSERVATION RESERVE PROGRAM**

DUPONT™ ESCORT® XP HERBICIDE

EPA REG. NO. 352-439

FOR USE ON PASTURE, RANGELAND AND CONSERVATION RESERVE PROGRAM

This supplemental label expires February 28, 2016 and must not be used or distributed after this date.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

DUPONT™ ESCORT® XP Herbicide is registered for the control of broadleaf weeds, brush and several woody vine species in forage grasses growing in pasture, rangeland, and Conservation Reserve Program (CRP).

ESCORT® XP may be tank mixed with other pesticides labeled for use in pasture, rangeland and CRP. Read and follow the labels on all products used in the tank mix. Observe the most restrictive precautions on each of the product's labels. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Application of ESCORT® XP to pasture, rangeland and CRP may be made by ground or air. Use a sufficient volume of water to ensure thorough coverage of the targeted weeds with the equipment being used. In Idaho, Oregon and Washington use a minimum application volume of 3 gallons of spray solution per acre.

APPLICATION INFORMATION FOR GRASS ESTABLISHMENT IN PASTURE, RANGELAND AND CRP

ESCORT® XP is registered for the control or suppression of broadleaf weeds to aid in the establishment of the following perennial native or improved grasses planted in pasture, rangeland or acres enrolled in the Conservation Reserve Program

(CRP):

| | | |
|-------------------|----------------|-----------------|
| Blue Grama | Atherstone | intermediate |
| Bluestems -- | Sand | pubescent |
| Big | Weeping | Siberian |
| Little | Wilman | slender |
| Plains | Orchardgrass | steambank |
| Sand | Sideoats grama | tall |
| W W spar | Switchgrass - | thickspike |
| Buffalograss | Blackwell | western |
| Green sprangletop | Wheatgrasses - | Wildrye grass - |
| Kleingrass | Bluebunch | Russian |
| Lovegrasses -- | crested | |

Maximize potential for grass establishment by consulting with the Natural Resource and Conservation Service of other government agencies or local experts concerning planting techniques and other cultural practices.

Performance from ESCORT® XP may not always be satisfactory due to the inability of newly planted grass stands to sufficiently compete with weeds, and the severity of weed pressure in new grass stands.

An additional herbicide application or mowing may be needed.

Use Rates and Application Timing for Grass Establishment in pasture, rangeland and CRP.

Preplant (prior to planting) or Preemergence (after planting but before grass emergence)

Do not use more than 1/10 ounce/acre of ESCORT® XP for grass establishment in pasture, rangeland and CRP.

Apply ESCORT® XP at 1/10 ounce/acre on all labeled grasses except orchardgrass and Russian wildrye grass. Do not apply ESCORT® XP preplant or preemergence to orchardgrass and Russian wildrye grass as severe crop injury may result.

For product information call 1-888-6-DUPONT

© 2015 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898

Early postemergence to new plantings

Apply ESCORT® XP at 1/10 ounce/acre, plus a non-ionic surfactant at the rate of 2 to 4 pints/100 gallons of spray solution on all labeled grasses anytime after grass emergence.

Do not use a spray adjuvant other than non-ionic surfactant. Because grass species differ in time of emergence, apply only after the majority of grasses are in the 3 to 4 leaf stage.

Postemergence to stands with 1 – 5 leaf grasses planted the previous season.

Apply ESCORT® XP at 1/10 ounce/acre plus a non-ionic surfactant at the rate of 2 to 4 pints/100 gallons of spray solution, on all labeled grasses when the majority of the grasses have one or more leaves.

Do not use a spray adjuvant other than non-ionic surfactant.

APPLICATION INFORMATION FOR ESTABLISHED GRASSES IN PASTURE, RANGELAND AND CRP

Use Rates for Established Grasses in Pasture and Rangeland

Apply up to 1 2/3 ounces ESCORT® XP per acre as a broadcast application to established grasses in pasture, rangeland and CRP. For spot applications, use 1 ounce per 100 gallons of water. Do not apply more than 1 2/3 ounces of ESCORT® XP per acre per year in pasture, rangeland and CRP.

Refer to the Weeds Controlled section of the section 3 label for a listing of the weeds controlled by ESCORT® XP and the appropriate use rate to obtain control.

Application Timing – Established Grasses in Pasture Rangeland and CRP

ESCORT® XP may be applied to established native grasses such as bluestems and grama, and on other established grasses such as bermudagrass, bluegrass, orchardgrass, bromegrass, fescue and timothy that were planted the previous growing season (or earlier) and are fully tillered, unless otherwise directed on this label. Specific application timing information on several of these grass species follows:

| <u>Grass</u> | <u>Minimum time from Grass establishment ESCORT® XP application</u> |
|--|---|
| Bermudagrass | 2 months |
| Bluegrass, bromegrass, Orchardgrass | 6 months |
| Timothy | 12 months |
| Fescue | 24 months |

Fescue Precautions:

Note that ESCORT® XP may temporarily stunt tall fescue, cause it to turn yellow, or cause seedhead suppression. To minimize these symptoms, take the following precautions:

- Do not use more than 4/10 ounce/acre of ESCORT® XP
- Tank mix ESCORT® XP with 2,4-D
- Use the lowest specified rate for target weeds
- Use a non-ionic surfactant at 1/2 to 1 pint per 100 gallons of spray solution
- Make application later in the spring after the new growth is 5

to 6 inches tall, or in the fall

- Do not use surfactant when liquid nitrogen is used as a carrier
- Do not use a spray adjuvant other than non-ionic surfactant

The first cutting yields may be reduced due to seedhead suppression resulting from treatment with ESCORT® XP.

Timothy Precautions:

Timothy should be at least 6 inches tall at application and be actively growing. Applications of ESCORT® XP to timothy under any other conditions may cause crop yellowing and/or stunting. To minimize these symptoms, take the following precautions:

- Do not use more than 4/10 ounce/acre ESCORT® XP
- Tank mix ESCORT® XP with 2, 4-D
- Use the lowest specified rate for target weeds
- Use a non-ionic surfactant at 1/2 pint per 100 gallons of spray solution (1/16%)
- Make applications in the late summer or fall
- Do not use surfactant when liquid nitrogen is used as a carrier
- Do not use spray adjuvant other than non-ionic surfactant

Application of ESCORT® XP to Pensacola bahiagrass, ryegrass (Italian or perennial) and Garrison's creeping foxtail may cause severe injury to and/or loss of forage.

Other Pasture and Rangeland Grasses

Varieties and species of forage grasses differ in their tolerance to herbicides. When using ESCORT® XP on a particular grass for the first time, limit use to a small area. In no injury occurs throughout the season, larger acreage may be treated the following season.

Broadleaf forage species, such as alfalfa and clover, are highly sensitive to ESCORT® XP and will be severely stunted or injured by ESCORT® XP.

SPOT TREATMENTS

ESCORT® XP Herbicide may be used for use as spot treatment to control noxious and troublesome weeds on pasture, rangeland and CRP.

Application Information

DuPont™ ESCORT® XP may be used to control many species of weeds, including noxious weeds, in forage grasses growing on pasture, rangeland and CRP. Refer to the "Weeds Controlled" section of the package label or supplemental labeling for a listing of susceptible weed species. If the sprayer is calibrated, consult the package label or other supplemental labeling to select the application rate per acre of ESCORT® XP appropriate for the target weeds. Or mix one gram of ESCORT® XP per one gallon of water along with a suitable surfactant. Spray to the point of wetting the entire surface of the target weeds, approximately 40 gallons of solution per acre. When applied in this manner there is no grazing restrictions following the use of ESCORT® XP. Applications may be made at anytime of the year, except when the soil is frozen.

CROP ROTATION

Before using ESCORT® XP, carefully consider your crop rotation plans and options. For rotational flexibility, do not treat

For product information call 1-888-6-DUPONT

© 2015 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898

Rotation Intervals in Pasture, Rangeland and CRP for Overseeding and Renovation

| Location | Crop or Grass Species | Maximum ESCORT®XP Rate on Pasture Rangeland, and CRP (oz per A) | Minimum Rotation Interval (months) |
|--|---|---|------------------------------------|
| AL, AR, FL, GA, KY, LA, MS, NC, OK, SC, TN, TX, VA, WV | Alfalfa, red clover, white clover, sweet clover, bermudagrass, bluegrass, ryegrass, tall fescue | 1/10 to 3/10 | 4 |
| | Wheat (except durum) | 1/10 to 3/10 | 1 |
| | Durum, barley, oat | 1/10 to 3/10 | 10 |
| ALL STATES NOT INCLUDED ABOVE | Red clover, white clover, and sweet clover | 1/10 to 2/10 | 12 |
| | Bermudagrass, bluegrass, ryegrass | 1/10 to 2/10 | 6 |
| | Tall Fescue | 1/10 to 2/10 | 18 |
| | Wheat (except durum) | 1/10 to 2/10 | 1 |
| | Durum, barley, oat | 1/10 to 2/10 | 10 |
| ALL AREAS WITH SOIL PH OF 7.5 OR LESS | Russian wildrye | 1/10 to 1/2 | 1 |
| | Green needlegrass, switchgrass, sheep fescue | 1/10 to 1 | 1 |
| | Meadow brome, smooth brome, alta fescue, red fescue, meadow foxtail, orchardgrass, Russian wildrye, timothy | 1/10 to 1 | 2 |
| | Alkali sacaton, mountain brome, blue grama thickspike wheatgrass | 1/10 to 1 | 1 |
| ALL AREAS WITH SOIL PH OF 7.9 OR LESS | Sideoats grama, switchgrass | 1/10 to 1/2 | 2 |
| | Western wheatgrass | 1/10 to 1 | 2 |
| | Sideoats grama, switchgrass, big bluestem | 1/10 to 1 | 3 |

all of your pasture, rangeland or CRP acres at the same time.

Minimum Rotational Intervals

Minimum rotation intervals* are determined by the rate of breakdown of ESCORT® XP applied. ESCORT® XP breakdown in the soil is affected by soil pH, presence of soil microorganisms, soil temperature, and soil moisture. Low soil pH, high soil temperature, and high soil moisture increase ESCORT® XP breakdown in soil, while high soil pH, low soil temperature, and low soil moisture slow ESCORT® XP breakdown.

Of these 3 factors, only soil pH remains relatively constant. Soil temperature, and to a greater extent, soil moisture, can vary significantly from year to year and from area to area. For this reason, soil temperatures and soil moisture should be monitored regularly when considering crop rotations.

* The minimum rotation interval represents the period of time from the last application to the anticipated date of the next planting.

Soil pH Limitations

ESCORT® XP should not be used on soils having a pH above 7.9, as extended soil residual activity could extend crop rotation intervals beyond normal. Under certain conditions, ESCORT® XP could remain in the soil for 34 months or more, injuring wheat and barley. In addition, other crops planted in high-pH soils can be extremely sensitive to low concentrations of ESCORT® XP.

Checking Soil pH

Before using ESCORT® XP, determine the soil pH of the areas of intended use. To obtain a representative pH value for the test

area, take several 0" to 4" samples from different areas of the field and analyze them separately. Consult local extension publications for additional information on recommended soil sampling procedures.

BIOASSAY

A field bioassay must be completed before rotating to any crop or grass species/variety not listed in the Rotation Intervals Table, or if the soil pH is not in the specified range, or if the use rate applied is not specified in the table.

To conduct a field bioassay, grow test strips of the crop(s) or grass(es) you plan to grow the following year in fields previously treated with ESCORT® XP. Crop or grass response to the bioassay will indicate whether or not to rotate to the crop(s) or grass(es) grown in the test strips.

If a field bioassay is planned, check with your local Agricultural dealer or DuPont representative for information detailing the field bioassay procedure.

GRAZING/HAYING

When used as directed, there is no grazing or haying restriction for use rates of 1 2/3 ounces per acre and less.

Coveralls, shoes plus socks must be worn if cutting within 4 hours of treatment.

IMPORTANT PRECAUTIONS

- Do not apply more than 1 2/3 ounces of ESCORT® XP per acre per year on pasture, rangeland or CRP.
- Grass species or varieties may differ in their response to

For product information call 1-888-6-DUPONT

© 2015 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898

various herbicides. DuPont recommends that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use of ESCORT® XP to a small area. Components in a grass seed mixture will vary in tolerance to ESCORT® XP so the final stand may not reflect the seed ratio.

- Under certain conditions such as heavy rainfall, high pH, prolonged cold weather, or wide fluctuations in day/night temperatures prior to or soon after ESCORT® XP application, temporary discoloration and/or grass injury may occur. ESCORT® XP should not be applied to grass that is stressed by severe weather conditions, drought, low fertility, water-saturated soils, disease, or insect damage as grass injury may result. Severe winter stress, drought, disease, or insect damage before or following application also may result in grass injury.
- Applications of ESCORT® XP to pasture, rangeland, and CRP undersown with legumes may cause injury to the legumes. Legumes in a seeding mixture may be severely injured or killed following an application of ESCORT® XP.
- Applications made to some established grasses may cause temporary stunting, yellowing or seedhead suppression (i.e. fescue, timothy).
- Applications made to newly established grasses less than 2 years from seeding may result in injury or loss.
- Do not apply to forage grasses known to be sensitive to ESCORT® XP such as ryegrass (Italian and perennial), bahia or Garrison's creeping foxtail.
- Broadleaf forage species, such as alfalfa and clover, are highly sensitive to ESCORT® XP and will be severely injured or killed.
- The control of weeds in wheel track areas may be reduced if ground applications are made when dry, dusty field conditions exist. The addition of 2,4-D or MCPA should improve weed control under these conditions.
- Do not use this product in the following counties of Colorado: Saguache, Rio Grande, Alamosa, Costilla and Conejos.
- Do not use this product in California.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS LABEL, READ THIS LABEL AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS LABEL, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

This label contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This label must be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

(Replaces: R-1282 060412 05-30-12)

(Ref: SL - 1612MSTR 022912 02-29-12)

For product information call 1-888-6-DUPONT

© 2015 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898