




REG. U.S. PAT. & TM. OFF.

Krovar II[®] DF[™]

HERBICIDE



TRADEMARK

DRY FLOWABLE

NET 6 POUNDS



HERBICIDE DISPERSIBLE GRANULE

ACTIVE INGREDIENTS	80%
Bromacil [5-bromo-3-sec-butyl-6-methyluracil]	53%
Difluron [3-(3,4-dichlorophenyl)-1,1-dimethylurea]	27%
INERT INGREDIENTS	20%

EPA Reg. No. 352-XXX

**Keep out of reach of children.
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS**

CAUTION! HARMFUL IF SWALLOWED. CAUSES MODERATE EYE INJURY.

Avoid breathing dust or spray mist. Avoid contact with skin, eyes or clothing. In case of contact with eyes, immediately flush with plenty of water. Get medical attention if irritation persists. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water. Call a physician if irritation persists.

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

E. I. Du Pont de Nemours & Co. (Inc.), Agricultural Chemicals Department, Wilmington, De

IMPORTANT—Injury to or loss of desirable trees or other plants may result from failure to observe the following:

Do not apply (except as recommended for crop use), or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks or driveways, tennis courts or similar areas. Do not use in home fruit plantings nor in citrus orchards interplanted to other trees or desirable plants. Prevent drift of dry powder or spray to desirable plants. Do not contaminate any body of water. Keep from contact with fertilizers, insecticides, fungicides and seeds.

Thoroughly clean all traces of "Krovar" II DF from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

GENERAL INFORMATION

Du Pont "Krovar" II DF Herbicide is a dry flowable granule to be mixed in water and applied as a spray for selective control of weeds in crops and for non-crop weed control.

"Krovar" II DF controls certain perennial grasses and annual weeds such as barnyardgrass (watergrass), bermudagrass, crabgrass, Florida pusley, foxtail, guinea grass (partial control), goosegrass, lambsquarters, natalgrass (red top), nightshade (annual), pigweed, purslane, ragweed, sandbur (sandspur), sowthistle (annual), and spanishneedles. Effects on perennial weeds are slow to appear, usually progressing over a period of several months. In Florida, it is also recommended for control of certain seeding vines.

Moisture is necessary to activate the herbicide, best results are obtained if treatment is made to moist soil, and moisture is supplied by rain, sprinkler irrigation within two weeks after application. The degree of control and duration of effect will vary with the amount applied, soil texture, rainfall, and other conditions.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

"Krovar" II DF should be used only in accordance with recommendations on this label, or in separate published Du Pont recommendations available through local dealers.

Du Pont will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by Du Pont. User assumes all risk associated with such non-recommended use.

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Warnings must include the following information: "CAUTION: Area treated with "Krovar" II DF on (date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of contact flush eyes with plenty of water; wash skin with plenty of soap and water. Get medical attention if irritation persists."

CITRUS

Use only where trees have been established in the orchard for at least four years. Apply with a properly calibrated fixed-boom power sprayer as a band or broadcast treatment beneath and/or between trees. Use sufficient water (min. 40 gals. per acre) to obtain uniform coverage; avoid contact of foliage and fruit with spray or mist. Avoid overlapping, and shut off spray boom while starting, turning, slowing or stopping, or injury to trees may result. Continuous agitation in the spray tank is required to keep the material in suspension. Best results are obtained if application is made shortly before or shortly after weed growth begins; if dense growth is present, remove tops and spray the ground. Do not graze cattle in treated areas. Dosages are expressed as broadcast rates; for band treatment, use proportionately less.

Do not use on citrus in Kern County, California.

Perennial Weeds: For bahiagrass, bermudagrass, nutsedge, pangolagrass, paragrass, and torpedograss, make a single application per year during the period from winter to early summer at the following rates.

Soil Texture	Lbs. Per Acre
Sand, loamy sand	4 to 6
Sandy loam	5 to 7
Silt loam, clay loam	6 to 8

Go to last page for Herbicide

Alternatively, make two applications of 3 to 4 lbs. per acre per year (Florida 4 to 8 lbs.) apply in spring and summer.

Partial control of perennials usually occurs with a single treatment; repeat applications are required to control perennials. Control may be improved by cultivation prior to treatment, otherwise, avoid working the soil as long as weed control continues or else effectiveness of the treatment may be reduced.

Annual Weeds: Apply 2 to 5 lbs. per acre using the lower rates on coarse soils. apply any time of year provided overhead moisture (rainfall or sprinkler irrigation) is available to activate the herbicide, just before or just after weeds germinate.

Note: Temporary yellowing of citrus leaves may occur following treatment. Because injury to citrus trees may result, do not use on soils low in organic matter (less than 1%) poorly drained soils, gravelly soils, not thinly covered or exposed subsoils, do not apply more than 10 lbs. per acre per year (Florida, 16 lbs.). do not treat trees planted in irrigation furrows, do not treat diseased trees such as those with foot rot. Do not replant treated areas to any crop within two years after last application as injury to subsequent crops may result, except that citrus trees may be planted one year after the last application.

Florida Only—

Trees Established Less Than Four Years

Trees Established Less Than One Year:

Apply 2 to 4 lbs. "Krovar" II DF per acre as needed to maintain weed control. Do not apply at less than 60-day intervals, do not apply more than 6 lbs. per acre during any 6-month period or more than 8 lbs. per acre during the first year.

Trees Established One to Four Years:

Apply 3 to 6 lbs. "Krovar" II DF per acre, for details on timing, equipment, spray volume and other use directions, refer to product label. Use the lower rate for 1 to 2 year old trees and the higher rate for trees more than 2 years old. A second application may be made if needed, but do not apply more than 8 lbs. "Krovar" II DF per acre per year as injury to the trees may result.

Tank Mixtures:

"Krovar" II DF is recommended for use on citrus for weed control in tank mixtures with paraquat. For rates of the individual products, refer to the respective product label. Do not apply tank mixture to foliage or green wood as injury may result.

Seedling Vines (balsam apple and milkweed)—

Florida: Remove established vines from trees prior to treatment. Apply 6 lbs. per acre after vine growth begins in the spring (or after cultivation or chopping), but before vines reach tree limbs. For band treatment, the treated area should extend at least 1 foot beyond the dripline of the tree. Add Du Pont Surfactant WK at a rate of 1 qt. per 50 gals. of spray mixture.

A repeat application may be made in late summer or early fall to control seedlings which germinate late in the season.

When used in conjunction with good mechanical cultivation practices, repeated annual applications aid in the suppression of established vines.

NON-CROP WEED CONTROL

For short-term control of annual weeds on non-cropland areas such as roadsides, storage areas and plant sites, apply 2 to 5 lbs. per acre in 40 to 100 gals. water. For extended control of annuals and partial control of perennials such as bermudagrass and nutsedge, apply 6 to 14 lbs. per acre. For control of hard-to-kill perennials such as bermudagrass, bouncing-bet, dogbane, johnsongrass, nutsedge, and saltgrass, apply 15 to 30 lbs. per acre.

Use the higher levels of dosage rates on adsorptive soils (high in organic matter or carbon). Best results occur when application is made just before weed emergence or in early stages of weed growth.

Retreatment: Apply 2 to 5 lbs. per acre when annual weeds and grasses reappear on sites where weed growth has been controlled.

Small Areas: 1 cupful of "Krovar" II DF per 200 sq. ft. is approximately 15 lbs. per acre.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

STORAGE AND DISPOSAL


STORAGE: Store product in original container only, away from other pesticides, fertilizer, food or feed.

DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Completely empty box into application equipment. Then dispose of empty box in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

AG 652 0485

Made in U.S.A.

Printed in U.S.A.



Reg no: 352-440

PM-25

192

SPECIMEN LABEL

86% OF ACTUAL SIZE

NOT REVIEWED

In Accordance with FR Notice S2-2.

Based on Draft Labeling Dated


2/11/76



REG US PAT & TM-OFF

Krovar II DF™

HERBICIDE



TRADE MARK

DRY FLOWABLE

NET 6 POUNDS



Krovar II DF™

HERBICIDE DISPERSIBLE GRANULE

ACTIVE INGREDIENTS

- Bromacil [5-bromo-3-sec-butyl-6-methyluracil]
- Diuron [3-(3,4-dichlorophenyl)-1,1-dimethylurea]

INERT INGREDIENTS

80%

53%

27%

20%

EPA REG. NO. 101-101-01

SPECIMEN

73% OF ACTUAL

Keep out of reach of children. PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION! HARMFUL IF SWALLOWED. CAUSES MODERATE EYE INJURY.

Avoid breathing dust, mist or gas. Avoid contact with skin, eyes or clothing. In case of contact with eyes, immediately flush with plenty of water. Get medical attention if irritation persists. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before use.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water. Call a physician if irritation persists.
If swallowed: Call a physician. Do not induce vomiting. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not eat, drink or use tobacco during treatment. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

IMPORTANT—Do not use Krovar II DF herbicide on citrus trees or plants in any area where the fruit will be harvested. Do not use Krovar II DF herbicide on citrus trees or plants in any area where the fruit will be harvested. Do not use Krovar II DF herbicide on citrus trees or plants in any area where the fruit will be harvested. Do not use Krovar II DF herbicide on citrus trees or plants in any area where the fruit will be harvested.

GENERAL INFORMATION

Krovar II DF herbicide is a selective herbicide which controls a wide range of broadleaf weeds and grasses in citrus groves. It is effective against many of the most troublesome weeds and grasses in citrus groves.

Krovar II DF herbicide is a selective herbicide which controls a wide range of broadleaf weeds and grasses in citrus groves. It is effective against many of the most troublesome weeds and grasses in citrus groves.

Krovar II DF herbicide is a selective herbicide which controls a wide range of broadleaf weeds and grasses in citrus groves. It is effective against many of the most troublesome weeds and grasses in citrus groves.

DIRECTIONS FOR USE

Apply Krovar II DF herbicide to citrus groves in the following manner: Apply 2 to 4 lbs. per acre as needed to maintain weed control. Do not apply at less than 15 day intervals. Do not apply more than 6 lbs. per acre during any 6 month period or more than 8 lbs. per acre during the first year.

Apply Krovar II DF herbicide to citrus groves in the following manner: Apply 2 to 4 lbs. per acre as needed to maintain weed control. Do not apply at less than 15 day intervals. Do not apply more than 6 lbs. per acre during any 6 month period or more than 8 lbs. per acre during the first year.

Apply Krovar II DF herbicide to citrus groves in the following manner: Apply 2 to 4 lbs. per acre as needed to maintain weed control. Do not apply at less than 15 day intervals. Do not apply more than 6 lbs. per acre during any 6 month period or more than 8 lbs. per acre during the first year.

Apply Krovar II DF herbicide to citrus groves in the following manner: Apply 2 to 4 lbs. per acre as needed to maintain weed control. Do not apply at less than 15 day intervals. Do not apply more than 6 lbs. per acre during any 6 month period or more than 8 lbs. per acre during the first year.

Apply Krovar II DF herbicide to citrus groves in the following manner: Apply 2 to 4 lbs. per acre as needed to maintain weed control. Do not apply at less than 15 day intervals. Do not apply more than 6 lbs. per acre during any 6 month period or more than 8 lbs. per acre during the first year.

Apply Krovar II DF herbicide to citrus groves in the following manner: Apply 2 to 4 lbs. per acre as needed to maintain weed control. Do not apply at less than 15 day intervals. Do not apply more than 6 lbs. per acre during any 6 month period or more than 8 lbs. per acre during the first year.

Apply Krovar II DF herbicide to citrus groves in the following manner: Apply 2 to 4 lbs. per acre as needed to maintain weed control. Do not apply at less than 15 day intervals. Do not apply more than 6 lbs. per acre during any 6 month period or more than 8 lbs. per acre during the first year.

Apply Krovar II DF herbicide to citrus groves in the following manner: Apply 2 to 4 lbs. per acre as needed to maintain weed control. Do not apply at less than 15 day intervals. Do not apply more than 6 lbs. per acre during any 6 month period or more than 8 lbs. per acre during the first year.

Apply Krovar II DF herbicide to citrus groves in the following manner: Apply 2 to 4 lbs. per acre as needed to maintain weed control. Do not apply at less than 15 day intervals. Do not apply more than 6 lbs. per acre during any 6 month period or more than 8 lbs. per acre during the first year.

Apply Krovar II DF herbicide to citrus groves in the following manner: Apply 2 to 4 lbs. per acre as needed to maintain weed control. Do not apply at less than 15 day intervals. Do not apply more than 6 lbs. per acre during any 6 month period or more than 8 lbs. per acre during the first year.

Apply Krovar II DF herbicide to citrus groves in the following manner: Apply 2 to 4 lbs. per acre as needed to maintain weed control. Do not apply at less than 15 day intervals. Do not apply more than 6 lbs. per acre during any 6 month period or more than 8 lbs. per acre during the first year.

Apply Krovar II DF herbicide to citrus groves in the following manner: Apply 2 to 4 lbs. per acre as needed to maintain weed control. Do not apply at less than 15 day intervals. Do not apply more than 6 lbs. per acre during any 6 month period or more than 8 lbs. per acre during the first year.

Apply Krovar II DF herbicide to citrus groves in the following manner: Apply 2 to 4 lbs. per acre as needed to maintain weed control. Do not apply at less than 15 day intervals. Do not apply more than 6 lbs. per acre during any 6 month period or more than 8 lbs. per acre during the first year.

Apply Krovar II DF herbicide to citrus groves in the following manner: Apply 2 to 4 lbs. per acre as needed to maintain weed control. Do not apply at less than 15 day intervals. Do not apply more than 6 lbs. per acre during any 6 month period or more than 8 lbs. per acre during the first year.

Apply Krovar II DF herbicide to citrus groves in the following manner: Apply 2 to 4 lbs. per acre as needed to maintain weed control. Do not apply at less than 15 day intervals. Do not apply more than 6 lbs. per acre during any 6 month period or more than 8 lbs. per acre during the first year.

Apply Krovar II DF herbicide to citrus groves in the following manner: Apply 2 to 4 lbs. per acre as needed to maintain weed control. Do not apply at less than 15 day intervals. Do not apply more than 6 lbs. per acre during any 6 month period or more than 8 lbs. per acre during the first year.

Apply Krovar II DF herbicide to citrus groves in the following manner: Apply 2 to 4 lbs. per acre as needed to maintain weed control. Do not apply at less than 15 day intervals. Do not apply more than 6 lbs. per acre during any 6 month period or more than 8 lbs. per acre during the first year.

Alternatively, make two applications of 3 to 4 lbs. per acre per year (Florida 4 to 8 lbs.) applying in spring and summer.

Partial control of perennials usually occurs with a single treatment. Repeat applications are required to control perennials. Control may be improved by cultivation prior to treatment. Otherwise, avoid working the soil as long as weed control continues or else effectiveness of the treatment may be reduced.

Annual Weeds: Apply 2 to 5 lbs. per acre using the lower rates on coarse soils; apply any time of year provided overhead moisture (rainfall or sprinkler irrigation) is available to activate the herbicide just before or just after weeds germinate.

Note—Temporary yellowing of citrus leaves may occur following treatment. Because injury to citrus trees may result, do not use on soils low in organic matter (less than 1%) poorly drained soils, gravelly soils, nor thinly covered or exposed subsoils; do not apply more than 10 lbs. per acre per year (Florida, 16 lbs.); do not treat trees planted in irrigation furrows; do not treat d-seeded trees such as those with footrot. Do not replant treated areas to any crop within two years after last application as injury to subsequent crops may result, except that citrus trees may be replanted one year after the last application.

Florida Only—

Trees Established Less Than Four Years

Trees Established Less Than One Year. Apply 2 to 4 lbs. Krovar II DF per acre as needed to maintain weed control. Do not apply at less than 15 day intervals. Do not apply more than 6 lbs. per acre during any 6 month period or more than 8 lbs. per acre during the first year.

Trees Established One to Four Years

Apply 3 to 6 lbs. Krovar II DF per acre for details on timing, equipment, spray volume and other use directions, refer to product label. Use the lower rate for 1 to 2 year old trees and the higher rate for trees more than 2 years old. A second application may be made if needed but do not apply more than 8 lbs. Krovar II DF per acre per year as shown in the table below.

Tank Mixtures:

Krovar II DF is recommended for use on citrus for weed control in tank mixtures with paraquat. For rates of the individual products, refer to the respective product label. Do not apply tank mixture to foliage or green wood as injury may result.

Seedling Vines (balsam apple and milkweed)—

Florida: Remove established vines from trees prior to treatment. Apply 6 lbs. per acre after vine growth begins in the spring for after-cultivation or chipping, but before vines reach tree limbs. For band treatment, the treated area should extend at least 1 foot beyond the drip line of the tree. Add DuPont Surfactant WK at a rate of 1 qt. per 50 gals. of spray mixture.

A repeat application may be made in late summer or early fall to control seedlings which germinate after the season.

When used in conjunction with good mechanical cultivation practices, repeated annual applications aid in the suppression of established vines.

- Soil Texture
- Sand, Heavy sand
- Sandy loam
- Silt with heavy sand

- Lbs. Per Acre
- 4 to 6
- 5 to 7
- 6 to 8
- 7 to 10

E. I. du Pont de Nemours & Co. (Inc.), Agricultural Products Department, Wilmington, DE