

"CANOPY" HERBICIDE
EPA.REG. NO. 352-444

SUPPLEMENTAL LABELING
"CANOPY" HERBICIDE REDUCED RATE
FOLLOWED BY "CLASSIC" HERBICIDE
FOR BROADLEAF WEED CONTROL IN SOYBEANS

ACCEPTED

Under the Federal Insecticide
Fungicide and Rodenticide Act
as amended by the pesticide
registered under
EPA Reg No. 352444

Du Pont "Canopy" Herbicide is recommended preplant incorporated or preemergence at reduced rates followed by "Classic" Herbicide postemergence for broadleaf weed control in soybeans. This treatment will control those weeds listed on the "Canopy" and "Classic" Herbicide product labels.

Directions For Use

Apply "Canopy" preplant incorporated or preemergence at reduced rates (see rate chart below) as directed on the product label. Follow the "Canopy" preplant incorporated or preemergence reduced rate application with a postemergence application of "Classic" Herbicide at its labeled rate when weeds are small and actively growing (usually 14-28 days after preplant incorporated or preemergence application of "Canopy" reduced rate). "Canopy" can be applied in tank mix with Lasso[2], Dual[3], Prowl[4], Surflan[5], Treflan[5], Sonalan[5] and Command[6]. Read and follow the product label for use of these products.

"Canopy" Reduced Rate

<u>Soil Texture</u>	<u>Rate in Ounces per acre</u>	
	<u>1/2 - 4% Organic Matter</u>	
<u>COURSE; Loamy Sand, Sandy Loam</u>	4-6	
<u>MEDIUM; Loam, Siltloam, Silt, Sandy Clay Loam</u>	4-6	
<u>FINE; Silty Clay Loam, Clay Loam, Clay</u>	6-8	

- Do not apply to soils with less than 1/2% organic matter or greater than 4% organic matter.

Rotational Crop Guidelines

(A) STATES OF IA, IL, IN, KS, MI, MO, MN, NE, OK, OH AND WI

The following "Canopy" rates followed by a postemergence "Classic" Herbicide application may be recropped as indicated in the chart below. For "Canopy" rates other than those listed or for more than one "Classic" application, refer to the "Canopy" and "Classic" Herbicide labels for rotational crop information.

- "Canopy" rate of 4-5 ounces per acre and a single application of "Classic" Herbicide postemergence of up to 3/4 ounce per acre.
- "Canopy" rate of 6 ounces per acre and a single application of "Classic" Herbicide postemergence of no more than 2/3 ounce per acre.
- "Canopy" rate of 7 ounces per acre and No "Classic" Herbicide postemergence.

<u>Crop</u>	<u>Recropping Interval in Months</u>
Soybeans	Anytime
Barley	4
Ryegrass	4
Wheat	4
Alfalfa	10
Cotton	10
Clover	12
Field Corn	10
Dry Beans	12
Sorghum	12
Rice	10
Tobacco (transplant)	10
Tomato (transplant)	10

(B) ALL STATES NOT LISTED ABOVE FOLLOW ROTATIONAL AND SOIL pH RESTRICTIONS ON THE "CANOPY" AND "CLASSIC" HERBICIDE LABELS.

IMPORTANT

Before using "Canopy" and "Classic" Herbicides, read and carefully note the cautionary statements and all other information appearing on the product labels.

This bulletin contains new or supplemental information for use of these products which does not appear on the product label.

Follow the instructions carefully.

This label must be in the possession of the user at the time of pesticide application.

- [1]Registered Trademark of E. I. du Pont de Nemours & Co. (Inc)
- [2]Registered Trademark of Monsanto Company
- [3]Registered trademark of Ciba-Giegy Corporation
- [4]Registered Trademark of American Cyanamid Company
- [5]Registered Trademark of Elanco Products Company
- [6]Registered Trademark of FMC Corporation

E. I. du Pont de Nemours & Co. (Inc.)
Agricultural Products Department
Wilmington, Delaware 19880-0038

