

"CANOPY" HERBICIDE  
EPA REG. NO. 352-444

SUPPLEMENTAL LABELING  
"CANOPY" HERBICIDE FOR COMMERCIAL IMPREGNATION  
AND APPLICATION ON DRY BULK FERTILIZER

DIRECTIONS FOR USE

It is a violation of federal law to use this product  
in a manner inconsistent with its labeling.

Do not "CANOPY" Herbicide may be applied with dry bulk fertilizer for  
preemergence weed control. Dry bulk fertilizer may be impregnated or coated  
with "CANOPY" for application to soybeans. All recommendations, cautions and  
special precautions on this label must be followed along with state  
regulations relating to dry bulk fertilizer blending, impregnating and  
labeling.

HOW TO USE

Application: Uniform application of "CANOPY" which has been impregnated or  
dry fertilizer is essential for satisfactory weed control. Accurate  
calibration of fertilizer application equipment is essential for uniform  
distribution to the soil. Air flow or auger metered application equipment  
is preferred (one pass application). If other equipment is used, the  
recommended method of application is to apply 1/2 the recommended rate and  
overlap 50 percent to double apply by splitting the middles to obtain the  
best distribution pattern.

If fertilizer materials are excessively dusty, use diesel oil or other  
suitable additive to reduce dust prior to impregnation as dusty fertilizer  
will result in poor distribution during application. Crop injury and poor  
weed control may occur where the impregnated fertilizer is not uniformly  
applied.

Incorporation and Combination Uses: When "CANOPY" is to be used in  
combination with another herbicide, follow directions on this label for  
combinations, rates, crops, incorporation and special precautions.

Impregnation: To impregnate, use a system consisting of a survey meter and  
trump, which is used for dry bulk fertilizer blending. Any commercial  
fertilizer can be impregnated with "CANOPY" except ammonium nitrate or  
fertilizers containing ammonium nitrate, potassium nitrate or sodium nitrate.  
Do not use on limestone.

ACCEPTED

U.S. Federal Government  
Registration Act as amended for L.  
pesticide registration under  
EPA Reg. No. 352-444

BEST AVAILABLE COPY

vary, and if the adsorptivity is not adequate, the use of an adsorptive powder may be required to produce a dry freeflowing mixture. Microcel E (Johns Manville Product Corporation) is the recommended absorbent powder. "CANOPY" should be pre-mixed with water to form a sprayable mixture. When another herbicide is used with "CANOPY", mix and impregnate immediately.

**Clean Out:** Before using equipment to apply fertilizer or herbicide impregnated fertilizer to another crop other than soybeans, at least 1,000 lbs. of fertilizer not treated with "CANOPY" must be run through the application equipment and impregnation drum. All equipment containing the herbicide slurry should be cleaned out in accordance with the "CANOPY" recommendations for application with liquid delivery systems.

Apply immediately after impregnation unless experience has shown that impregnated fertilizer can be stored without becoming lumpy and difficult to spread.

**Rates:** Select the rate of "CANOPY" to apply per acre from the appropriate section of this label and refer to the rate table to determine the amount of "CANOPY" which is to be impregnated on a ton of dry bulk fertilizer based on the amount of fertilizer which will be distributed on one acre.

**RATE CHART FOR IMPREGNATING FERTILIZER WITH "CANOPY"**

FERTILIZER RATE (LB./ACRE)	4 OZ. ****	6 OZ. ****	8 OZ. ****	10 OZ. ****	12 OZ. ****
	(LB. "CANOPY" PER TON OF FERTILIZER)				
*****					
200 POUNDS	2.50	3.75	5.00	6.25	7.50
250 POUNDS	2.00	3.00	4.00	5.00	6.00
300 POUNDS	1.67	2.50	3.33	4.17	5.00
350 POUNDS	1.43	2.14	2.86	3.57	4.29
400 POUNDS	1.25	1.88	2.50	3.13	3.75
450 POUNDS	1.11	1.67	2.22	2.78	3.33

For rates other than those listed, use the following formula to calculate the amount of "CANOPY" to be impregnated per ton of dry fertilizer.

$$\begin{aligned} \text{lb. "CANOPY"/ton} &= \frac{2000 \text{ lb./ton} \times \text{"CANOPY" rate (oz./acre)}}{\text{Fertilizer rate (lb./acre)} \times 16 \text{ oz./lb.}} \\ &= \frac{125 \times \text{"CANOPY" rate (oz./acre)}}{\text{Fertilizer rate (lb./acre)}} \end{aligned}$$

IMPORTANT

BEFORE USING "CANOPY", ALONE OR IN COMBINATIONS, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABELS.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package labels. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

[1] Registered trademark of E.I. du Pont de Nemours and Company

E.I. du Pont de Nemours and Company  
Agricultural Products Department  
Wilmington, DE 19880-0038

040490

---

"CANOPY" HERBICIDE  
EPA REG. NO. 352-444

SUPPLEMENTAL LABELING  
"CANOPY" HERBICIDE  
AERIAL APPLICATION ON SOYBEANS

DIRECTIONS FOR USE

It is a violation of federal law to use this product  
in a manner inconsistent with its labeling.

Du Pont "Canopy" Herbicide is recommended for use by aerial application on  
soybeans.


HOW TO USE

"Canopy" may be applied by air for early preplant, preplant incorporated or  
preemergence use on soybeans. Apply uniformly with properly calibrated  
aerial equipment. Use a minimum of 2 gallons of water per acre. Avoid over  
applying of spray swath. Continuous agitation of the spray tank is required  
to keep the material in suspension.

Do not allow spray to drift onto adjacent crops or land as injury to other  
plants may occur. To avoid drift, applicators must observe the following  
precautions:

- Do not spray when wind speeds exceed 10 mph.
- Do not spray when winds are gusty.
- Do not spray when a temperature inversion exists.
- Use nozzles which produce a coarse spray. Nozzles should  
be oriented in the direction of the air stream. Do not  
exceed 40 psi.

Refer to the "Canopy" label for additional use instructions, limitations and  
precautions.



This labeling must be in the possession of the user at the time of pesticide application.

[1] Registered trademark of E.I. du Pont de Nemours and Company

E.I. du Pont de Nemours and Company  
Agricultural Products Department  
Wilmington, DE 19880-0038

040990