



**DuPont Agricultural
Products**

..... A Growing Partnership With Nature



SUPPLEMENTAL LABELING

**CANOPY® HERBICIDE
TANK MIXES FOR BROADLEAF
WEED CONTROL IN SOYBEANS
IN CERTAIN MIDWEST STATES**

CANOPY® HERBICIDE

EPA Reg. No. 352-444

"CANOPY" HERBICIDE TANK MIXES FOR BROADLEAF WEED CONTROL IN THE STATES OF INDIANA, ILLINOIS, IOWA, KANSAS, KENTUCKY, MICHIGAN, MISSOURI, NEBRASKA, AND OHIO

DuPont "Canopy" Herbicide is recommended at reduced rates for pre-emergence broadleaf weed control when tank mixed with either "Lexone" DF, "Lexone" SP, "Lorox" DF or "Lorox" SP in the states of Indiana, Illinois, Iowa, Kansas, Kentucky, Michigan, Missouri, Nebraska, and Ohio.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

For use on soybeans only.

Do not apply this product through any type of irrigation equipment.

Do not graze treated fields or harvest forage for hay.

HOW TO USE

APPLICATION TIMING AND METHODS

"Canopy" tank mixed with either "Lexone" DF, "Lexone" SP, "Lorox" DF or "Lorox" SP may be applied at planting or at any time up to 45 days prior to planting. Do not apply after the soybean crop has emerged. These tank mixes may be preemergence surface applied or preplant incorporated. Refer to the "Canopy" label for specific instructions on incorporation and moisture requirements. These tank mixes may also be applied over existing weed vegetation as in no-till soybean production.

APPLICATION EQUIPMENT AND SPRAY VOLUMES

Apply uniformly by ground equipment using a properly calibrated fixed boom sprayer. Use 10-40 gallons per acre at 20-40 psi. In no-till situations, where these tank mixes are being applied over existing vegetation, use flat fan nozzles and a minimum of 20 gallons per acre.

APPLICATION RATES

Select the rate of "Canopy" + "Lexone" DF, "Lexone" SP, "Lorox" DF or "Lorox" SP from the chart below, according to soil type:

Soil Texture	Broadcast Rate (oz/A)				
	"CANOPY" + "LEXONE" OR "LOROX"				
Coarse: Loamy sand, Sandy loam	2.5	+	4	OR	8
Medium: Loam, Silt Loam, Silt, Sandy clay loam	3	+	6	OR	10
Fine: Silty clay loam, Clay loam, Clay	3-4	+	8	OR	12

Do not apply to soils with less than 1/2% or more than 4% organic matter.

*Use the rates listed in the table when tank mixing "Canopy" with either "Lexone" DF or "Lexone" SP or when tank mixing "Canopy" with either "Lorox" DF or "Lorox" SP. "Lexone" SP and "Lorox" SP are water dispersible granules packaged in water soluble film. Refer to the product labels for specific instructions on using and measuring this type of packaging.

WEEDS CONTROLLED

Preemergence:

When used according to this label these "Canopy" tank mixes will generally provide season-long preemergence control of the following weeds:

Lambsquarters
Mustard, wild
Pigweeds:
Palmer
Redroot
Smooth
Spiny amaranth
Tall waterhemp

Rigweed, common
Smartweeds, annual
Velvetleaf

ACCEPTED

MAR 13 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 352-444

© 1993 E. I. du Pont de Nemours and Company, DuPont Agricultural Products, Wilmington, Delaware 19898