




DuPont Agricultural Products

"..... A Growing Partnership With Nature"



PIL 35 352-444 P. 17
SUPPLEMENTAL LABELING

**CANOPY® HERBICIDE
TANK MIXES FOR BROADLEAF
WEED CONTROL IN SOYBEANS
IN CERTAIN MIDWEST STATES**

CANOPY® HERBICIDE

EPA Reg. No. 352-444

CANOPY HERBICIDE TANK MIXES FOR BROADLEAF WEED CONTROL IN THE STATES OF INDIANA AND OHIO

DuPont CANOPY Herbicide is recommended at reduced rates for preemergence broadleaf weed control when tank mixed with DuPont NEW LOROX PLUS® Herbicide in the states of Indiana and Ohio.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

For use on soybeans only.

Do not apply this product through any type of irrigation equipment.

Do not graze treated fields or harvest forage for hay.

HOW TO USE

APPLICATION TIMING AND METHODS

CANOPY tank mixed with NEW LOROX PLUS may be applied up to just before soybean emergence, but no earlier than 45 days before planting. Do not apply after the soybean crop has emerged. These tank mixes may be preemergence surface applied or preplant incorporated. Refer to the CANOPY label for specific instructions on incorporation and moisture requirements. These tank mixes may also be applied over existing weed vegetation as in no-till soybean production.

APPLICATION EQUIPMENT AND SPRAY VOLUMES

Apply uniformly by ground equipment using a properly calibrated fixed boom sprayer. Use 10-40 gallons per acre at 20-40 psi. In no-till situations, where these tank mixes are being applied over existing vegetation, use flat fan nozzles and a minimum of 20 gallons per acre.

APPLICATION RATES

Select the rate of CANOPY + NEW LOROX PLUS from the chart below, according to soil type:

Soil Texture	Broadcast Rate (oz/A) CANOPY+NEW LOROX PLUS		
	2	+	6
Coarse:			
Loamy sand, Sandy loam			
Medium:	2.5	+	7
Loam, Silt Loam,			
Silt, Sandy clay loam			
Fine:	3	+	8
Silty clay loam,			
Clay loam, Clay			

Do not apply to soils with less than 1/2% or more than 4% organic matter.

WEEDS CONTROLLED

Preemergence:

When used according to this label these CANOPY tank mixes will generally provide season-long preemergence control of the following weeds.

Carpetweed	Prickly sida
Lambsquarters	Ragweed, common
Mustard, wild	Smartweeds, annual
Pigweeds:	Velvetleaf
Palmer	
Redroot	
Smooth	
Spiny amaranth	
Tall waterhemp	

These CANOPY tank mixes will also provide partial control (suppression) preemergence of the following weeds:

Cocklebur	Morningglories
Crabgrass	Entireleaf
Eastern black nightshade	Ivyleaf
Foxtail species	Pitted
Immunweed	Tall
	Nutsedge species
	Panicum species

Postemergence (for No-till Burndown):

CANOPY tank mixes will provide burndown control of certain small broadleaf weeds and annual grasses. To obtain burndown control of the species listed below:

Annual Grasses	Pigweeds:
Chickweed	Palmer amaranth
Dandelion	Redroot
Henbit	Smooth
Lettuce, prickly	Spiny amaranth
Lambsquarters	Tall waterhemp
Marestail	Ragweed, common
Mustard, wild	Ragweed, giant
Pennycress	Shepherd's-purse
	Smartweeds, annual

- o The broadleaf weeds listed above should not exceed 3" in height, the annual grasses should not exceed 2" in height. To control weeds exceeding these sizes, CANOPY + NEW LOROX PLUS may be tank mixed with Roundup¹, Bronco², Gramoxone³ Extra, or 2,4-D (LVE).
- o Always include crop oil concentrate at 1% v/v (1 gallon per 100 gallons of spray mix.) When tank mixing with "Roundup" use nonionic surfactant at 0.25% v/v (1 quart per 100 gallons of spray mix).
- o Use flat fan nozzles and a minimum of 20 gallons of water per acre.

Grass Control:

These tank mixes may not provide adequate preemergence control of grasses. For grass control they may be:

- o Followed, as needed, by a postemergence grass herbicide such as DuPont ASSURE[®] II Herbicide.
- o Tank mixed with Lasso⁴ (or other products containing the active ingredient alachlor and registered for use on soybeans), "Bronco", Dual⁵, Prowl⁶, Command⁷, Treflan⁸ (or other products containing the active ingredient trifluralin and registered for use on soybeans, or Sonalan⁹).

Refer to the label of the tank mix partner or the post grass herbicide for specific information regarding use, rates, pre-plant intervals, application timing, grasses controlled, precautions, restrictions and other information.

No-Till Burndown:

These tank mixes, when mixed with crop oil concentrate, will control (burndown) many small weeds. However, to burndown those weeds listed on this label which exceed 2-3" or to burndown other species not listed, they may be tank mixed with one of the following products:

"Roundup"
"Bronco"
"Gramoxone" Extra
2,4-D LVE*

To select the proper burndown product, identify the weeds which need to be controlled and consult the product labels to determine which product is needed. Consult the companion burndown herbicide label for use instructions, rates, precautions, restrictions and other use information.

*2,4-D LVE is the isooctyl (2-ethylhexyl) ester of 2,4-Dichlorophenoxyacetic acid. This product is sold under a variety of trade names. It has a minimum preplant interval of 7 to 30 days and is based on the rate used. Consult the label of the product used for specific information on this interval.

IMPORTANT

BEFORE USING "CANOPY", READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

¹ Trademark of Monsanto Company

² Trademark of Zeneca

³ Trademark of Ciba-Geigy Corporation

⁴ Trademark of American Cyanamid Company

⁵ Trademark of FMC Corporation

⁶ Trademark of DowElanco

D-061995/072595/072695

