

PM 25


352-444

Page 1 of 2



# DuPont Agricultural Products

..... A Growing Partnership With Nature™



## SUPPLEMENTAL LABELING

CANOPY® HERBICIDE  
FALL APPLICATIONS

REC'D EPA OPP/EDD1  
'96 SEP 19 P 1034

### CANOPY® HERBICIDE

(EPA REG. NO. 352-444)

FALL APPLICATIONS TO BE USED IN NO-TILL SOYBEAN PRODUCTION IN THE STATES OF ILLINOIS, INDIANA, KANSAS, MISSOURI, OHIO AND PENNSYLVANIA

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont CANOPY Herbicide is recommended for fall burndown of existing vegetation and preemergence broadleaf control the following spring in no-till soybean production systems in the states of Illinois, Indiana, Kansas, Missouri, Ohio, and Pennsylvania.

For use in soybeans only and in soils with a composite pH of 6.8 or less.

Do not apply this product through an irrigation system.

Do not graze treated fields or harvest forage for hay.

#### Application Timing

CANOPY can be applied to no-till fields in the fall between November 1 and December 31 and when soil temperatures have dropped below 50° F in the top 1" of soil. DO NOT APPLY TO FROZEN GROUND. If a fall application of CANOPY is used, do not apply CANOPY again in the spring.

#### Postemergence or Burndown control of existing fall weeds

CANOPY applied in the fall will provide burndown and subsequent spring control of certain broadleaf weeds and annual grasses. To obtain burndown of the species listed below, always use CANOPY in 1% COC.

- |                       |                        |
|-----------------------|------------------------|
| Winter annual grasses | Mustard, wild          |
| Chickweed             | Pennycress             |
| Dandelion             | Thistle, Canadian      |
| Henbits               | Shepard's-purse        |
| Lettuce, prickly      | Winter annual mustards |
| Marestail (horseweed) |                        |

The broadleaf weeds listed above should not exceed 3" in height; the annual grasses should not exceed 2" in height. To burndown weeds exceeding these heights, CANOPY may be tank mixed, with Roundup<sup>1</sup>, Gramoxone<sup>2</sup> Extra, Bronco<sup>1</sup>, or 2,4-D LVE. Always include crop oil concentrate at V/V (1 gallon per 100 gallons of spray mix). When tank mixing with "Roundup" or "Bronco", use a nonionic surfactant at 0.25% V/V (1 quart per 100 gallons of spray mix). To select the proper burndown product, identify the weeds that need to be controlled and consult the product labels to determine which product is needed. Consult the companion burndown herbicide for use instructions, rates, precautions, restrictions and other use information.

#### For Improved Grass Control

CANOPY applied in the fall may not provide adequate season-long preemergence control of annual grasses. For improved season-long grass control, CANOPY may be:

- followed, as needed, by a spring-applied postemergence grass herbicide such as DuPont Assure® II Herbicide.
- followed, as needed, by a spring-applied preemergence grass herbicide such as Lasso<sup>1</sup>, Dual<sup>3</sup>, Dual<sup>3</sup> II, Prowl<sup>4</sup>, Command<sup>5</sup>, or Frontier<sup>6</sup>.

Refer to the preemergence or postemergence grass herbicide for specific information regarding use rates, pre-plant intervals, application timing, grasses controlled, precautions, restrictions, and other information.

#### Application Equipment and Spray Volumes

Apply uniformly by ground equipment using a properly calibrated fixed boom sprayer. Use 10-40 gallons per acre at 20-40 psi. For postemergence or burndown applications over existing vegetation, use flat fan nozzles and a minimum of 20 gallons per acre.

292

**Application Rates**

Select the rate of CANOPY from the chart below, according to soil type.

Soil Texture	Broadcast Rate
<b>1.5 - 4% Organic Matter</b>	
<b>Medium:</b>	
Loam, Silt Loam, Silt, Silty Clay Loam	5 - 6 oz
<b>Fine:</b>	
Silty Clay Loam, Clay Loam, Clay	5 - 7 oz

**Medium:**

**Fine:**

**IMPORTANT**

**BEFORE USING "CANOPY", READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.**

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

**WEEDS CONTROLLED**

**Preemergence**

When used according to this label, fall-applied CANOPY will generally provide extended preemergence control of the weeds listed below. Subsequent flushes the following spring after soybean emergence may require an application of a postemergence herbicide for season-long control.

<sup>1</sup> Registered trademark of Monsanto Company

<sup>2</sup> Registered trademark of Zeneca Inc.

<sup>3</sup> Registered trademark of Ciba-Geigy Corp.

<sup>4</sup> Registered trademark of American Cyanamid Company

<sup>5</sup> Registered trademark of FMC Corp.

<sup>6</sup> Registered trademark of Sandoz Agro, Inc.

Cocklebur	Ragweed, common
Lambsquarters	Smartweeds, annual
Pigweeds, redroot and smooth	Speedwell, purslane
	Velvetleaf

Fall-applied CANOPY will provide partial control (suppression) preemergence of the following weeds:

Crabgrass	Nutsedge, yellow
Foxtail, species	Panicum, fall
Jimsonweed	Ragweed, giant
Morningglories, entireleaf, ivyleaf, tall	Sida, prickly (teaweed)

D-091096/3  
(Replaces H-62758)

