




DuPont Agricultural  
Products

"..... A Growing Partnership With Nature"



SUPPLEMENTAL LABELING

CANOPY HERBICIDE

PMJS 352-444 11-17-97 1044  
Accepted stamp on p8. Jaf2

## CANOPY® HERBICIDE

Reg. No. 352-444

### CANOPY APPLIED REDUCED RATES FOR BURNDOWN AND LIMITED RESIDUAL CONTROL ON VARIABLE pH SOILS

Canopy at 2 ounces per acre can be used in soybean production for burndown and limited residual weed control regardless of soil type and pH. Canopy at 2 ounces per acre is recommended for use in the states of Illinois north of Route 36, Indiana north of I-70, Iowa outside the Nicollett-Clarrion-Webster and Hamburg-Ida-Monana soil associations, Missouri north of Route 36, Ohio north of I-70, and Pennsylvania north of I-70.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labelling.

- Before using Canopy, read and carefully observe the cautionary statements and all other information appearing on the Canopy Specimen Label.
- Do not apply this product through an irrigation system.
- For use on soybeans only.
- Do not graze treated fields or harvest forage for hay.

#### APPLICATION RATES AND ADJUVANTS

Canopy applied at 2 ounces per acre is recommended for burndown of existing summer and winter annuals and limited preemergence control of certain broadleaf weeds (see Weeds Controlled list below). For burndown, always include a Crop Oil Concentrate at 1% v/v (1 gallon per 100 gallons of finished spray).

#### APPLICATION TIMING

Canopy at 2 ounces can be applied up to 45 days before planting soybeans, at-planting, or immediately after planting but prior to soybean emergence.

#### APPLICATION EQUIPMENT SPRAY VOLUMES

Apply uniformly by ground equipment using a properly calibrated fixed boom sprayer. Use 10-40 gallons per acre at 20-40 psi. For burndown applications of existing vegetation, use flat fan nozzles and a minimum of 20 gallons per acre.

#### WEEDS CONTROLLED - BURNDOWN

Burndown control of existing winter and summer annuals with 2 oz Canopy applied prior to planting will provide burndown control of certain 1-3" annual broadleaf weeds and winter annuals. To obtain burndown of the species listed below, use Canopy plus 1% COC. For enhanced burndown control, 1 pint of 2,4-D LVE (0.5 lbs a.e./A) may be tank mixed with Canopy and applied from 45 days up to 7 days prior to planting.

Annual grasses	Marestail	Shepherd's purse
Chickweed* annual	Mustard, wild	Smartweeds,
Dandelion	Pennycress	Sunflower
Henbits	Pigweeds	Velvetleaf
Lambsquarters*	Ragweed, Giant	Wild Garlic*
Lettuce, Prickly	Ragweed, Common	

\* always add 1 pt 2,4-D LVE

The annual grasses should not exceed 1" in height. For burndown of annual grasses taller than 1", add 2.5 - 5.0 oz Assure II. The broadleaf weeds should not exceed 3" in height. To burndown weeds exceeding 3", 2 oz Canopy may be tankmixed with Roundup Ultra<sup>1</sup>, Gramoxone Extra<sup>2</sup>, Bronco<sup>1</sup>, or 2,4-D LVE. With Canopy plus Gramoxone Extra or 2,4-D LVE tankmixes, always include crop oil concentrate at 1% V/V (1 gallon per 100 gallons of spray mix). When tank mixing with Roundup Ultra or

Bronco, refer to these product labels for specific adjuvant recommendations, however to optimize burndown from the Canopy, add 0.25% NIS when tankmixing Canopy with Roundup Ultra or Bronco. To select the proper burndown product, identify the weeds that need to be controlled and consult the product labels to determine which product is needed. Consult the companion burndown herbicide for use instructions, rates, precautions, restrictions and other use information.

**PREEMERGENCE**

When used according to this label, 2 oz of Canopy will provide limited preemergence control of the weeds below to contribute to a clean seed bed at planting. For season-long control, a planned PRE or POST sequential program is required.

Pigweeds, Redroot and smooth

Ragweed, Common

Smartweeds, annual

2 oz Canopy will provide partial preemergence control \* (suppression) of the following weeds:

Cocklebur Ragweed, Giant

Jimsonweed Foxtail, species

Lambsquarters Fall Panicum

Morningglories,  
annual,  
ivyleaf,  
tall

Nutsedge, yellow Velvetleaf

\*Due to heavy weed pressure, delayed planting, or adverse environmental conditions, additional burndown control measures may be required at planting.

For enhanced residual control, 2-4 oz Lexone may be tankmixed with 2 oz Canopy.

For Season-long Grass and Broadleaf Control 2 oz Canopy will not provide season-long preemergence control of annual grasses and broadleaf weeds.

For season-long control, 2 oz Canopy may be:

- used on soils with pH's above 6.8, follow as needed, by postemergence grass and/or broadleaf herbicides such as Dupont Assure II, or Pinnacle herbicides. Do not apply any additional CE containing herbicides within the same year if pH's exceed 6.8.

- used on soils with pH less than 6.8, follow as needed, by sequential applications of Classic, Synchrony STS,

Concert, or Pinnacle. Refer to the Classic, Synchrony STS, Concert, and Pinnacle labels for all claims, use directions, and restrictions associated with these products.

- tank mixed or followed as needed with a preemergence grass herbicide such as Lasso<sup>1</sup>, Dual<sup>2</sup>,

Dual II<sup>3</sup>, Prowl<sup>4</sup>, Command<sup>5</sup>, or Frontier<sup>6</sup>.

- used alone or in a tank mix with Roundup Ultra<sup>7</sup> as a burndown, and followed by Roundup Ultra in Roundup Ready<sup>8</sup> soybeans.

Refer to the preemergence or postemergence herbicide labels for specific information regarding use rates, pre-plant intervals, application timing, grasses controlled, precautions, restrictions, and other information.

**IMPORTANT**

Before using Canopy, read and follow all applicable directions, restrictions and precautions on the EPA-Registered label.

Follow rotational crop interval recommendations on the Canopy EPA-Registered label. Before using Canopy, read and carefully observe the cautionary statements and all other information appearing on the Canopy label. For recropping information after 2 oz. Canopy, use rotational guideline 1 on the Canopy label.

This label contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

<sup>1</sup>Trademark of Monsanto Agricultural Products, Co.

<sup>2</sup>Trademark of ICI Americas Corp.

<sup>3</sup>Trademark of Novartis Corp.


<sup>4</sup>Trademark of Sandoz Chemicals, Inc.

<sup>5</sup>Trademark of FMC Corp.

<sup>6</sup>Trademark of American Cyanamid Corp.

<sup>7</sup>Trademark of Monsanto Agricultural Products Co.

<sup>8</sup>Trademark of Monsanto Agricultural Products Co.



D-092597

PM 25


352-444

11-17-97

314



DuPont Agricultural Products



SUPPLEMENTAL LABELING

CANOPY HERBICIDE

"..... A Growing Partnership With Nature"

CANOPY® HERBICIDE

Reg. No. 352-444

CANOPY HERBICIDE TANKMIXES WITH SCEPTER<sup>1</sup> DG OR SQUADRON<sup>2</sup> FOR BROADLEAF WEED CONTROL IN THE STATES OF INDIANA, MICHIGAN, OHIO, AND PENNSYLVANIA

DuPont Canopy herbicide is recommended at a reduced rate for preemergence broadleaf weed control when tankmixed with a reduced rate of either Scepter DG or Squadron in the states of Indiana, Michigan, Ohio and Pennsylvania.

APPLICATION RATES

Broadcast Rate (ounces or pints per acre)
1/2 - 4% Organic Matter

Table with columns: Soil texture, Canopy (oz), Scepter DG (oz), Squadron (pt). Rows include Coarse, Loamy sand, sandy loam, Medium or Fine, loam, silt loam, silt, sandy clay loam, silty clay loam, clay.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

For use on soybeans only.

Do not apply this product through any type of irrigation equipment.

Do not graze treated fields or harvest forage for hay.

APPLICATION TIMING AND METHODS

Canopy tankmixed with either Scepter DG or Squadron may be applied at planting or at any time up to 45 days prior to planting. Do not apply after the soybean crop has emerged. These tankmixes may be preemergence surface applied or preplant incorporated. Refer to the Canopy label for specific instructions on incorporation and moisture requirements. These tankmixes may also be applied over existing weed vegetation as in no-till soybean production.


APPLICATION EQUIPMENT AND SPRAY VOLUMES

Apply uniformly by ground equipment using a properly calibrated fixed boom sprayer. Use 10-40 gallons per acre at 20-40 psi. In no-till situations, where these tankmixes are being applied over existing vegetation, use flat fan nozzles and a minimum of 20 gallons per acre.

WEEDS CONTROLLED - PREEMERGENCE

When used according to this label, these Canopy tankmixes will generally provide season-long preemergence control of the following weeds:

- Cocklebur\*
Jimsonweed
Lambsquarters
Nightshade, Eastern Black
Pigweed, Redroot and smartweed
Ragweed, Common and Giant
Velvetleaf
Smartweed, annual



These Canopy tankmixes will also provide partial preemergence control of the following weeds:

- Annual grasses
Morningglories, entireleaf, ivyleaf\*

\* Large-seeded weeds germinating deep in the soil such as cocklebur, giant ragweed, and Morningglories or other weeds which may emerge at various times during the growing season may require a cultivation or an application of a postemergent herbicide such as DuPont Classic® herbicide for season-long control.

**POSTEMERGENCE (FOR NO-TILL BURNDOWN)**

These Canopy tankmixes will provide burndown control of the weeds listed below.

To obtain burndown control of these weeds:

- Always include crop oil concentrate at 1% v/v (1 gallon per 100 gallons of spray mix). When tankmixing with Roundup Ultra<sup>3</sup>, use nonionic surfactant at 0.25% v/v (1 quart per 100 gallons of spray mix).
- Make applications before broadleaf weeds exceed 3" in height and before annual grasses exceed 1" in height. To control weeds exceeding these sizes, Canopy + Scepter DG or Squadron may be tankmixed with Roundup Ultra, Bronco<sup>4</sup>, Gramoxone Extra<sup>5</sup>, or 2,4-D (LVE).

annual grasses	pigweeds, redroot and smooth
chickweed	ragweed, common
dandelion	ragweed, giant
henbit	shepherd's purse
lambsquarters**	smartweeds, annual
lettuce, prickly	velvetleaf
maretail	wild garlic**
mustard, wild	
nightshade, E.B.**	
pennycress	

\*\* always add 1 pt 2,4-D LVE

**FOR IMPROVED GRASS CONTROL**

The Canopy + Scepter or Squadron tankmixes may not provide adequate preemergence control of grasses. For improved grass control:

- follow, as needed, with a postemergence grass herbicide such as DuPont Assure<sup>®</sup> II herbicide.
- tankmix with a preemergence grass control herbicide registered for use in soybeans.

When tankmixing Canopy + Scepter or Squadron with another tankmix partner, refer to the label of the tankmix partner for specific information regarding use, rates, pre-plant intervals, application timing, grasses controlled, precautions, restrictions and other information.

**IMPORTANT**

- Before using Canopy, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label.
- This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow instructions carefully.
- This labeling must be in the possession of the user at the time of pesticide application.

<sup>1</sup>Trademark of American Cyanamid Corp.

<sup>2</sup>Trademark of American Cyanamid Corp.

<sup>3</sup>Trademark of Monsanto Agricultural Products Co.

<sup>4</sup>Trademark of Monsanto Agricultural Products Co.

<sup>5</sup>Trademark of ICI Americas Corp.

D-101497