




**DuPont Agricultural
Products**

"..... A Growing Partnership With Nature"



SUPPLEMENTAL LABELING

**CANOPY® HERBICIDE
REDUCED RATES FOR BURNDOWN
AND LIMITED RESIDUAL CONTROL
ON VARIABLE pH SOILS
IN THE STATE OF MICHIGAN**

CANOPY® HERBICIDE

Reg. No. 352-444

CANOPY APPLIED REDUCED RATES FOR BURNDOWN AND LIMITED RESIDUAL CONTROL ON VARIABLE pH SOILS IN THE STATE OF MI

Canopy at 2 ounces per acre can be used in soybean production for burndown and limited residual weed control in the State of Michigan on soils with pH of 7.6 or less.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labelling.

- Before using Canopy, read and carefully observe the cautionary statements and all other information appearing on the Canopy Specimen Label.
- Do not apply this product through an irrigation system.
- For use on soybeans only.
- Do not graze treated fields or harvest forage for hay.

APPLICATION RATES AND ADJUVANTS

Canopy applied at 2 ounces per acre is recommended for burndown of existing summer and winter annuals and limited pre-emergence control of certain broadleaf weeds (see Weeds Controlled list below). For burndown, always include a Crop Oil Concentrate at 1% v/v (1 gallon per 100 gallons of finished spray).

APPLICATION TIMING

Canopy at 2 ounces can be applied up to 45 days before planting soybeans, at-planting, or immediately after planting but prior to soybean emergence.

APPLICATION EQUIPMENT SPRAY VOLUMES

Apply uniformly by ground equipment using a properly calibrated fixed boom sprayer. Use 10-40 gallons per acre at 20-40 psi. For burndown applications of existing vegetation, use flat fan nozzles and a minimum of 20 gallons per acre.


WEEDS CONTROLLED - BURNDOWN

Burndown control of existing winter and summer annuals with 2 oz Canopy applied prior to planting will provide burndown control of certain 1-3" annual broadleaf weeds and winter annuals. To obtain burndown of the species listed below, use Canopy plus 1% COC. For enhanced burndown control, 1 pint of 2,4-D LVE (0.5 lbs a.e./A) may be tank mixed with Canopy and applied from 45 days up to 7 days prior to planting.

Annual grasses	Marestail	Shepherd's purse
Chickweed*	Mustard, wild	Smartweeds, annual
Dandelion	Pennycress	Sunflower
Henbits	Pigweeds	Velvetleaf
Lambsquarters*	Ragweed, Giant	Wild Garlic*
Lettuce, Prickly	Ragweed, Common	

* always add 1 pt 2,4-D LVE

The annual grasses should not exceed 1" in height. For burndown of annual grasses taller than 1", add 2.5 - 5.0 oz Assure II. The broadleaf weeds should not exceed 3" in height. To burndown weeds exceeding 3", 2 oz Canopy may be tankmixed with Roundup Ultra¹, Gramoxone Extra², Bronco¹, or 2,4-D LVE. With Canopy plus Gramoxone Extra or 2,4-D LVE tankmixes, always include crop oil concentrate at 1% V/V (1 gallon per 100 gallons of spray mix). When tank mixing with Roundup Ultra or Bronco, refer to these product labels for specific adjuvant recommendations, however to optimize burndown from the Canopy, add 0.25% NIS when tankmixing Canopy with Roundup Ultra or Bronco. To select the proper burndown product, identify the weeds that need to be controlled and consult the product labels to determine which product is needed. Consult the companion burndown herbicide for use instructions, rates, precautions, restrictions and other use information.



PREEMERGENCE

When used according to this label, 2 oz of Canopy will provide limited preemergence control of the weeds below to contribute to a clean seed bed at planting. For season-long control, a planned PRE or POST sequential program is required.

- Pigweeds, Redroot and smooth
- Ragweed, Common
- Smartweeds, annual

2 oz Canopy will provide partial preemergence control * (suppression) of the following weeds:

- | | |
|------------------|------------------|
| Cocklebur | Ragweed, Giant |
| Jimsonweed | Foxtail, species |
| Lambsquarters | Fall Panicum |
| Morningglories, | |
| annual, | |
| ivyleaf, | |
| tall | |
| Nutsedge, yellow | Velvetleaf |

*Due to heavy weed pressure, delayed planting, or adverse environmental conditions, additional burndown control measures may be required at planting.

For enhanced residual control, 2-4 oz Lexone DF or 8-16 oz of Lorox DF may be tankmixed with 2 oz Canopy.

Canopy at 2 oz rate will not provide season-long preemergence control of annual grasses and broadleaf weeds.

For season-long control, 2 oz Canopy may be:

- used on soils with pH's between 6.8 and 7.6, and follow as needed, by postemergence grass and/or broadleaf herbicides such as Dupont Assure II, or Pinnacle herbicides. Do not apply Classic, Synchrony STS, or Reliance STS within the same year if pH's exceed 6.8.
- used on soils with pH less than 6.8, follow as needed, by sequential applications of Classic, Synchrony STS, Reliance STS, or Pinnacle. Refer to the Classic, Synchrony STS, Reliance STS and Pinnacle labels for all claims, use directions, and restrictions associated with these products.
- tank mixed or followed as needed with a preemergence grass herbicide such as Lasso¹, Dual², Dual II³, Prowl⁶, Command⁵, or Frontier⁴.
- used alone or in a tank mix with Roundup Ultra⁷ as a burn-down, and followed by Roundup Ultra in Roundup Ready⁸ soybeans.

Refer to the preemergence or postemergence herbicide labels for specific information regarding use rates, pre-plant intervals, application timing, grasses controlled, precautions, restrictions, and other information.

IMPORTANT

Before using Canopy, read and follow all applicable directions, restrictions and precautions on the EPA-Registered label.

Follow rotational crop interval recommendations on the Canopy EPA-Registered label. Before using Canopy, read and carefully observe the cautionary statements and all other information appearing on the Canopy label. For recropping information after 2 oz. Canopy, use rotational guideline 1 on the Canopy label.

This label contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

- ¹Trademark of Monsanto Agricultural Products, Co.
- ²Trademark of ICI Americas Corp.
- ³Trademark of Novartis Corp.
- ⁴Trademark of Sandoz Chemicals, Inc.
- ⁵Trademark of FMC Corp.
- ⁶Trademark of American Cyanamid Corp.
- ⁷Trademark of Monsanto Agricultural Products Co.
- ⁸Trademark of Monsanto Agricultural Products Co.

