




DuPont Agricultural  
Products

"..... A Growing Partnership With Nature"



SUPPLEMENTAL LABELING

CANOPY® HERBICIDE  
EXPANDED APPLICATION  
TIMING

## CANOPY® HERBICIDE

EPA Reg. No. 352-444

### EXPANDED APPLICATION TIMING FOR USE IN NO-TILL OR CONSERVATION TILLAGE\* SOYBEAN PRODUCTION IN THE STATES OF INDIANA, ILLINOIS, KANSAS, KENTUCKY, MISSOURI, OHIO, PENNSYLVANIA, AND TENNESSEE

Dupont Canopy herbicide is recommended for fall burndown of existing vegetation and preemergence broadleaf control the following spring in no-till and conservation tillage soybean production systems in the states of Indiana, Illinois, Kansas, Kentucky, Missouri, Ohio, Pennsylvania, and Tennessee.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through an irrigation system.

For use in soybeans only and in soils with a composite pH of 6.8 or less.

Do not graze treated fields or harvest forage for hay.

#### APPLICATION TIMING

Canopy can be applied to No-till or conservation tillage fields after soil temperatures have dropped below 50° F in the top 1" of soil. **DO NOT APPLY TO FROZEN GROUND.** Split applications of Canopy may be used as long as the full labeled rate for the geography is not exceeded.

#### APPLICATION EQUIPMENT AND SPRAY VOLUME

Apply uniformly by ground equipment using a properly calibrated fixed boom sprayer. Use 10-40 gallons per acre at 20-40 psi. For postemergence or burndown applications over existing vegetation, use flat fan nozzles and a minimum of 20 gallons per acre.

\* Conservation Tillage is defined as greater than or equal to 30% surface residue at the time of Canopy application

#### APPLICATION RATES

Select a Broadcast rate of Canopy from the chart below, according to soil type and length of residual control desired.

Soil Texture  
(1.5 - 4% organic matter)

	<u>Full season Rate</u>	<u>Planned sequential Rate</u>
<b>Medium:</b>		
Loam,		
Silt Loam,		
Silt,		
Sandy Clay Loam	5 - 6 oz	3 oz
<b>Fine:</b>		
Silty Clay Loam,		
Clay Loam,		
Clay	5 - 7 oz	3 - 4 oz

#### WEEDS CONTROLLED

##### Preemergence

When used according to this label, Fall-applied 5-7 oz Canopy will generally provide season-long preemergence control of the following weeds.

Fall-applied 3-4 oz Canopy will generally provide acceptable preemergence control of the following weeds through 2-3 weeks after planting.

For season-long control following 3-4 oz Canopy, a planned PRE or POST sequential program is required.

Lambsquarters	Ragweeds, Common
Smartweeds, annual	Velvetleaf
Pigweeds, Redroot and smooth	Purselane Speedwell
Cocklebur **	Marestail

For product information call 1-888-6-DUPONT

© 1998 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898

Fall-applied Canopy will provide partial control \*\* (suppression) preemergence of the following weeds:

- |                  |                             |
|------------------|-----------------------------|
| Ragweed, Giant   | Jimsonweed                  |
| Foxtail, species | Picky Sida (teaweed)        |
| Yellow Nutsedge  | Morningglories,             |
|                  | (Entireleaf, Ivyleaf, Tall) |
| Crabgrass        | Fall Panicum                |

\*\* Subsequent flushes the following spring after soybean emergence may require an application of a postemergence herbicide for season-long control.

**Postemergence or Burndown Control of Existing Fall Weeds**  
Canopy applied in the fall will provide burndown and subsequent spring control of certain broadleaf weeds and annual grasses.

To obtain burndown of the species listed below, always use Canopy in 1% COC. A pint of 2,4-D may be added for extra control.

- |                  |                                   |
|------------------|-----------------------------------|
| Annual grasses   | Marestail                         |
| Chickweed        | Mustards                          |
| Dandelion        | (Pennycress, Pepperweed)          |
| Henbits          | Shepherdspurse, Bushy Wallflower) |
| Lettuce, prickly | Thistle, Canadian                 |

The broadleaf weeds listed above should not exceed 3" in height; the annual grasses should not exceed 1" in height. To burndown weeds exceeding these heights, Canopy may be tank mixed with Roundup<sup>1</sup> Ultra, Gramoxone<sup>2</sup> Extra, Bronco<sup>3</sup>, or 2,4-D LVE. Always include crop oil concentrate at 1% V/V (1 gallon

per 100 gallons of spray mix). When tank mixing with "Roundup Ultra" or "Bronco", use a nonionic surfactant at 0.25% V/V (1 quart per 100 gallons of spray mix). To select the proper burndown product, identify the weeds that need to be controlled and consult the product labels to determine which product is needed. Consult the companion burndown herbicide for use instructions, rates, precautions, restrictions and other use information.

**For Improved Grass Control**

Canopy applied in the Fall may not provide adequate season-long preemergence control of annual grasses. For improved season-long grass control, Canopy may be:

- followed, as needed, by a spring-applied postemergence grass herbicide such as Dupont Assure<sup>®</sup> II herbicide.
- followed, as needed, by a spring-applied preemergence grass herbicide such as Lasso<sup>1</sup>, Dual<sup>3</sup>, Dual<sup>3</sup> II, Prowl<sup>4</sup>, Command<sup>5</sup>, or Frontier<sup>6</sup>.


Refer to the preemergence or postemergence grass herbicide for specific information regarding use rates, pre-plant intervals, application timing, grasses controlled, precautions, restrictions, and other information.

**IMPORTANT**  
**BEFORE USING "CANOPY", READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.**

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

<sup>1</sup> Registered trademark of Monsanto Agricultural Products, Co.  
<sup>2</sup> Registered trademark of Zeneca, Inc.  
<sup>3</sup> Registered trademark of Novartis  
<sup>4</sup> Registered trademark of American Cyanamid Co.  
<sup>5</sup> Registered trademark of FMC Corporation.  
<sup>6</sup> Registered trademark of Sandoz Agro, Inc.



(Replaces H-63025)

D-071498

For product information call 1-888-6-DUPONT

© 1998 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898