



PM 25


352-444

11-3-98

SUPPLEMENTAL LABELING

104-2

DuPont Agricultural Products



CANOPY HERBICIDE

"..... A Growing Partnership With Nature"

CANOPY® HERBICIDE

Reg. No.352-444

CANOPY APPLIED REDUCED RATES FOR BURNDOWN AND LIMITED RESIDUAL CONTROL ON VARIABLE pH SOILS

Canopy at 2 ounces per acre can be used in soybean production for burndown and limited residual weed control regardless of soil type and pH. Canopy at 2 ounces per acre is recommended for use in the states of Illinois north of Route 36, Indiana north of I-70, Iowa outside the Nicollett-Clarrion-Webster and Hamburg-Ida-Monana soil associations, Missouri north of Route 36, Ohio north of I-70, and Pennsylvania north of I-70.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labelling.

- Before using Canopy, read and carefully observe the cautionary statements and all other information appearing on the Canopy Specimen Label.
• Do not apply this product through an irrigation system.
• For use on soybeans only.
• Do not graze treated fields or harvest forage for hay.

APPLICATION RATES AND ADJUVANTS

Canopy applied at 2 ounces per acre is recommended for burndown of existing summer and winter annuals and limited pre-emergence control of certain broadleaf weeds (see Weeds Controlled list below). For burndown, always include a Crop Oil Concentrate at 1% v/v (1 gallon per 100 gallons of finished spray).

APPLICATION TIMING

Canopy at 2 ounces can be applied before planting soybeans, at planting, or immediately after planting but prior to soybean emergence. DO NOT APPLY TO FROZEN GROUND.

APPLICATION EQUIPMENT SPRAY VOLUMES

Apply uniformly by ground equipment using a properly calibrated fixed boom sprayer. Use 10-40 gallons per acre at 20-40 psi. For burndown applications of existing vegetation, use flat fan nozzles and a minimum of 20 gallons per acre.

WEEDS CONTROLLED - BURNDOWN

Burndown control of existing winter and summer annuals with 2 oz Canopy applied prior to planting will provide burndown control of certain 1-3" annual broadleaf weeds and winter annuals. To obtain burndown of the species listed below, use Canopy plus 1% COC. For enhanced burndown control, 1 pint of 2,4-D LVE (0.5 lbs a.e./A) may be tank mixed with Canopy and applied from 25 days up to 7 days prior to planting.

Table listing weeds controlled: Annual grasses, Chickweed**, Dandelion, Henbits, Lambsquarters*, Lettuce, Prickly, Maretail, Mustard, wild, Pennycress, Pigweeds, Ragweed, Giant, Ragweed, Common, Shepherd's purse, Smartweeds, annual, Sunflower, Velvetleaf, Wild Garlic*.

* always add 1 pt 2,4-D LVE

** For enhanced burndown of chickweed, Harmony Extra at 0.5-0.6 oz/A may be tankmixed with Canopy. Apply Harmony Extra at least 45 days prior to planting soybeans.

The annual grasses should not exceed 1" in height. For burndown of annual grasses taller than 1", add 2.5 - 5.0 oz Assure II. The broadleaf weeds should not exceed 3" in height. To burndown weeds exceeding 3", 2 oz Canopy may be tankmixed with Roundup Ultra, Touchdown, Gramoxone Extra, Bronco, or 2,4-D LVE. With Canopy plus Gramoxone Extra or 2,4-D LVE tankmixes, always include crop oil concentrate at 1% V/V (1 gallon per 100 gallons of spray mix). When tank mixing with Roundup Ultra, Touchdown, or Bronco, refer to these product labels for specific adjuvant recommendations, however to optimize burndown from the Canopy, add 0.25% NIS when tankmixing Canopy with Roundup Ultra, Touchdown, or Bronco. To select the proper burndown product, identify the weeds that need to be controlled and consult the product labels to determine which product is needed. Consult the companion burndown herbicide for use instructions, rates, precautions, restrictions and other use information.

For product information call 1-888-6-DUPONT

© 1998 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898

2 of 2

PREEMERGENCE

When used according to this label, 2 oz of Canopy will provide limited preemergence control of the weeds below to contribute to a clean seed bed at planting. For season-long control, a planned PRE or POST sequential program is required.

- Pigweeds, Redroot and smooth
- Ragweed, Common
- Smartweeds, annual

2 oz Canopy will provide partial preemergence control * (see expression) of the following weeds:

- | | |
|------------------|------------------|
| Cocklebur | Ragweed, Giant |
| Jimsonweed | Foxtail, species |
| Lambsquarters | Fall Panicum |
| Morningglories, | Velvetleaf |
| annual, | |
| ivyleaf, | |
| tall | |
| Nutsedge, yellow | |

*Due to heavy weed pressure, delayed planting, or adverse environmental conditions, additional burndown control measures may be required at planting.

For enhanced residual control, 2-4 oz Lexone or Senecor¹ may be tankmixed with 2 oz Canopy.

For Season-long Grass and Broadleaf Control 2 oz Canopy will not provide season-long preemergence control of annual grasses and broadleaf weeds.

For season-long control, 2 oz Canopy may be:

- used on soils with pH's above 6.8, follow as needed, by post-emergence grass and/or broadleaf herbicides such as DuPont Assure II, or Pinnacle herbicides. Do not apply any additional chlorimuron ethyl, such as DuPont Classic, Reliance STS, or Synchrony STS, or Concert, containing herbicides, within the same year if pH's exceed 6.8.

- used on soils with pH less than 6.8, follow as needed, by sequential applications of Classic, Synchrony STS, Concert, or Pinnacle. Refer to the Classic, Synchrony STS, Concert, and Pinnacle labels for all claims, use directions, and restrictions associated with these products.

- tank mixed or followed as needed with a preemergence grass herbicide such as Lasso¹, Dual², Dual II², Prowl³, Command⁴, or Frontier⁵.

- used alone or in a tank mix with Roundup Ultra⁶, or Touchdown as a burndown, and followed by Roundup Ultra or Roundup Ready⁶ soybeans.

Refer to the preemergence or postemergence herbicide labels for specific information regarding use rates, pre-plant intervals, application timing, grasses controlled, precautions, restrictions, and other information.

IMPORTANT

Before using Canopy, read and follow all applicable directions, restrictions and precautions on the EPA-Registered label.

Follow rotational crop interval recommendations on the Canopy EPA-Registered label. Before using Canopy, read and carefully observe the cautionary statements and all other information appearing on the Canopy label. For recropping information after 2 oz. Canopy, use rotational guideline 1 on the Canopy label.

This label contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

¹Trademark of Monsanto Agricultural Products, Co.

²Trademark of Zeneca, Inc.

³Trademark of Novartis Corp.

⁴Trademark of Sandoz Chemicals, Inc.


⁵Trademark of FMC Corp.

⁶Trademark of American Cyanamid Corp.

⁷Trademark of Bayer AG

(Replaces H-63283)

D-091198



For product information call 1-888-6-DUPONT

© 1998 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898