



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

December 19, 2023

Terri Moss
Regulatory Specialist
Corteva Agriscience LLC
9330 Zionsville Road
Indianapolis, IN 46268

Subject: Notification per PRN 98-10 – Minor label revisions
Product Name: Canopy
EPA Registration Number: 352-444
Application Date: November 4, 2022
Case Number: 475893

Dear Terri Moss:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “NOTIFICATION” and placed in our records.

The primary brand name, “*Canopy*” has been made active in the product’s record.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

Page 2 of 2
EPA Reg. No. 352-444
Case No. 475893

If you have any questions, please contact Jamie Millard at (202) 566-2726 or by email at millard.jamie@epa.gov.

Sincerely,



Emily Schmid, Product Manager 25
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

NOTIFICATION

352-444

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

12/19/2023

*(Bulk/Tote Base Label/Base-label):***DuPont™ Canopy®****HERBICIDE****[Alternate Brand Name: DuPont Canopy®]**

METRIBUZIN	GROUP	5	HERBICIDE
CHLORIMURON ETHYL	GROUP	2	HERBICIDE

Dispersible Granules

Active Ingredients	By Weight
	75%
Metribuzin	
4-Amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4H)-one	64.3%
Chlorimuron Ethyl	
Ethyl 2-[[[(4-chloro-6-methoxypyrimidin-2-yl)amino]carbonyl]amino]sulfonyl]benzoate	10.7%
Other Ingredients	25%
TOTAL	100%

Keep Out of Reach of Children**CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

Precautionary Statements**Hazards to Humans and Domestic Animals**

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye irritation. Avoid breathing dust, vapor or spray mist. Avoid contact with skin, eye or clothing.

In case of contact with eyes, immediately flush with plenty of water. Get medical attention if irritation persists.

PERSONAL PROTECTIVE EQUIPMENT

Some of the material that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistant category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical resistant gloves made of any water proof material such as polyethylene or polyvinylchloride.
- Shoes plus socks.

Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Important: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "Applicators and Other Handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/ PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF SWALLOWED: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. **DO NOT** induce vomiting unless told to do so by a poison control center or doctor.

IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor

For medical emergencies involving this product, call toll free 1-800-992-5994.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate. **DO NOT** apply where/when conditions favor runoff.

Ground Water Advisory: This product contains metribuzin and chlorimuron-ethyl, chemicals which can travel (seep or leach) through soil and can contaminate ground water which may be used as drinking water. Metribuzin has been found in ground water as a result of agricultural use. Users are advised not to apply this product where the water table (ground water) is close to the surface, and where the soils are very permeable, i.e., well drained soils such as loamy sands. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

Surface Water Advisory: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as

having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this product from runoff water and sediment. Runoff of this product will be greatly reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Windblown Soil Particles: This product has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying this product if prevailing local conditions may be expected to result in off-site movement.

Non-target Organism Advisory: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by minimizing spray drift. For further guidance and instructions on how to minimize spray drift, refer to the Spray Drift Management section of this label.

PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Ensure that all operation employees accurately measure pesticides.
- Mix only enough product for the job at hand.
- Avoid overfilling of spray tank.
- **DO NOT** discharge excess material on the soil at a single spot in the field or mixing/loading station.
- Dilute and agitate excess solution and apply at labeled rates or uses.
- Avoid storage of pesticides near well sites.
- When triple-rinsing the pesticide container, be sure to add the rinsate to the spray mix.

IMPORTANT

Injury to or loss of desirable trees or vegetation may result from failure to observe the following: **DO NOT** apply or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. **DO NOT** use on lawns, walks, driveways, tennis courts or similar areas. Prevent drift of spray to desirable plants. **DO NOT** contaminate any body of water. Keep from contact with fertilizers, insecticides, fungicides and seeds during storage.

Prior to using **DuPont™ Canopy CANOPY®** herbicide, consideration should be given to crop rotation plans. Crops other than soybeans may be extremely sensitive to low concentrations of **CANOPY® Canopy** remaining in the soil the next planting season. Choice of rotation crop is restricted following application of **Canopy CANOPY®**. (See "ROTATIONAL CROP GUIDELINES" for your geographical region.)

Thoroughly clean **Canopy CANOPY®** from application equipment immediately after use and prior to spraying crops other than soybeans. Failure to remove even small amounts of **Canopy CANOPY®** from application equipment may result in injury to subsequently sprayed crops.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

STORAGE AND DISPOSAL

Pesticide Storage: Store product in original container only. **DO NOT** contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a cool, dry place.

Pesticide Disposal: **DO NOT** contaminate water, food, or feed by disposal. Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Refillable Containers: Refilling Container: Refill this container with [Canopy CANOPY®](#) containing metribuzin and chlorimuron ethyl only. **DO NOT** reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, **DO NOT** use the container, contact [DuPont-Corteva Agriscience](#) at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, **DO NOT** reuse or transport container, contact [DuPont-Corteva Agriscience](#) at the number below for instructions. Disposing of Container: **DO NOT** reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

DO NOT transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact 1-800-992-5994, day or night.

~~See back and side panels for additional precautionary statements.~~

Refer to accompanying labels for additional ~~precautions~~[precautionary information including](#), ~~complete~~ Directions for Use and Storage and Disposal.

~~THIS UNIT NOT FOR RESALE. FOR DISPENSING INTO DEDICATED CONTAINERS ONLY.~~

[Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.](#)

~~NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.~~

[In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.](#)

[Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.](#)

EPA Reg. No. 352-444

EPA Est. No. _____

[™ Trademarks of Corteva Agriscience and its affiliated companies](#)

~~® Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners~~

[Produced for
Corteva Agriscience LLC
9330 Zionsville Road](#)

Indianapolis, IN 46268 ~~Produced for~~
~~E.I du Pont de Nemours & Company~~
~~Chestnut Run Plaza~~
~~974 Centre Road~~
~~Wilmington, DE 19805~~

Nonrefillable Container

Net: _____

OR

Refillable Container

Net: _____

(Booklet Cover / Small Container Labelcover/shipping label):

DuPont™ Canopy®

HERBICIDE

[Alternate Brand Name: [DuPont Canopy®](#)]

METRIBUZIN	GROUP	5	HERBICIDE
CHLORIMURON ETHYL	GROUP	2	HERBICIDE

Dispersible Granules

Active Ingredients	By Weight
Metribuzin	75%
4-Amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4H)-one	64.3%
Chlorimuron Ethyl	
Ethyl 2-[[[(4-chloro-6-methoxypyrimidin-2-yl)amino]carbonyl]amino]sulfonyl]benzoate	10.7%
Other Ingredients	25%
TOTAL	100%

Keep Out of Reach of Children

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

~~See back and side panels for additional precautionary statements.~~

Refer to accompanying labels for additional precautionary information including, ~~complete~~ Directions for Use and Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. Before using this product, read [Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.](#)

~~NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.~~

[In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.](#)

[Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.](#)

EPA Reg. No. 352-444

EPA Est. No. _____

™ Trademarks of Corteva Agriscience and its affiliated companies

~~® Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners~~

[Produced for
Corteva Agriscience LLC
9330 Zionsville Road
Indianapolis, IN 46268](#) Produced for

~~E.I du Pont de Nemours & Company
Chestnut Run Plaza
974 Centre Road
Wilmington, DE 19805~~

Nonrefillable Container

Net: _____

OR

Refillable Container

Net: _____

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye irritation. Avoid breathing dust, vapor or spray mist. Avoid contact with skin, eye or clothing.

In case of contact with eyes, immediately flush with plenty of water. Get medical attention if irritation persists.

PERSONAL PROTECTIVE EQUIPMENT

Some of the material that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistant category selection chart.

Applicators and other handlers must wear:

Long-sleeved shirt and long pants.
Chemical resistant gloves made of any water proof material such as polyethylene or polyvinylchloride.
Shoes plus socks.

Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Important: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "Applicators and Other Handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/ PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF SWALLOWED: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. **DO NOT** induce vomiting unless told to do so by a poison control center or doctor.

IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor

For medical emergencies involving this product, call toll free 1-800-992-5994.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate. **DO NOT** apply where/when conditions favor runoff.

Ground Water Advisory: This product contains metribuzin and chlorimuron-ethyl, chemicals which can travel (seep or leach) through soil and can contaminate ground water which may be used as drinking water. Metribuzin has been found in ground water as a result of agricultural use. Users are advised not to apply this product where the water table (ground water) is close to the surface, and where the soils are very permeable, i.e., well drained soils such as loamy sands. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

Surface Water Advisory: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this product from runoff water and sediment. Runoff of this product will be greatly reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Windblown Soil Particles: This product has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying this product if prevailing local conditions may be expected to result in off-site movement.

Non-target Organism Advisory: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by minimizing spray drift. For further guidance and instructions on how to minimize spray drift, refer to the Spray Drift Management section of this label.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

DuPont™ ~~Canopy~~ [CANOPY®](#) herbicide must be used only in accordance with instructions on this label or in separately published [DuPont-Corteva Agriscience](#) instructions. ~~DuPont-Corteva Agriscience~~ will not be responsible for losses or damage resulting from the use of this product in any manner not specifically recommended by [DuPont-Corteva Agriscience](#).

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls.

Chemical resistant gloves made of any water proof material such as polyethylene or polyvinylchloride.

Shoes plus socks.

STORAGE AND DISPOSAL

Pesticide Storage: Store product in original container only. **DO NOT** contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a cool, dry place.

Pesticide Disposal: **DO NOT** contaminate water, food, or feed by disposal. Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Nonrefillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds): Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities

Nonrefillable Plastic and Metal Containers (Capacity Greater Than 50 Pounds): Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or

Shape Too Large to be Tipped, Rolled or Turned Upside Down): Nonrefillable container. **DO NOT** reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Paper or Plastic Bags, Fiber Sacks including Flexible Intermediate Bulk Containers (FIBC) or Fiber Drums With Liners: Nonrefillable container. **DO NOT** reuse or refill this container. Completely empty paper or plastic bag, fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances.

Refillable Fiber Drums With Liners: Refillable container (fiber drum only). Refilling Fiber Drum: Refill this fiber drum with Canopy containing metribuzin and chlorimuron ethyl only. **DO NOT** reuse this fiber drum for any other purpose. Cleaning before refilling is the responsibility of the refiller. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Disposing of Fiber Drum and/or Liner: **DO NOT** reuse this fiber drum for any other purpose other than refilling (see preceding). Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container. Offer the liner for recycling if available or dispose of liner in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner. To clean the fiber drum before final disposal, completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances.

All Other Refillable Containers: Refillable container. Refilling Container: Refill this container with Canopy containing metribuzin and chlorimuron ethyl only. **DO NOT** reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, **DO NOT** use the container, contact DuPont at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, **DO NOT** reuse or transport container, contact DuPont at the number below for instructions. Disposing of Container: **DO NOT** reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Outer Foil Pouches of Water Soluble Packets (WSP): Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available or, dispose of the empty outer foil pouch in the trash as long as WSP is unbroken. If the outer pouch contacts the formulated product in any way, the pouch must be triple rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer pouch as described previously.

DO NOT transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact 1-800-992-5994, day or night.

FOR USE ON SOYBEANS ONLY

DO NOT apply this product through any type of irrigation system. **DO NOT** graze treated fields or harvest for forage or hay.

Low pressure and high volume hand wand equipment is prohibited.

Single Application: **DO NOT** apply a full rate of [CanopyCANOPY®](#) more than once per soybean cropping cycle.

Split Application: Two applications totaling the fully labeled [CanopyCANOPY®](#) rate may be made per soybean cropping cycle. **DO NOT** exceed the full labeled rate for the geography.

PRODUCT INFORMATION

[CanopyCANOPY®](#) herbicide is a dispersible granule formulation to be mixed with water and sprayed for selective burndown and residual weed control in soybeans. When applied according to the instructions on this label, it will control many broadleaf weeds and provide partial control of nutsedge and annual grasses.

Residual applications of [CanopyCANOPY®](#) require rainfall or sprinkler irrigation to activate the herbicide. Degree of control and duration of effect depend on: rate used, weed spectrum, growing conditions at and following time of treatment, soil pH, texture, organic matter, moisture and precipitation.

Best residual control is obtained if [CanopyCANOPY®](#) is applied to moist soil and followed by rainfall or irrigation (~1") before weeds germinate. Several small rainfalls of less than 1/4" each are not as beneficial as one large rainfall of 1/2-1". On dry soil, more moisture is required for activation (1-2") before weed emergence. If moisture is insufficient to activate the herbicide, a rotary hoeing or shallow cultivation should be made after emergence of the crop while weeds are small enough to be controlled by mechanical means.

BIOLOGICAL ACTIVITY

[CanopyCANOPY®](#) rapidly inhibits the growth of susceptible weeds. Following application of preplant incorporation or preemergence treatment, susceptible weeds may germinate and emerge, but growth then ceases, and leaves become yellow 3-5 days after emergence. Death of leaf tissue and growing point will follow in some species while others will remain green but stunted and noncompetitive. Following a burndown application, growth of susceptible weeds ceases followed by tissue yellowing and browning and death of the growing point. [CanopyCANOPY®](#) provides partial control of some annual grasses when used preplant or preemergence but other products may be needed to ensure adequate grass control.

USE RESTRICTIONS

- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** graze treated fields or harvest for forage or hay.
- Low pressure and high volume hand wand equipment is prohibited.

- Single Application: DO NOT apply a full rate of [CanopyCANOPY®](#) more than once per acre per soybean cropping cycle.
- Split Application: A maximum of two applications, not to exceed the total fully labeled rate of [CanopyCANOPY®](#), may be made per year.
- **DO NOT** exceed the full labeled rate for the geography.
- **DO NOT** apply in land that has been or will be treated with metsulfuron and/or chlorsulfuron-containing herbicides in Nebraska and Kansas without observing the rotational crop intervals for those products.
- Injury to or loss of desirable trees or vegetation may result from failure to observe the following:
- **DO NOT** apply or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Prevent drift of spray to desirable plants.
- **DO NOT** use on lawns, walks, driveways, tennis courts or similar areas.
- **DO NOT** discharge excess material on the soil at a single spot in the field or mixing/loading station.
- **DO NOT** contaminate any body of water.
- **DO NOT** tank mix [CanopyCANOPY®](#) with organophosphate insecticides. **DO NOT** apply [CanopyCANOPY®](#) within 14 days before or after an application or organophosphate insecticide, as severe crop injury may occur. Injury to soybeans may occur if [CanopyCANOPY®](#) is used in conjunction with soil-applied organophosphate pesticides.

USE PRECAUTIONS

- Prior to using [CanopyCANOPY®](#) herbicide, consideration should be given to crop rotation plans. Crops other than soybeans may be extremely sensitive to low concentrations of [CanopyCANOPY®](#) remaining in the soil the next planting season. Choice of rotation crop is restricted following application of [DuPont™ CanopyCANOPY®](#). (See "ROTATIONAL CROP GUIDELINES" for your geographical region.)
- Thoroughly clean [CanopyCANOPY®](#) from application equipment immediately after use and prior to spraying crops other than soybeans. Failure to remove even small amounts of [CanopyCANOPY®](#) from application equipment may result in injury to subsequently sprayed crops.
- Keep from contact with fertilizers, insecticides, fungicides and seeds during storage.
- Calibrate sprayers only with clean water away from the well site. Make scheduled checks of spray equipment.
- Ensure that all operation employees accurately measure pesticides.
- Because most crops are highly sensitive to [CanopyCANOPY®](#), all direct or indirect contact (such as spray drift) to crops or to land scheduled to be planted to crops other than soybeans should be avoided.
- If a soybean variety is suspected of being sensitive to metribuzin, check with the soybean seed company before treating a field of that soybean variety with [CanopyCANOPY®](#) (contains metribuzin).
- Soybean stunting may occur if excessive rainfall occurs after application but before soybeans germinate. Injury is more prevalent under poor drainage or compacted conditions or when soil is saturated for long periods of time. Soybeans rapidly outgrow stunting once favorable growing conditions return.
- Seedling disease, nematodes, cold weather, deep planting (more than 2"), excessive moisture, high salt concentration, or drought may weaken soybean seedlings and increase possibility of crop injury.
- Mix only enough product for the job at hand.
- Avoid overfilling of spray tank.
- Dilute and agitate excess solution and apply at labeled rates or uses.
- Avoid storage of pesticides near well sites.
- When triple-rinsing the pesticide container, be sure to add the rinsate to the spray mix.

WEED RESISTANCE MANAGEMENT

[CanopyCANOPY®](#), which contains the active ingredients chlorimuron-ethyl and metribuzin, is both a Group 2 and a Group 5 herbicide based on the mode of action classification system of the Weed Science Society of America.

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is a best practice. A diversified weed management program may include the use of multiple herbicides with different sites of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistance.

The continued effectiveness of this product depends on the successful implementation of a weed resistance management program.

To aid in the prevention of developing weeds resistant to this product, users should:

- Scout fields before application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.
- Start with a clean field, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small (less than 4 inches).
- Apply full rates of [CanopyCANOPY®](#) herbicide for the most difficult to control weed in the field at the specified time (correct weed size) to minimize weed escapes.
- Scout fields after application to detect weed escapes or shifts in control of weed species.
- Control weed escapes before they reproduce by seed or proliferate vegetatively.
- Report any incidence of non-performance of this product against a particular weed to your [DuPont company](#) representative, local retailer, or county extension agent.
- Contact your [DuPont company](#) representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective sites of actions for each target weed.
- If resistance is suspected, treat weed escapes with an herbicide having a site of action other than Group 2 or 5 and/or use nonchemical methods to remove escapes, as practical, with the goal of preventing further seed production.
- Suspected herbicide-resistant weeds may be identified by these indicators:
- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

Additionally, users should follow as many of the following herbicide resistance management practices as is practical:

- Use a broad spectrum soil-applied herbicide with other sites of action as a foundation in a weed control program.
- Utilize sequential applications of herbicides with alternative sites of action.
- Rotate the use of this product with non-Group 2 or 5 herbicides.
- Avoid making more than two applications of [DuPont™ CanopyCANOPY®](#) herbicide and any other Group 2 or 5 herbicides within a single growing season unless mixed with an herbicide with a different site of action with an overlapping spectrum for the difficult-to-control weeds.
- Incorporate non-chemical weed control practices, such as mechanical cultivation, crop rotation, cover crops and weedfree crop seeds, as part of an integrated weed control program.
- Use good agronomic principles that enhance crop development and crop competitiveness.
- Thoroughly clean plant residues from equipment before leaving fields suspected to contain resistant weeds.
- Manage weeds in and around fields, during and after harvest to reduce weed seed production.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

MANDATORY SPRAY DRIFT MANAGEMENT

Ground Boom Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy unless making a rangeland application, in which case applicators may apply with a nozzle height no more than 4 feet above the ground.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- DO NOT apply when wind speeds exceed 10 miles per hour at the application site.
- DO NOT apply during temperature inversions.

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 feet above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use one-half swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented, so the spray is directed toward the back of the aircraft.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Boom-less Ground Applications:

- Applicators are required to use a Medium or coarser droplet size (ASABE S572.1) for all applications.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT MANAGEMENT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- **Adjust Nozzles** - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT – Aircraft

Higher release heights increase the potential for spray drift. When applying aurally to crops, **DO NOT** release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.

Boom-less Ground Applications:

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

APPLICATION INFORMATION

Rate Summary for DuPont ~~Canopy~~ CANOPY® Herbicide

Rate of DuPont Canopy CANOPY® per acre	Pounds of Active Ingredient Metribuzin per acre	Pounds of Active Ingredient Chlorimuron Ethyl per acre
2.25 oz	0.090	0.015

3.0 oz	0.121	0.020
3.5 oz	0.141	0.023
4.0 oz	0.161	0.027
5.0 oz	0.201	0.033
6.0 oz	0.241	0.040
7.0 oz	0.281	0.047
8.0 oz	0.322	0.054
10.0 oz	0.402	0.067
12.0 oz	0.482	0.080

APPLICATION INFORMATION – ALL USES

Geographic Use Regions

The geographical use regions for ~~DuPont™~~ [CanopyCANOPY®](#) are defined below:

Central Region: The states of Delaware, Illinois, Indiana, Iowa (fields east of State Route 63 or south of I-80), Kansas, Maryland, Michigan, Missouri (except the Bootheel), Nebraska (fields south of Route 30 and east of Route 281), New Jersey, New York (fields south of I-90), Ohio, Pennsylvania, Virginia, West Virginia and Wisconsin (fields south of I-90 between Lacrosse and Madison and fields south of I-94 between Madison and Milwaukee).

- On soils with a composite pH greater than 7.0, **DO NOT** exceed 2.25 oz/acre [CanopyCANOPY®](#).
- In the states of Michigan, New York, and Wisconsin, **DO NOT** use the 2.25 oz/acre rate on soils where the composite pH exceeds 7.6.
- In the states of New York and Wisconsin, **DO NOT** exceed 2.25 oz/acre per season.

[CanopyCANOPY®](#) may be used on fields which are composite pH 7.0 or less, but which may contain isolated areas where the pH exceeds 7.0. Use of [CanopyCANOPY®](#) at rates exceeding 2.25 oz/acre on soils which exceed composite pH 7.0 may result in unacceptable injury to the following crop.

Southern Region: The states of Alabama (except the “Black Belt” where soil pH must be less than 7.0), Arkansas, Florida, Georgia, Kentucky, Louisiana, Missouri (Bootheel region only), Mississippi (except the “Black Belt” where soil pH must be less than 7.0), North Carolina, Oklahoma, South Carolina, Tennessee and Texas (fields east of Route 183).

- On soils with a composite pH greater than 7.0 **DO NOT** exceed 3.5 oz/acre [CanopyCANOPY®](#).
- **DO NOT** apply to Black Belt Soils of Alabama and Mississippi with a soil pH greater than 7.0 or history of nutrient deficiency such as iron chlorosis, as injury may occur.
- Injury to soybeans may occur if [CanopyCANOPY®](#) is used on soils having a calcareous surface layer or pH greater than 7.5.

Application Methods

- Fall-applied, early pre-plant, pre-plant and preemergence, including burndown.
- Pre-plant incorporated. Incorporate uniformly, no deeper than the top 1-2" of soil prior to planting soybeans.
- [CanopyCANOPY®](#) may be followed sequentially by many postemergence herbicides, such as glyphosate, ~~DuPont™~~ [SYNCHRONY Synchrony®](#) XP, quizalofop, or fomesafen. See Rate Tables 4 and 6 for recommended sequential rates for [CanopyCANOPY®](#).

- Spring-applied [CanopyCANOPY®](#) may follow fall applications of [CanopyCANOPY®](#) EX.
- For sequential programs using chlorimuron ethyl-containing herbicides (including [CanopyCANOPY®](#), [CanopyCANOPY®](#) EX, and [SynchronySYNCHRONY®](#) XP), **DO NOT** exceed 0.82 oz ai/acre (0.051 lb ai/acre) chlorimuron ethyl in the Central Region States or 1.07 oz ai/acre (0.067 lb ai/acre) chlorimuron ethyl in the Southern Region States in any one soybean growing cycle.

Timing To Crop Stage

- After fall harvest, [CanopyCANOPY®](#) may be applied any time prior to soybean emergence, except on frozen ground.
- **DO NOT** apply [CanopyCANOPY®](#) after the soybean crop has emerged or severe injury or death of the crop will occur.
- **DO NOT** apply [CanopyCANOPY®](#) to frozen ground.

Burndown Information

Apply [CanopyCANOPY®](#) when weeds are young and actively growing. Applications made to weeds larger than the indicated sizes, or to weeds under stress, may result in unsatisfactory control.

When used for burndown, [CanopyCANOPY®](#) is rainfast after one hour.

- Use a minimum of 15 gallons per acre to ensure thorough coverage of the weeds and the best performance. For small weeds and/or heavy crop residue, increase the gallonage to ensure coverage.
- For best performance, select nozzle and pressure combinations that deliver medium spray droplets, as indicated, for example, by ASAE standard S572.

Spray Additives

Applications of [DuPont™ CanopyCANOPY®](#) used for burndown must include either a crop oil concentrate or a nonionic surfactant. Crop oil concentrate is the required adjuvant system unless tank mixing with a product that precludes use of crop oil concentrate.

Consult local [DuPont-Corteva Agriscience](#) fact sheets, technical bulletins, and service policies prior to using other adjuvant systems. If another herbicide is tank mixed with [CanopyCANOPY®](#), select adjuvants authorized for use with both products. Adjuvants must contain only EPA-exempt ingredients (40 CFR 1001).

Crop Oil Concentrate (COC) - Petroleum or Modified Seed Oil (MSO)

- Apply at 1% v/v (1 gal per 100 gal spray solution) or 2% under arid conditions.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

Nonionic Surfactant (NIS)

- Apply at 0.25% v/v (1 qt per 100 gal spray solution) or 0.5% under arid conditions.
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.

Tank Mixes

Other than the exceptions noted, and in addition to the tank mix partners and rates indicated in this label, [CanopyCANOPY®](#) may be tank mixed or followed with sequential applications of other products registered for use in soybeans. [CanopyCANOPY®](#) may be applied in tank mix combinations with full or reduced rates of other products provided:

- The tank mix product is labeled for the same timing, method of application, adjuvants, and use restrictions as [CanopyCANOPY®](#).
- The tank mix is not specifically prohibited on the label of the tank mix product.
- The tank mix combination is compatible as determined by a “jar test” described in the TANK MIX COMPATIBILITY TESTING section below.

Weed control and crop safety resulting from the use of tank mixtures not specifically noted on this label, or in separately published [DuPont-Corteva Agriscience](#) recommendations, are the responsibility of the user.

To select the proper tank mix product, identify the weeds which need to be controlled and consult the product labels to determine which product is needed. Consult the companion tank mix herbicide label for use instructions, rates, precautions, restrictions, and other use information. For [CanopyCANOPY®](#) tank mixes with glyphosate substitute 0.25% NIS for the 1% COC.

2,4-D (LVE) is the isooctyl (2-ethylhexyl) ester of 2,4-Dichlorophenoxyacetic acid. This product is sold under a variety of trade names. It has a minimum preplant interval of 7-30 days based on the rate used. Consult the label of the product used for specific information on this interval.

Tank Mix Compatibility Testing

Perform a jar test prior to tank mixing to ensure compatibility of [CanopyCANOPY®](#) and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, gels, oily film or layers, or other precipitates, it is not compatible.

FALL APPLICATIONS

Timing

- [CanopyCANOPY®](#) can be applied to no-till or conservation fields anytime after the fall harvest.

Timing to Weeds: Burndown

For best results, apply to annual broadleaf weeds that are up to 3 inches in height or diameter and to perennial broadleaf weeds that are up to 6 inches in height or diameter. Annual grasses should not exceed 1 inch in height. Where the rate is not restricted by soil pH, use higher [CanopyCANOPY®](#) rates for improved and longer residual activity.

Rate table 1 – Fall or Early Spring Use Rates by Region

In medium and fine soils of 1.5-4% organic matter	Rate* oz/acre
Central region Delaware, Illinois, Indiana, Iowa, Kansas, Maryland, Michigan, Missouri (except the bootheel), Nebraska, New Jersey, New York, Ohio, Pennsylvania, Virginia, West Virginia and Wisconsin no soil pH restriction	2.25
Composite soil pH of 7 or less	3-7
Southern Region Alabama, Arkansas, Georgia, Kentucky, Louisiana, Missouri (bootheel region only), Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas (fields east of Rte 183) No soil pH restriction	2.25 to 3.5
Composite soil pH of 7 or less	>3.5-7
*see , 'Geographic Use Regions', for state specific restrictions for Alabama, Iowa, Michigan, Mississippi, Missouri, Nebraska, New York, Texas, and Wisconsin.	

Weeds controlled – burndown

For the best burndown results, the addition of 2,4-D LVE is recommended, and is required for control of some weeds

DuPont™ [Canopy](#) **CANOPY**® herbicide, applied at 2.25 - 7 oz/acre, will burndown the following weeds.

Burndown Control of Emerged Winter Annual, Perennial, and Summer Annual Weeds

Annual Grasses	Mustard, tansy, wild
Bittercress, smallflowered	Pennycress, field
Bushy wallflower	Pepperweed, Virginia
Buttercup, smallflower	Pigweed, redroot
Butterweed, (Cressleaf groundsel)	Ragweed, common, giant
Dandelion	Shepherd's purse
Deadnettle, purple	Smartweed, Pennsylvania
Garlic, wild*	Speedwell, field and purslane
Henbit	Sunflower
Ladysthumb	Thistle, Canada (above ground portion)
Lambsquarters*	Velvetleaf
Lettuce, prickly	Whitlowgrass
Marestail (horseweed)*	Yellow rocket

* Addition of 8 oz ai/acre 2,4-D LVE is required for all [Canopy](#) **CANOPY**® rates.

For adjuvant and gallonage requirements for burndown applications, refer to the 'Burndown Information', 'Spray Additives', and 'Tank Mixes' sections of this label under the 'Application Information - All Uses' section.

Chickweed Burndown

- For best results: add tribenuron-methyl herbicide to [Canopy](#) **CANOPY**® for control of up to 6-inch common chickweed.
- Alternatively, metribuzin or glyphosate-containing products registered for soybeans may be added for chickweed burndown.

Weeds Controlled – Preemergence

Fall through early Spring applications of 2.25 oz/acre [Canopy](#) **CANOPY**® will provide limited residual control of listed weeds to contribute to a clean seedbed at normal planting times.

Fall through early Spring applications of 3-7 oz/acre [Canopy](#) **CANOPY**® will provide acceptable preemergence control, or partial control (suppression), of the following weeds through normal planting dates.

Weeds Controlled or Suppressed Preemergence

Control	Suppression
Cocklebur	Annual grasses* (foxtails, barnyardgrass, crabgrass, panicum)
Ladysthumb	Chickweed, common
Lambsquarters	Jimsonweed
Henbit	Morningglory, annual*
Marestail	Nutsedge, yellow*
Pigweed, redroot and smooth	Prickly sida (teaweed)*
Purslane speedwell	Ragweed, giant*
Ragweed, common	Velvetleaf
Smartweed, Pennsylvania	

Winter annual mustards (pennycress, bittercress, shepherd's-purse, whitlowgrass, yellow rocket)	
---	--

* With 2.25 oz/acre applications of [DuPont™ Canopy CANOPY®](#), heavy weed pressure, delayed planting, or adverse environmental conditions may require additional burndown control measures at planting.

In addition to the weeds noted in the lists above, [Canopy CANOPY®](#) has activity on a range of other weeds. Consult [DuPont-Corteva Agriscience](#) Fact Sheets, technical bulletins, and service policies for information on other weeds controlled.

SPRING APPLICATIONS

Application Methods

Apply [Canopy CANOPY®](#) or [Canopy CANOPY®](#) tank mixes using one of the following application methods.

- Early preplant or preplant in conservation tillage, no-till or stale seedbed systems.
- Pre-plant incorporated (incorporate uniformly, no deeper than the top 1-2" of soil prior to planting soybeans).
- Preemergence application.
- Sequential applications followed by planned postemergence treatments.

Weeds Controlled Preemergence

When used as directed, [Canopy CANOPY®](#) will provide residual control of the following weeds. Lower rates are recommended for planned sequential programs and higher rates are recommended for full-season programs. See the Rate Tables below.

*Cocklebur	Pigweed
Florida beggarwood	Palmer
Hemp sesbania	Redroot
Hophornbeam copperleaf	Smooth
Jimsonweed	Spiny amaranth
Ladysthumb	Poinsetta (wild)
Mustard, wild	Prickly sida (teaweed)
*Morningglory	Purslane, common
Annual	*Ragweed, giant
Ivyleaf	*Sicklepod
Entireleaf	Smartweed, Pennsylvania
Pitted	Spotted spurge
Smallflower	Sunflower
Tall	Velvetleaf

* Large-seeded weeds germinating deep in the soil, such as morningglory, sicklepod, cocklebur and giant ragweed, or other weeds which may emerge at various times during the growing season may require a cultivation or a postemergence herbicide application for season-long control.

When used as directed [Canopy CANOPY®](#) will provide partial control of the following weeds:

Annual grasses (Barnyardgrass, Broadleaf signalgrass, Crabgrass, Foxtail species, Panicum, Texas and Fall)	Burcucumber Chickweed, common Johnsongrass (seedling) Mexicanweed Nutsedge, purple, yellow
---	--

CENTRAL REGION STATES

Specific Use Directions

- Spring applications of [CanopyCANOPY®](#) may be applied at planting or up to 45 days before planting.

Giving careful consideration to soil type, soil pH, organic matter, rotational crop intervals, geographic location, and weed pressure, select a rate of ~~DuPont™~~ [CanopyCANOPY®](#) from Rate Table 2.

Rate Table 2 – Early Pre-plant Burndown, Pre-plant Incorporated and Preemergence:

Soil Texture	Broadcast Rate (Ounces per Acre) 0.5 - 4% Organic Matter
Coarse: Loamy Sand, Sandy Loam	4 - 5
Medium: Loam, Silt loam, Silt, Sandy Clay Loam	5 - 6
Fine: Silty Clay Loam, Clay Loam, Clay	5 - 7

For Season-long Grass Control

- **Central Region States**

[CanopyCANOPY®](#) may not provide season-long preemergence control of grasses. For improved grass control, [CanopyCANOPY®](#) may be:

- Followed as needed by a postemergence grass herbicide such as quizalofop herbicide or, in glyphosate-resistant tolerant soybeans, [CanopyCANOPY®](#) may be followed with an in-season glyphosate application.
- Tank mixed with other grass herbicides including s-metolachlor and pendimethalin.

Pre-plant Burndown - Central Region States

In addition to providing season-long preemergence control of certain broadleaf weeds and partial control of other broadleaf weeds and annual grasses, [CanopyCANOPY®](#) will provide burndown control of the following broadleaf weeds up to 3" in diameter or height and annual grasses up to 1".

Annual grasses	Mustard, wild, tansy
Bittercress, smallflowered	Pennycress, field
Bushy wallflower	Pepperweed, Virginia
Buttercup, smallflower	Pigweed, redroot
Butterweed (Cressleaf groundsel)	Aagweed, common, giant
Dandelion	Shepherd's-purse
Deadnettle, purple	Smartweed, Pennsylvania
Garlic, wild*	Speedwell, purslane
Henbit	Sunflower
Ladysthumb	Thistle, Canada (above ground portion)
Lambsquarters*	Velvetleaf
Lettuce, prickly	Whitlowgrass
Marestail (horseweed)*	Yellow rocket

* Addition of 8 oz ai/ac 2,4-D LVE is required for all [CanopyCANOPY®](#) rates

For Spring Burndown control, pick the appropriate rate from Rate Table 2, 3 or 4

For burndown of larger annual grasses or broadleaf weeds exceeding 1-3", or for burndown of weeds not

listed above, [CanopyCANOPY[®]](#) may be tank mixed with one or more of the following:

quizalofop
glyphosate
paraquat
2,4-D (LVE)

For adjuvant and gallonage requirements for burndown applications, refer to the 'Burndown Information', 'Spray Additives', and 'Tank Mixes' sections of this label under the 'Application Information - All Uses' section.

[CanopyCANOPY[®]](#) Tank mixes with Metribuzin - Central Region States

[CanopyCANOPY[®]](#) may be applied at reduced rates when tank mixed with metribuzin or metribuzin-containing products. These tank mixes will generally provide season-long preemergence weed control for the weeds listed below. When used according to the directions in the previous section for Burndown control, these tank mixes will also provide Burndown control of the weeds claimed in the previous section.

Reduced rates of [CanopyCANOPY[®]](#) tank mixed with metribuzin or metribuzin-containing products will generally provide season-long preemergence control of the following weeds:

Ladysthumb	Ragweed, common
Lambsquarters	Smartweed, Pennsylvania
Mustard, wild	Velvetleaf
Pigweeds: Palmer Redroot Smooth Spiny amaranth	

[DuPont™ CanopyCANOPY[®]](#) + metribuzin tank mixes will provide partial control (suppression) preemergence of the following weeds:

Cocklebur Craggrass Eastern black nightshade Foxtail species Jimsonweed	Morningglories Entireleaf Ivyleaf Pitted Tall Waterhemp
---	--

Choose a reduced rate of [CanopyCANOPY[®]](#) and a rate of metribuzin from Rate Table 3 below.

Rate Table 3 - Reduced Rate [CanopyCANOPY[®]](#) Tank mixes with metribuzin

Soil Texture	Broadcast Rate (per acre) 0.5 - 4% Organic Matter CanopyCANOPY[®] + metribuzin		
	Coarse: Loamy Sand, Sandy Loam	2.25* - 4 oz	1.5 – 3 oz ai
Medium or Fine: Loam, Silt Loam, Silt, Sandy Clay Loam, Silty Clay Loam, Clay Loam, Clay	2.25* - 4 oz	3 - 4.5 oz ai	.

*2.25 oz/acre is the maximum rate on soil with a composite pH greater than 7.0

Sequential Applications - Central Region States

Reduced rates of [CanopyCANOPY®](#), from 2.25 - 7 oz/acre, may be followed, as needed, by sequential applications of many postemergence herbicides such as [SynchronySYNCHRONY®](#) XP or chlorimuron-ethyl. Reduced rates of [CanopyCANOPY®](#) in Rate Table 4 below, will provide early-season residual control (of the weeds listed under “Weeds Controlled-Preemergence: Spring Applications”) prior to the planned postemergence program.

Rate Table 4 - Sequential Applications: [CanopyCANOPY®](#) followed by Postemergence

CanopyCANOPY® Broadcast (oz/a)	Sequential rate limits for chlorimuron-ethyl (oz ai/acre) or SynchronySYNCHRONY® XP (oz product/acre)
2.25*	On soils with composite pH greater than 7.0, DO NOT follow with any chlorimuron-ethyl containing herbicide (including SynchronySYNCHRONY® XP)
3 - 5	Chlorimuron-ethyl up to 0.1875 oz ai (0.0117 lb) or SynchronySYNCHRONY® XP up to 0.75 oz
6	Chlorimuron-ethyl up to 0.165 oz ai (0.0103 lb) or SynchronySYNCHRONY® XP up to 0.75 oz.
7	Chlorimuron-ethyl up to 0.0625 oz ai (0.0039 lb)

* 2.25 oz/acre is the maximum rate on soil with composite pH greater than 7.0.

ROTATIONAL GUIDELINES FOR FALL AND SPRING **DUPONT™** CANOPY® APPLICATIONS –

Central Region States

Central Region: The states of Delaware, Illinois, Indiana, Iowa (east of State Route 63 or south of I-80), Kansas, Maryland, Michigan, Missouri (except the Bootheel), Nebraska (fields south of Route 30 and east of Route 281), New Jersey, New York (fields south of Interstate 90), Ohio, Pennsylvania, Virginia, West Virginia and Wisconsin (fields south of Interstate 90 between Lacrosse and Madison and fields south of Interstate 94 between Madison and Milwaukee).

When used as described in the Central Region section of this label, Rotational Guideline 1 describes the minimum length in months from the time of [CanopyCANOPY®](#) application until [CanopyCANOPY®](#) treated soil can be replanted to the crops listed in the table. For Fall applications, begin counting the re-cropping interval from the normal Spring planting time for soybeans in your area.

Crop rotation intervals noted below are based on crops grown under favorable growing conditions. Crops grown under unfavorable environmental conditions, such as drought, nutrient deficiency, high salts, disease and insect pressure may demonstrate reduced tolerance to crop protection chemicals. When deciding on a particular crop to replant in your fields, carefully consider your particular soil and other field conditions. When a recommended tank mix is used, consult the tank mix partner labels for recropping instructions and follow the directions that are most restrictive.

Rotational Guideline 1 - Central Region

For all recommended Fall and Spring [CanopyCANOPY®](#) uses, including sequentials with

CanopyCANOPY® EX or DuPont™ SynchronySYNCHRONY® XP

Crop	Recropping Interval in Months
Soybeans	Anytime
Barley, Ryegrass, Wheat, Winter Rye	4
Alfalfa	10
Cotton	10
Rice	10
Tobacco (transplant)	10
Tomato (transplant)	10
Field Corn*	10
Clover	12
Dry Beans, Kidney Beans, Snap Beans, Peas	12
Sorghum	12
Cucumber, Flax, Peanuts, Pumpkin, Sunflower, Sweet Corn, Watermelon, Cabbage, Canola (rapeseed), Lentils, Mustard	18
Carrot, Onion, Potato, and Sugar Beets and any other crop not listed	30

* Field Corn is defined to include only that corn grown for grain or silage, popcorn, and seed corn. However, because seed corn inbred lines may vary in their sensitivity to trace amounts of herbicide carryover, [DuPont-Corteva Agriscience](#) cannot warrant that seed corn can be recropped without damage or yield loss. Users should seek the advice of their seed corn company agronomists regarding inbred sensitivity to herbicides prior to planting any inbred lines.

SOUTHERN REGION STATES

Specific use directions

Spring applications of [CanopyCANOPY®](#) may be applied at planting or up to 45 days prior to planting.

Giving careful consideration to soil type, soil pH, organic matter, rotational crop intervals, geographic location, and weed pressure, select a rate of [CanopyCANOPY®](#) from **Rate Table 5** below. Apply [CanopyCANOPY®](#) Early Preplant, Preplant Incorporated or Preemergence as directed in the 'Application Information - All Uses' section of this label.

Rate Table 5 – Early Pre-plant, Pre-plant, Pre-plant Incorporated and Preemergence

	Broadcast Rate	
	Percent Organic Matter in Soil*	
	0.5 - 3%	3 - 5%
Coarse: Loamy Sand, Sandy Loam	6	8
Medium: Loam, Silt Loam*, Silt, Sandy Clay Loam	8	10
Fine: Silty Clay Loam, Clay Loam, Clay	10	12

*On silt loam soils in TN and KY use 6-8oz

For Season-long Grass Control

- Southern Region States

DuPont™ CanopyCANOPY® may not provide season-long preemergence control of grasses. For improved grass control, CanopyCANOPY® may be:

- Followed as needed by a postemergence grass herbicide such as quizalofop herbicide or, in glyphosate-resistant soybeans, CanopyCANOPY® may be followed with an in-season glyphosate application.
- Tank mixed with such herbicides as s-metolachlor and pendimethalin.

Use in Stale Seedbed or Conservation Tillage - Southern Region States

For Burndown control of small annual grasses and broadleaf weeds, use 3-4 oz of CanopyCANOPY® and apply up to 45 days prior to planting. Select the higher rate for larger weeds. For burndown weeds controlled, see the “Pre-plant Burndown - Central Region States” section in this label.

When burndown plus residual control is desired, CanopyCANOPY® may be applied at-planting or up to 45 days prior to planting at a rate of 4 to 12 oz. Select a rate, based on soil type from either Rate Table 5 or Rate Table 6.

For burndown of weeds and grasses not listed above, or for burndown of larger weeds and grasses, it is recommended that CanopyCANOPY® be tank mixed with such herbicides as 2,4-D LVE, paraquat, and/or glyphosate.

For adjuvant and gallonage requirements for burndown applications, refer to the ‘Burndown Information’, ‘Spray Additives’, and ‘Tank Mixes’ sections of this label under the ‘Application Information - All Uses’ section.

Sequential Applications - Southern Region States

CanopyCANOPY® may be applied at reduced rates when followed by one planned postemergence treatment of DuPont™ SynchronySYNCHRONY® XP, chlorimuron-ethyl, or by other herbicides registered for soybeans. Select a rate of CanopyCANOPY®, according to soil type, from Rate Table 6.

Rate Table 6 - Sequential Applications - CanopyCANOPY® followed by Postemergence

Soil Texture	Broadcast Rate (Ounces per Acre) 0.5 - 4% Organic Matter
Any*	3 - 3.5
Coarse: Loamy Sand, Sandy Loam	4 - 6
Medium: Loam, Silt loam, Silt, Sandy Clay Loam	4 - 6
Fine: Silty Clay Loam, Clay Loam, Clay	6 - 8

* 3.5 oz/acre is the maximum rate that may be used on soils with a composite pH greater than 7.0. When re-cropping to rice and using 3-3.5 oz/acre on soils with pH greater than 7.0, the recrop interval is 18 months.

Rotational Guidelines for Fall and Spring DuPont™ CanopyCANOPY® Applications - Southern Region States

Southern Region: The states of Alabama (except the “Black Belt” where soil pH must be less than 7.0), Arkansas, Florida, Georgia, Kentucky, Louisiana, Missouri (Bootheel region only), Mississippi (except the “Black Belt” where soil pH must be less than 7.0), North Carolina, Oklahoma, South Carolina, Tennessee

and Texas (fields east of Route 183).

When used as described in the Southern section of this label, the table describes the minimum length in months from the time of [CanopyCANOPY®](#) application before [CanopyCANOPY®](#) treated soil can be replanted to the crops listed in the table. For Fall applications, begin counting the re-cropping interval from the normal Spring planting time for soybeans in your area.

Crop rotation intervals noted below are based on crops grown under favorable growing conditions. Crops grown under unfavorable environmental conditions, such as drought, nutrient deficiency, high salts, disease and insect pressure may demonstrate reduced tolerance to crop protection chemicals. When deciding on a particular crop to replant in your fields, carefully consider your particular soil and other field conditions. **When a recommended tank mix is used, consult the tank mix partner labels for recropping instructions and follow the directions that are most restrictive.**

Rotational Guideline 2 - Southern Region

For all recommended Fall and Spring [CanopyCANOPY®](#) uses, including sequentials with [CanopyCANOPY® EX](#), [DuPont™ SynchronySYNCHRONY® XP](#), or Chlorimuron-ethyl

Group I - composite soil pH greater than 7.0, maximum 3.5 oz/acre [CanopyCANOPY®](#), no postemergence or [SynchronySYNCHRONY® XP](#) or chlorimuron-ethyl

- All southern states

Group I - soil pH 7.0 or less

- States of AL, AR, FL, GA, LA, MS or TX
- States of KY, MO Bootheel, NC, OK, SC, TN – Use rate less than 10 oz./A
(if rate is 10 oz/A or greater, use Group II Guideline)

Group II - soil pH greater than 7.0 and [CanopyCANOPY®](#) rate greater than 3.5 oz/acre

All southern states		
Crops	Group I	Group II
Soybeans	Anytime	Anytime
Barley, Ryegrass, Wheat, Winter Rye	4	4
Alfalfa	10	18
Clover	12	18
Field Corn *	9/10 [†]	18
Cotton	10	18
Peanuts	8	18
Rice [§]	10	18
Sorghum	10	18
Tobacco (Transplant)	10	18
Tomato (Transplant)	10	18
Cucumber, Flax, Pumpkin, Sunflower, Sweet Corn, Watermelon, Cabbage, Canola (rapeseed), Lentils, Mustard, Carrot, Onion, Potato, Sugar Beets and any crop not listed above	18	30

* Field Corn is defined to include only that corn grown for grain or silage, popcorn, and seed corn. However, because seed corn inbred lines may vary in their sensitivity to trace amounts of herbicide carryover, [DuPont-Corteva Agriscience](#) cannot warrant that seed corn can be recropped without damage or yield loss. Users should seek the advice of their seed corn company agronomists regarding inbred sensitivity to herbicides prior to planting any inbred lines.

[†] May be recropped to field corn after 9 months if the [CanopyCANOPY®](#) rate does not exceed 6 oz/acre.

[§] The recrop to rice is 18 months after 3-3.5 oz/acre is used on soils with a composite pH greater than 7.0.

APPLICATION EQUIPMENT

SPRAY TANK PREPARATION

It is important that spray equipment is clean and free of existing pesticide deposits before using [DuPont™ CanopyCANOPY®](#). Follow the spray tank cleanout procedures specified on the label of product previously sprayed. If no cleanout procedure is provided, follow the cleanout procedure below for all application equipment.

1. Thoroughly rinse sprayer, tanks, boom, and hoses with clean water.
2. Partially fill the tank with water and add one of the cleaning agents listed in the SPRAYER CLEANUP section of this label. Complete filling the tank and flush the cleaning solution through the boom and hoses. Let stand for 15 minutes with agitation or recirculation and then drain the tank after flushing the hoses, boom, and nozzles.
3. Thoroughly rinse sprayer, tanks, boom, and hoses with clean water.
4. Follow label directions of the product previously sprayed for rinsate disposal.

Notes: During an extended period where spraying or mixing equipment will be used to apply multiple loads of [CanopyCANOPY®](#), at the end of each day of spraying partially fill the tank with fresh water, flush the boom and hoses and allow to sit overnight.

A steam cleaning of aerial spray tanks is recommended to dislodge any visible pesticide deposits.

EQUIPMENT/ SPRAY VOLUMES

Ground Application, conventional tillage:

- Use a minimum of 10 gallons per acre to ensure uniform coverage of soil and the best performance.
- For best performance, select nozzle and pressure combinations that deliver coarse to very coarse spray droplets, as indicated, for example, by ASAE standard S572.

Ground Application, conservation tillage- burndown:

- Use a minimum of 15 gallons per acre to ensure thorough coverage of the weeds and the best performance. For small weeds and/or heavy crop residue, increase the gallonage to ensure coverage.
- For best performance, select nozzle and pressure combinations that deliver medium spray droplets, as indicated, for example, by ASAE standard S572.

Aerial Application: [CanopyCANOPY®](#) may be applied by air for early preplant, preplant incorporated or preemergence use on soybeans. Apply uniformly with properly calibrated aerial equipment. Use a minimum of 2 gallons of water per acre. Avoid overlapping. Continuous agitation of the spray tank is required to keep the material in suspension.

MIXING INSTRUCTIONS

Fill tank 1/4 full with water. Start agitation system, add [CanopyCANOPY®](#) and continue adding water. Add separately each additional component of any tank mix while adding water. Continue agitation throughout. If poor mixing should occur with any component, premix the component with two parts water before adding to the spray tank.

A fertilizer solution may be used in the spray mixture. Small quantities should be tested for compatibility by the following procedures before full-scale mixing.

1. Put 1 pint of fertilizer solution in a quart jar.
2. Mix 2 teaspoons [CanopyCANOPY®](#) with 2 tablespoons of water; mix thoroughly and add to fertilizer solution.
3. Close jar and shake well.
4. If other herbicides are to be used in the mixture, premix 2 teaspoons of wettable powder or 1 teaspoon of liquid with 2 tablespoons of water; add to [CanopyCANOPY®](#)/fertilizer solution mixture.
5. Close jar and shake well.
6. Watch mixture for several seconds; check again in 30 minutes.
7. If mixture does not separate, foam, gel, or become lumpy, it may be used.
8. Mixing ability may be improved by adding compatibility agents.

Provided the above procedure shows the mixture to be compatible, prepare the tank mixture as follows: Add the fertilizer solution to the spray tank first, with the agitator running, add the required amount of [CanopyCANOPY®](#) and thoroughly mix. For tank mixtures with other herbicides, follow directions above. For tank mixtures with other herbicides, all applicable directions, restrictions and precautions for the additional herbicides are also to be followed.

Use [DuPont™ CanopyCANOPY®](#) spray preparations the same day as mixed or product degradation may occur. Thoroughly reagitrate and remix before using, if allowed to settle. When tank mixing with other herbicides, all applicable directions, restrictions and precautions for the additional herbicides are also to be followed.

SPRAYER CLEANUP

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of [CanopyCANOPY®](#) as follows: *

1. Drain tank; thoroughly hose down the interior surfaces of the tank; then flush tank, boom, and hoses with clean water for a minimum of 5 minutes.
2. Partially fill the tank with water and add one gallon of household ammonia* (containing 3% active) for every 100 gallons of water. Complete filling the tank with water, then flush the cleaning solution through the boom, hoses, and nozzles. Add water to completely fill the tank and allow to agitate or recirculate for at least 15 minutes. Again, flush the boom, hoses and nozzles, and drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing water and the cleaning agent.
4. Repeat Step 2.
5. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing water through the boom and hoses.

*Equivalent amounts of an alternate strength ammonia solution or a tank cleaner recommended in separately published DuPont bulletins may be used.

THE IMPORTANCE OF SOIL PH

Soil pH varies greatly, even within the same field. pH variations as much as 2 pH units are common. Composite soil samples taken across an entire field, such as those samples taken for soil fertility recommendations, may not detect areas of high pH. Sub-sampling is recommended for areas likely to have pH values higher than the field average. The following is a non-inclusive list of potential high pH areas where subsampling is recommended.

- Where different soil types are evident within a field, sample soil types separately.
- Where conditions vary within a field, sample areas separately, such as:
 - areas bordered by limestone gravel roads,
 - river bottoms subject to flooding,
 - low areas in hardpan soils where evaporative ponds may occur,
 - eroded hillsides,
 - along drain tile lines, and

- areas where drainage ditch spoil has been spread.
- Where lime has not been deeply incorporated, soil may exhibit significantly higher pH values in the upper 3 inches of soil. Composite soil samples taken at a 6-8 inch depth may not reflect the elevated pH near the surface. In these cases, shallow sampling, the upper 3 inches, is advised.

Determine soil pH by laboratory analysis using a 1:1 soil:water suspension.

IDENTIFICATION INFORMATION FOR OTHER PRODUCTS INCLUDED IN THIS LABEL

USEPA REGISTERED PRODUCTS MENTIONED IN THIS LABEL FOR USE IN TANK MIXTURES OR OTHER REASONS		
PRODUCT BRAND NAME	ACTIVE INGREDIENT(S)	EPA REGISTRATION NUMBER
Canopy EX	Chlorimuron Ethyl + Tribenuron methyl	352-635
Synchrony XP	Chlorimuron Ethyl + Thifensulfuron methyl	352-648

STORAGE AND DISPOSAL

Pesticide Storage: ~~Store product in original container only. DO NOT contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a cool, dry place.~~

Pesticide Disposal: ~~DO NOT contaminate water, food, or feed by disposal. Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.~~

Container Handling: ~~Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.~~

Nonrefillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds): ~~Nonrefillable container. DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. DO NOT burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities~~

Nonrefillable Plastic and Metal Containers (Capacity Greater Than 50 Pounds): ~~Nonrefillable container. DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. DO NOT burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.~~

Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down): ~~Nonrefillable container. DO NOT reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that~~

the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Paper or Plastic Bags, Fiber Sacks including Flexible Intermediate Bulk Containers (FIBC) or Fiber Drums With Liners: Nonrefillable container. **DO NOT** reuse or refill this container. Completely empty paper or plastic bag, fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances.

Refillable Fiber Drums With Liners: Refillable container (fiber drum only). Refilling Fiber Drum: Refill this fiber drum with DuPont™ CANOPY® containing metribuzin and chlorimuron ethyl only. **DO NOT** reuse this fiber drum for any other purpose. Cleaning before refilling is the responsibility of the refiller. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Disposing of Fiber Drum and/or Liner: **DO NOT** reuse this fiber drum for any other purpose other than refilling (see preceding). Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container. Offer the liner for recycling if available or dispose of liner in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner. To clean the fiber drum before final disposal, completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances.

All Other Refillable Containers: Refillable container. Refilling Container: Refill this container with DuPont™ CANOPY® containing metribuzin and chlorimuron ethyl only. **DO NOT** reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, **DO NOT** use the container, contact DuPont at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, **DO NOT** reuse or transport container, contact DuPont at the number below for instructions. Disposing of Container: **DO NOT** reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Outer Foil Pouches of Water Soluble Packets (WSP): Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available or, dispose of the empty outer foil pouch in the trash

~~as long as WSP is unbroken. If the outer pouch contacts the formulated product in any way, the pouch must be triple rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer pouch as described previously.~~

~~**DO NOT** transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact 1-800-992-5994, day or night.~~

~~**NOTICE TO BUYER:** Purchase of this material does not confer any rights under patents of countries outside of the United States.~~

~~®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners~~

~~“Flexstar” is a trademark of Syngenta Crop Protection Inc.~~

~~“Lorsban” is a trademark of Dow AgroSciences LLC~~

~~“Mocap”, “Nemacur” and “Di-Syston” are registered trademarks of Bayer CropScience~~

~~“Thimet” is a registered trademark of Micro Flo Company LLC~~

~~**LIMITATION OF WARRANTY AND LIABILITY**~~

~~NOTICE: Read this Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.~~

~~It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product, crop injury, or injury to non-target crops or plants. WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.~~

~~DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.~~

~~TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.~~

~~To the extent consistent with applicable law that allows such requirement, DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise, or be barred from any remedy.~~

~~This Limitation of Warranty and Liability may not be amended by any oral or written agreement.~~

Terms and Conditions of Use

If terms of the following [Warranty Disclaimer](#), [Inherent Risks of Use](#), and [Limitation of Remedies](#) are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent consistent with applicable law ~~To the extent permitted by law~~, use by the buyer or any other user constitutes acceptance of the terms under [Warranty Disclaimer](#), [Inherent Risks of Use](#) and [Limitations of Remedies](#).

Warranty Disclaimer

Corteva Agriscience warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions for use, subject to the inherent risks set forth below. To the extent consistent with applicable law ~~To the extent permitted by law~~, Corteva Agriscience **MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application or other factors, all of which are beyond the control of Corteva Agriscience or the seller. To the extent consistent with applicable law, Corteva Agriscience will not be responsible for losses or damages resulting from the use of this product in any manner not specifically directed by Corteva Agriscience. To the extent consistent with applicable law ~~To the extent permitted by law~~, all such risks associated with non-directed use shall be assumed by buyer and/or user.

Limitation of Remedies

To the extent consistent with applicable law ~~To the extent permitted by law~~, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, tort, strict liability, or other legal theories), shall be limited to, at Corteva Agriscience's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of product used.

~~To the extent permitted by law~~To the extent consistent with applicable law ~~w~~, Corteva Agriscience shall not be liable for losses or damages resulting from handling or use of this product unless Corteva Agriscience is promptly notified of such loss or damage in writing. To the extent consistent with applicable law ~~To the extent permitted by law~~, in no case shall Corteva Agriscience be liable for consequential, incidental or special damages or losses.

The terms of the [Warranty Disclaimer](#), [Inherent Risks of Use](#), and [Limitation of Remedies](#) cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Corteva Agriscience or the seller is authorized to vary or exceed the terms of the [Warranty Disclaimer](#) or this [Limitation of Remedies](#) in any manner.

For product information call: 1-800-258-3033

Produced for

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268~~E. I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Road, Wilmington, Delaware 19805 U.S.A.~~

All rights reserved.

TM® Trademarks of Corteva Agriscience and its affiliated companies

~~®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners~~

EPA accepted 06/30/2021