

352-446

10-15-2003

1/15



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OCT 15 2003

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Mr. Jacob J. Vukich
E.I. DuPont de Nemours & Company
DuPont Crop Protection
Stine Research Center
P. O. Box 30
Newark, DE 19714-0030

Dear Mr. Vukich:

Subject: DuPont Harmony GT Herbicide (Update First Aid Statements)
EPA Registration No. 352-446
Application Dated August 1, 2003

The labeling referred to above submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended is acceptable provided you make the following changes before you release the product for shipment.

1. At the beginning of the list of Personal Protective Equipment (PPE) within the Precautionary Statements, add the statements "Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart. In addition, revise the requirement for "chemical-resistant gloves made of any waterproof material to a requirement for "chemical-resistant gloves made of any waterproof material **such as polyethylene or polyvinyl chloride.**
2. Revise the last sentence of your Environmental Hazard section to read "Do not contaminate water by **cleaning of equipment or** disposing of equipment washwaters or wastes.
3. Under Storage and Disposal revise "Storage" to read "Pesticide Storage".

Submit three (3) copies of final printed labeling incorporating the above changes before you release the product for shipment. Amended labeling supercedes all previously accepted ones. A stamped copy of labeling is enclosed for your records.

Sincerely,

James A. Tompkins for
James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505C)

2/15

H-



Harmony[®] GT

herbicide

DRAFT LABEL



“..... A Growing Partnership With Nature”

DUPONT HARMONY® GT HIGHLIGHTS

- May be applied by ground or by air.
- Wheat, Barley, Oat, Soybeans and Field Corn may be replanted anytime after the application of HARMONY® GT. Any other crop may be planted 45 days after the application of HARMONY® GT.
- Certain environmental conditions, such as cool and dry, or hot and humid weather, affect the performance of HARMONY® GT. (See Environmental Conditions.)
- Consult label text for complete instructions. Always read and follow label directions for use.

Cereals

- For selective postemergence broadleaf weed control in Wheat (including Durum wheat), Barley, Oat and Fallow.
- Apply at the rate of 0.3 to 0.6 ounce per acre on Wheat, Barley and Fallow; 0.3 to 0.4 ounce per acre on Oat (see Cereals Application Information).
- Apply after the crop is in the 2-leaf stage, but before the flag leaf is visible on Wheat, Barley and Winter Oat. On Spring Oat, apply after the crop is in the 3 leaf stage, but before jointing.
- Use in tank mixtures with other registered herbicides for broader spectrum weed control (see Cereals Tank Mixtures).

Soybeans

- For selective postemergence broadleaf weed control in soybeans.
- Apply at the rate of 0.083 (1/12) ounce per acre.
- Include a spray additive recommended in this label. (See Soybeans Spray Additives)
- Include a nitrogen fertilizer (example: 4-8 pints of 28-0-0). (See Soybeans Spray Additives.)
- For ground application to optimize HARMONY® GT performance, use flat fan nozzles and apply in 10-25 gallons of water at 25-60 psi.
- Apply to actively growing weeds at the recommended sizes. (See Soybeans Weeds Controlled.)
- Tank mix only with pesticides specified by this or other supplemental labeling. (See Soybeans Tank Mix Applications.)

TABLE OF CONTENTS

PRECAUTIONARY STATEMENTS	1
GENERAL INFORMATION	2
DIRECTIONS FOR USE	2
CEREALS	
Application Timing	2
Use Rate	2
Spray Additives	3
Weeds Controlled	3
Specific Weed Problems	3
Tank Mixtures	4
SOYBEANS	
Application Timing	6
Use Rate	6
Spray Additives	6
Cultivation	6
Weeds Controlled	6
Tank Mixtures	6
GENERAL USE AND APPLICATION	
DIRECTIONS - ALL CROPS	
Ground Application	8
Aerial Application	8
Environmental Conditions.....	8
Crop Rotation	8
Grazing	8
Mixing Instructions.....	8
Spray Equipment	8
SPRAYER CLEANUP	9
At the End of the Day	9
After Spraying HARMONY® GT and before Spraying Crops Other Than Wheat, Barley, Oat and Soybeans	9
SPRAY DRIFT MANAGEMENT	9
Importance of Droplet Size	9
Controlling Droplet Size - General Techniques	9
Controlling Droplet Size - Aircraft	10
Boom Height	10
Wind	10
Temperature and Humidity	10
Temperature Inversions	10
Shielded Sprayers	10
Air-Assisted (Air Blast) Field Crop Sprayers	10
WEED RESISTANCE	10
INTEGRATED PEST MANAGEMENT	11
PRECAUTIONS	11
STORAGE AND DISPOSAL	11
NOTICE OF WARRANTY	12



Harmony[®] GT

herbicide

Dry flowable

For Use on Wheat, Barley, Oat, Fallow and Soybeans

Active Ingredients	By Weight
Thifensulfuron-methyl	
Methyl 3-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl) amino]carbonyl]amino] sulfonyl]-2-thiophenecarboxylate	75%
Inert Ingredients	25%
TOTAL	100%

EPA Reg. No. 352-446

ACCEPTED
with COMMENTS
In EPA Letter Dated:
OCT 15 2003

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 352-446

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

For medical emergencies involving this product, call toll-free 1-800-441-3637.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution! Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical resistant gloves made of any waterproof material.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR part 170 Section 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply where/when conditions favor runoff.

4/15

5/15

PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Assure accurate measurement of pesticides by all operation employees.
- Mix only enough product for the job at hand.
- Avoid over-filling of spray tank.
- Do not discharge excess material on the soil at a single spot in the field/grove or mixing/loading station.
- Dilute and agitate excess solution and apply at labeled rates/uses.
- Avoid storage of pesticides near well sites.
- When triple rinsing the pesticide container, be sure to add the rinsate to the spray mix.

GENERAL INFORMATION

DuPont HARMONY® GT herbicide is recommended for selective postemergence control of certain broadleaf weeds in wheat (including durum), barley, oat, fallow and soybeans. HARMONY® GT is a dry flowable granule to be mixed in water or other recommended carrier and applied as a uniform broadcast spray. It is noncorrosive, nonflammable, nonvolatile and does not freeze.

Best results are obtained when HARMONY® GT is applied to young, actively growing weeds. The use rate will depend on weed spectrum and size of weed at time of application. The degree of control and duration of effect are dependent on rate used, sensitivity and size of target weed and environmental conditions at the time of and following application.

HARMONY® GT stops growth of susceptible weeds rapidly. However, typical symptoms of dying weeds (discoloration) may not be noticeable for 1-3 weeks after application (2-5 weeks for wild garlic) depending on the environmental conditions and weed susceptibility. Warm, moist conditions following treatment promote the activity of HARMONY® GT, while cold, dry conditions delay the activity. Weeds hardened-off by cold weather or drought stress will be less susceptible.

A vigorous growing crop will aid weed control by shading and providing competition for weeds. However, a dense crop canopy at time of application can intercept spray and result in reduced weed control. Weeds may not be adequately controlled in areas of thin crop stand or seeding skips.

Applications made to weeds that are in the cotyledon stage, larger than the size indicated, or to weeds under stress may result in unsatisfactory control.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Chemical resistant gloves made of any waterproof material.
- Shoes plus socks.

Do not apply this product through any type of irrigation system. HARMONY® GT herbicide must be used only in accordance with recommendations on this label or in separately published DuPont recommendations.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by DuPont.

HARMONY® GT is recommended for use on wheat, barley, oat, fallow and soybeans in most states. Check with your state extension or Dept. of Agriculture before use, to be certain HARMONY® GT is registered in your state.

CEREALS

APPLICATION TIMING

Wheat (Including Durum), Barley and Winter Oat

Make applications after the crop is in the 2-leaf stage, but before the flag leaf is visible.

Spring Oat

Make applications after the crop is in the 3-leaf stage, but before jointing. Do not use on "Ogle", "Porter" or "Premier" varieties as crop injury can occur.

Fallow

Apply HARMONY® GT in the spring or fall when the majority of weeds have emerged and are actively growing.

Pre-Plant Burndown

For burndown of emerged weeds, broadcast applications of HARMONY® GT may be applied up through planting, but before wheat (including durum), barley and oat plants emerge.

USE RATES

In cereals, do not use less than 0.3 ounce HARMONY® GT per acre.

6/15

If predominant weed(s) in field is (are) one of those listed in WEEDS PARTIALLY CONTROLLED table below, always include a tank mix partner (refer to TANK MIXTURES).

Wheat and Barley

Apply 0.5 ounce HARMONY® GT per acre to wheat (including durum) or barley for control or partial control of the weeds listed below.

Use 0.6 ounce HARMONY® GT per acre when weed infestation is heavy and predominately consists of those weeds listed under partial control, or when application timing and environmental conditions are marginal (refer to the sections APPLICATION TIMING AND GENERAL INFORMATION).

Use 0.3 ounce HARMONY® GT per acre when weed infestation is light and predominately consists of those weeds listed under weeds controlled, and when optimum application conditions occur.

Sequential treatments of HARMONY® GT may be made provided the total amount of HARMONY® GT applied to the crop does not exceed 1.0 ounce per acre.

Oat (Spring and Winter)

Apply 0.3 to 0.4 ounce HARMONY® GT per acre for control of the weeds listed in WEEDS CONTROLLED table.

If predominant weed(s) in field is(are) one of those listed in WEEDS PARTIALLY CONTROLLED table below, always include a tank mix partner (refer to TANK MIXTURES).

Do not make more than one application of HARMONY® GT per crop season on oat.

Fallow

HARMONY® GT may be used as a postemergence fallow treatment, in combination with other suitable registered fallow herbicides such as "Landmaster" BW, "Fallow Master", "Roundup" plus 2,4-D (ester formulations work best), "Roundup" plus "Banvel"/"Banvel" SGF/"Clarity", or with 2,4-D, or "Banvel"/"Banvel" SGF/"Clarity". Apply HARMONY® GT at 0.3 to 0.6 ounce per acre to fallow for control or partial control of the weeds listed below.

Read and follow all manufacturer's label recommendations for the companion herbicide. If those recommendations conflict with this label, do not tank mix the herbicide with HARMONY® GT.

Pre-Plant Burndown

HARMONY® GT may be used as a pre-plant burndown treatment, in combination with other suitable registered pre-plant herbicides (such as "Roundup").

Apply HARMONY® GT at 0.3 to 0.6 ounce per acre for control or partial control of the weeds listed below.

Read and follow all manufacturer's label recommendations for the companion herbicide. If those recommendations conflict with this label, do not tank mix the herbicide with HARMONY® GT.

SPRAY ADDITIVES

Consult your agricultural dealer, applicator, or DuPont representative for a listing of recommended surfactants.

Antifoaming agents may be used if needed.

Do not use low rates of liquid nitrogen fertilizer solution as a substitute for surfactant.

Cereals and Fallow

Unless otherwise specified, add a DuPont recommended nonionic surfactant having at least 80% active ingredient at 1 to 2 qt per 100 gal of spray solution (0.25 to 0.5% v/v-refer to TANK MIXTURES for specific adjuvant recommendations when HARMONY® GT is used in a tank mix).

WEEDS CONTROLLED

- | | |
|----------------------|-------------------------|
| Annual knawel | Miners lettuce |
| Annual sowthistle | Mouseear chickweed |
| Black mustard | Pennsylvania smartweed |
| Bushy wallflower | Prostrate knotweed |
| /Treacle mustard | Redmaids |
| Carolina geranium | Redroot pigweed |
| Coast fiddleneck | Russian thistle* |
| Common buckwheat | Scentless |
| Common chickweed* | chamomile/mayweed |
| Common groundsel | Shepherdspurse |
| Common lambsquarters | Smallflower buttercup |
| Corn chamomile | Stinking mayweed |
| Corn spurry | /Dogfennel |
| Cress (mouse-ear) | Swinecress |
| Curly dock | Tarweed fiddleneck |
| False chamomile | Tumble/Jim Hill mustard |
| Field pennycress | Volunteer lentils |
| Flixweed | Volunteer peas |
| Green smartweed | Volunteer sunflower |
| Kochia * | Wild buckwheat* |
| Ladysthumb | Wild chamomile |
| London rocket | Wild garlic* |
| Mallow (little) | Wild mustard |
| Marshelder | |

PARTIAL CONTROL**

- | | |
|-------------------------|------------------|
| Common cocklebur | Mallow (common) |
| Common sunflower | Prickly lettuce* |
| Cutleaf eveningprimrose | Tansymustard* |
| Henbit | Wild radish* |

* See SPECIFIC WEED PROBLEMS for more information.

**Partial control: A visual reduction of weed population as well as a significant loss of vigor for individual weed plants. For better results, use 0.5 or 0.6 ounce HARMONY® GT per acre and include a tank mix partner such as 2,4-D, MCPA, Bromoxynil (such as "Buctril" EC or "Bronate") or "Banvel"/"Banvel" SGF/"Clarity" (refer to TANK MIXTURES).

SPECIFIC WEED PROBLEMS

Common chickweed and wild buckwheat: For best results, apply a minimum of 0.5 ounce HARMONY® GT per acre plus surfactant when all or the majority of weeds have germinated and are past the cotyledon stage. Weeds should be less than 3 inches tall or across at the time of HARMONY® GT application.

Kochia, Russian thistle, Prickly lettuce: Naturally occurring resistant biotypes of these weeds are known to occur. For best results, use HARMONY® GT in a tank mix with "Banvel"/"Banvel" SGF/"Clarity" and 2,4-D, or Bromoxynil (such as "Buctril") and 2,4-D (3/4 - 1 pint

7/15

"Buctril" + 1/4 - 3/8 lb active 2,4-D ester). HARMONY® GT should be applied in the spring when kochia, Russian thistle, and prickly lettuce are less than 2" tall or 2" across and are actively growing (refer to the Tank Mixtures section of this label for additional details).

Tansymustard: For best results, use 0.5 ounce to 0.6 ounce HARMONY® GT per acre plus 2,4-D or MCPA. Refer to TANK MIXTURES for more information.

Wild garlic: For best results, apply 0.5 to 0.6 ounce HARMONY® GT per acre plus surfactant when wild garlic plants are less than 12 inches tall with 2 to 4 inches of new growth. For severe infestations, use the 0.6 ounce per acre rate of HARMONY® GT. Control may be reduced when plants are hardened-off by cold weather and/or drought stress. Control is enhanced when applications are made during warm temperatures to actively growing wild garlic plants. Typical symptoms of dying wild garlic plants (discoloration and collapse) may not be noticeable for 2-5 weeks.

Thorough coverage of all garlic plants is essential.

Tank mixes of HARMONY® GT plus metribuzin may result in reduced control of wild garlic.

Wild radish: For best results, apply 0.5 to 0.6 ounce HARMONY® GT per acre plus surfactant either in the fall or spring to wild radish rosettes less than 6 inches in diameter. Applications made later than 30 days after weed emergence will result in partial control. Fall applications should be made prior to hardening-off of plants.

TANK MIXTURES

In cereals, fallow and pre-plant burndown, HARMONY® GT may be tank mixed with other suitable registered herbicides to control weeds listed as suppressed, weeds resistant to HARMONY® GT or weeds not listed under Weeds Controlled. HARMONY® GT can also be mixed with registered fungicides, insecticides, or liquid fertilizer for use on wheat, barley, oat, or fallow.

Read and follow all manufacturer's label recommendations for the companion products. If those recommendations conflict with this label, do not tank mix the product with HARMONY® GT.

With 2,4-D (amine or ester) or MCPA (amine or ester)

HARMONY® GT may be tank mixed with the amine and ester formulations 2,4-D and MCPA herbicides for use on wheat, barley, oat, or fallow.

For best results in the Red River Valley and adjacent areas of North Dakota and Minnesota, add the ester formulations of 2,4-D or MCPA herbicides to the tank at 3/8 lb active ingredient (such as 3/4 pint of a 4 lb/gal product, 1/2 pint of a 6 lb/gal product). No additional surfactant is needed with this mixture.

For best results, in other areas, add the ester formulations of 2,4-D or MCPA herbicides to the tank at 1/4 to 3/8 lb active ingredient (such as 1/2-3/4 pint of a 4 lb/gal product, 1/3-1/2 pint of a 6 lb/gal product). Surfactant may be added to the mixture at 1/2 to 1 quart per 100 gal of spray solution (0.125 to 0.25% v/v); however, adding surfactant may increase the potential for crop injury, especially at the higher phenoxy rates. Higher rates of 2,4-D or MCPA may be used, but do not exceed the highest rate allowed by those respective labels. Read and

follow all label instructions on timing, precautions, and warnings for these herbicides before using these tank mixtures.

With dicamba (such as "Banvel"/"Banvel" SGF/"Clarity")

HARMONY® GT may be tank mixed with 1/16 to 1/8 lb active ingredient dicamba (such as 2-4 fluid ounce "Banvel", 4-8 fluid ounce "Banvel" SGF, 2-4 fluid ounce "Clarity"). Use higher rates when weed infestation is heavy. Surfactant may be added to the mixture at 1/2 to 1 quart per 100 gal of spray solution (0.125 to 0.25% v/v); however, adding surfactant may increase the potential for crop injury. Refer to the specific dicamba label for application timing and restrictions. Tank mixes of HARMONY® GT plus dicamba may result in reduced control of some broadleaf weeds.

With 2,4-D (amine or ester) and "Banvel"/"Clarity"

HARMONY® GT may be applied in a 3-way tank mix with formulations of dicamba and 2,4-D. Make application of HARMONY® GT plus 1/16 to 1/8 lb active ingredient dicamba (such as 2-4 fluid ounce "Banvel", 4-8 fluid ounce "Banvel" SGF, 2-4 fluid ounce "Clarity") plus 1/4-3/8 lb active ingredient 2,4-D Ester or Amine per acre. Use higher rates when weed infestation is heavy. Surfactant may be added to the mixture at 1/2 to 1 quart per 100 gal of spray solution (0.125 to 0.25% v/v); however, adding surfactant may increase the potential for crop injury. Consult the specific 2,4-D label, dicamba label, or local recommendations for more information and restrictions. Apply this three-way combination to winter wheat and winter oat after the crop is tillering and prior to jointing (first node). In Spring Wheat (including Durum) and Spring Oat, apply after the crop is tillering and before it exceeds the 5-leaf stage. In Spring Barley, apply after the crop is tillering and before it exceeds the 4-leaf stage.

With Bromoxynil (such as "Buctril", or "Bronate")

HARMONY® GT may be tank mixed with bromoxynil containing herbicides registered for use on wheat, barley, or fallow. For best results, add bromoxynil containing herbicides to the tank at 3/16 to 3/8 lb active per acre (such as "Bronate" or "Buctril" at 3/4-1 1/2 pint per acre).

Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using these tank mixtures. Follow the most restrictive labeling. Tank mixes of HARMONY® GT plus "Buctril" may result in reduced control of Canada thistle.

With DuPont EXPRESS® or EXPRESS® XP Herbicide

HARMONY® GT may be tank mixed with EXPRESS® or EXPRESS® XP based on local recommendations. Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using this tank mixture.

With DuPont ALLY® or ALLY® XP Herbicide

HARMONY® GT may be tank mixed with ALLY® or ALLY® XP based on local recommendations. Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using this tank mixture.

With "Assert" or "Avenge" Herbicide

HARMONY® GT can be tank mixed with "Avenge" or "Assert". When tank mixing HARMONY® GT with "Assert", always include another broadleaf weed herbicide with a different mode of action (for example: 2,4-D ester, MCPA ester, "Buctril", or "Bronate"). Tank-mixed applications of

8/15

HARMONY® GT plus "Assert" may cause temporary crop discoloration, stunting, or injury when heavy rainfall occurs shortly after application.

With "Hoelon"

A tankmix of "Hoelon" 3EC herbicide + HARMONY® GT herbicide can be applied for annual ryegrass (in the Pacific Northwest only), wild oat and broadleaf weed control in winter and spring wheat, and spring barley. The "Hoelon" 3EC herbicide rate should be 2 2/3 pints per acre with up to 0.5 ounce per acre HARMONY® GT herbicide in spring and winter wheat.

A three-way tankmix of "Hoelon" 3EC herbicide + "Buctril" herbicide + HARMONY® GT herbicide can be applied for annual ryegrass (in the Pacific Northwest only), wild oat and broadleaf weed control in winter and spring wheat, and spring barley. The "Hoelon" 3EC herbicide rate should be 2 2/3 pints per acre with up to 0.5 ounce per acre HARMONY® GT herbicide in winter wheat (up to 0.4 ounce per acre in spring wheat and spring barley). "Buctril" herbicide should be used at 1 pint per acre.

This tank mixture should only be used under good soil moisture conditions when wild oats are in the 1 to 4 leaf stage. Reduced control of foxtail is likely when tank mixing "Hoelon" with HARMONY® GT herbicide. When foxtail is the major grassy weed in the field, DO NOT tank mix "Hoelon" 3EC herbicide + HARMONY® GT herbicide - Use sequential treatments. Be sure to follow all use directions, warnings and cautions on the EPA approved "Hoelon" 3EC and "Buctril" labels.

With "Achieve"

HARMONY® GT can be tankmixed with "Achieve" for wild oat control. This tankmix may also include 2,4-D ester, MCPA ester, bromoxynil or bromoxynil/MCPA for greater spectrum of broadleaf control - see "Achieve" label for specific use directions and restrictions on tank mixes.

To minimize the reduction in wild oat control, use the higher rates of "Achieve" when using rates of HARMONY® GT greater than 0.3 ounce per acre.

Note: Green foxtail, yellow foxtail, Persian darnel and other grass weeds will not be controlled by this tankmix.

Read and follow all label instructions on tank mixes, application timing, precautions, and warnings on the "Achieve" label.

With "Puma"

HARMONY® GT herbicide can be tankmixed with "Puma" 1EC for control of some annual grass weeds. This tankmix may also include MCPA ester, bromoxynil or bromoxynil/MCPA for greater spectrum of broadleaf control - see "Puma" 1EC label for specific use directions and restrictions on tank mixes.

Read and follow all label instructions on the EPA approved "Puma" 1EC label for tank mixes, application timing, precautions, and restrictions. If those recommendations conflict with this label, do not tank mix the product with HARMONY® GT herbicide.

With "Tiller"

HARMONY® GT can be tankmixed with "Tiller" for green foxtail, foxtail millets and volunteer corn control. Read and follow all label instructions on tank mixes, application timing, precautions, and warnings on the "Tiller" label.

With Other Grass Control Products

HARMONY® GT can be tankmixed with grass control products. Antagonism generally does not occur. However, DuPont recommends that you first consult your state experiment station, university, or extension agent, Agricultural dealer, or DuPont representative as to the potential for antagonism before using the mixture. If no information is available, limit the initial use of HARMONY® GT and the grass product to a small area.

With Insecticides and Fungicides

HARMONY® GT may be tank mixed or used sequentially with insecticides and fungicides registered for use on cereal grains.

However, under certain conditions (drought stress, cold weather, or if the crop is in the 2-4 leaf stage), tank mixes or sequential applications of HARMONY® GT with organophosphate insecticides (such as parathion, "Di-Syston") may produce temporary crop yellowing or, in severe cases, crop injury. The potential for crop injury is greatest when wide fluctuations in day/night temperatures occur just prior to or soon after application. Test these mixtures in a small area before treating large areas.

Do not apply HARMONY® GT within 60 days of crop emergence where an organophosphate insecticide (such as "Di-Syston") has been applied as an in-furrow treatment, as crop injury may result.

Do not use HARMONY GT plus "Malathion", as crop injury will result.

With Liquid Nitrogen Solution Fertilizer

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing HARMONY® GT in fertilizer solution.

HARMONY® GT must first be slurried with water and then added to liquid nitrogen solutions (e.g., 28-0-0, 32-0-0). Ensure that the agitator is running while the HARMONY® GT is added. Use of this mixture may result in temporary crop yellowing and stunting.

If using low rates of liquid nitrogen fertilizer in the spray solution (less than 50% of the spray solution volume), the addition of surfactant is necessary. Add surfactant at 1/2 pint - 1 quart per 100 gal of spray solution (0.06 -0.25% v/v) based on local recommendations.

When using high rates of liquid nitrogen fertilizer in the spray solution, adding surfactant increases the risk of crop injury. Consult your agricultural dealer, consultant, fieldsman, or DuPont representative for a specific recommendation before adding an adjuvant to these tank mixtures.

If 2,4-D or MCPA is included with HARMONY® GT and the fertilizer mixture, ester formulations tend to be more compatible (See manufacturer's label). Do not add surfactant when using HARMONY® GT in tank mix with 2,4-D ester or MCPA ester and liquid nitrogen fertilizer solutions.

Do not use low rates of liquid fertilizer as a substitute for a surfactant.

Do not use with liquid fertilizer solutions with a pH less than 3.0.

9/15

SOYBEANS

APPLICATION TIMING

HARMONY® GT may be applied to soybeans any time after the first trifoliolate has expanded fully. Apply no later than 60 days before harvest.

USE RATES

Make a single application of HARMONY® GT at a rate of 0.083 (1/12) ounce per acre for selective postemergence broadleaf weed control.

Note: DuPont PINNACLE® was formulated as 25% DF, HARMONY® GT is more concentrated, be certain to use rate noted above.

SPRAY ADDITIVES

Applications of HARMONY® GT in soybeans must include a nonionic surfactant or crop oil concentrate, and an ammonium nitrogen fertilizer.

Nonionic Surfactant

- Apply at the rate of 1 to 2 pt per 100 gal of spray solution (0.125%-0.25% v/v concentration of formulated product). Surfactants must contain at least 50% of the formulated product as actual nonionic surfactant. Avoid products that do not accurately define their ingredients on the product label.
- Using the higher rate of nonionic surfactant, particularly under hot, humid conditions, may result in temporary crop injury.
- Do Not Use "Dash" unless specified on other DuPont supplemental labeling.
- In the States of Arkansas and South Carolina on soybeans, use only nonionic surfactant at a rate of 0.125% V/V (1 pt/100 gal of spray solution) - unless specified on other DuPont supplemental labeling.

Crop Oil Concentrate

Under dry conditions or during cool weather, a crop oil concentrate at 4 pt/100 gal of spray solution (0.5% v/v) may be used in place of a nonionic surfactant to enhance weed control.

- Use a petroleum-based crop oil concentrate with at least 14% emulsifiers/surfactant and 80% oil.

The use of crop oil concentrate may result in temporary crop injury.

Ammonium Nitrogen Fertilizer

An ammonium nitrogen fertilizer is recommended in addition to a surfactant or a crop oil concentrate and required where velvetleaf is present.

- Use a high-quality liquid nitrogen fertilizer such as 28-0-0 or 32-0-0 at a rate of 4-8 pints per acre, or 10-34-0 at a rate of 2-4 pints per acre.
- Alternatively, a high-quality, sprayable grade of ammonium sulfate (21-0-0) may be used at a rate of 2-4 pounds per acre.
- Use the lower rate for spray volumes less than 15 gallons per acre.

CULTIVATION

A timely cultivation may be necessary to control suppressed weeds, weeds that were beyond the maximum size at the time of application, or weeds that emerge after an application of HARMONY® GT.

- Do not cultivate before, during, or within 7 days after the application.
- Cultivation may decrease weed control by pruning roots and placing the weed under stress.
- The best time to cultivate is approximately 14 days after application.

WEEDS CONTROLLED

When applied to soybeans as directed, HARMONY® GT will control the following weeds:

Weeds Controlled	Maximum Height (Inches) at Application
Annual Smartweeds	6
Lambsquarters	4
Pigweed	
Rough (red root)	12
Other species	8
Velvetleaf	6
Wild Mustard	up to 4" in dia.

Partial Control*	Maximum Height (Inches) at Application
Cocklebur	6
Jimsonweed	4
Wild Sunflower	6

*Partial Control: A visual reduction of weed population as well as a significant loss of vigor for individual weed plants.

TANK MIXTURES

In soybeans, DuPont will not warrant crop safety of HARMONY® GT tank mixtures with any other pesticide or spray adjuvant except as specified in this or other DuPont supplemental labeling.

Do not tank mix HARMONY® GT with organophosphate insecticides, or apply HARMONY® GT within 14 days before or after an application of an organophosphate insecticide, as severe crop injury may occur.

With Postemergence Grass Herbicides

HARMONY® GT may be tank mixed with postemergence grass herbicides such as DuPont ASSURE® II herbicide. Do not tank mix with "Poast Plus" unless specified on other DuPont supplemental labeling.

Under certain conditions, HARMONY® GT may reduce the activity of the postemergent grass herbicide. The broadleaf activity of HARMONY® GT will not be affected. Refer to the postemergent grass herbicide label for specific use information and precautions.

10/15

With post grass herbicides, surfactant rate (concentration) should be 1-2 pints per 100 gallons of spray solution (0.125%-0.25% v/v concentration). Use of a higher rate of nonionic surfactant, particularly under hot, humid conditions, may result in temporary crop injury. Do not use "Dash" or crop oil concentrate when tank mixing DuPont HARMONY® GT with postemergence grass herbicides unless specified on other DuPont supplemental labeling. Include a nonionic surfactant with the tank mix of HARMONY® GT and post grass herbicides such as DuPont ASSURE® II.

With "Basagran"

HARMONY® GT may be tank mixed with "Basagran" herbicide at the rate of 0.083 (1/12) ounce HARMONY® GT plus 1 pint "Basagran" per acre for control of these weeds in addition to those listed as controlled by HARMONY® GT alone:

<i>Species</i>	<i>Maximum Height (inches)</i>
cocklebur	4
jimsonweed	6
venice mallow	2
wild sunflower	4

Applications of HARMONY® GT + "Basagran" must include a nonionic surfactant or crop oil concentrate and an ammonium nitrogen fertilizer. See "Soybeans - Spray Additives" section.

With "Galaxy"

HARMONY® GT herbicide at 0.083 (1/12) ounce may be tankmixed with "Galaxy" herbicide at 2 pints per acre for improved control of black nightshade. Consult the "Galaxy" label for additional weeds controlled by "Galaxy". Best results are obtained when the HARMONY® GT + "Galaxy" tankmix is applied to weeds that are young and actively growing and before weeds exceed the size limits on the respective labels.

Applications of HARMONY® GT + "Galaxy" must include a non-ionic surfactant or crop oil concentrate and an ammonium nitrogen fertilizer. See Soybeans - Spray Additives section. Use of the higher rate of non-ionic surfactant, particularly under hot, humid conditions may increase temporary crop injury.

Considerable early season crop injury may result from applications of this tank mix. The potential for adverse crop response is most pronounced during hot, humid conditions, under widely fluctuating climactic conditions, or with applications to soybeans under stress. Symptoms may appear as, but are not limited to, leaf speckling, leaf bronzing, and/or plant stunting.

SEQUENTIAL APPLICATIONS

HARMONY® GT following "Pursuit"

HARMONY® GT may be used as a sequential treatment to control newly emerged weeds following a soil application (preemerge, preplant, or preplant incorporated) of "Pursuit" or imazethapyr-containing products.

Sequential applications of HARMONY® GT following postemergent "Pursuit" treatments are not recommended because:

- Crop injury from sequential postemergence applications of HARMONY® GT following "Pursuit" is greater than from the use of either product applied alone. The first application interferes with the soybean plant's ability to metabolize the second herbicide treatment. Sequential applications may result in severe crop injury.
- Any weeds not controlled by the "Pursuit" application will be stressed at the time of the sequential treatment. This will result in unsatisfactory weed control, particularly for stress sensitive weeds such as lambsquarters.
- Weeds that have recovered from a "Pursuit" application will typically be larger than labeled size by the time soybeans may be safely treated with a HARMONY® GT application. This will result in unsatisfactory weed control.

Even though not recommended for sequential application, a minimum interval of at least 14 days between applications of HARMONY® GT following "Pursuit" is advised to reduce the potential for crop injury and unsatisfactory weed control. The soybeans should be free from stress (herbicide or environmental) and actively growing. Weeds should be free from stress and not exceed the labeled size (height) at the time of HARMONY® GT application.

ENVIRONMENTAL CONDITIONS

Applications made during or immediately following periods of abnormally cold weather for soybeans may result in less than satisfactory weed control.

Poor weed control or crop injury may result from applications made to plants under stress from:

- abnormal hot or cold weather,
- growing conditions such as drought or water-saturated soil,
- soil nutrient deficiencies such as iron chlorosis,
- disease,
- injury from cultivation,
- nematode, insect, or prior herbicide injury.

Delay application until stress passes and weeds and soybeans resume growth. Severe stress from conditions immediately following application may also result in crop injury or poor weed control.

Applications during periods of hot and humid weather increase the risk of crop injury.

Wilting, temporary leaf yellowing, reddened veins, and/or growth retardation of soybeans may follow application of HARMONY® GT. The growth retardation is generally in the form of shortened internode spacing. These effects will generally be most evident 5-7 days after application. The soybeans will recover quickly under favorable growing conditions.

GENERAL USE AND APPLICATION DIRECTIONS - ALL CROPS

GROUND APPLICATION

Wheat, Barley, Oat and Fallow

- For flat-fan nozzles, use a spray volume of at least 5 gal per acre (GPA).
- For flood nozzles on 30" spacings, use at least 10 GPA, flood nozzles no larger than TK10 (or the equivalent), and a pressure of at least 30 psi. For 40" nozzle spacings, use at least 13 GPA; for 60" spacings use at least 20 GPA. It is essential to overlap the nozzles 100% for all spacings.
- Raindrop "RA" nozzles are not recommended for HARMONY® GT applications, as weed control performance may be reduced.
- Use screens that are 50-mesh or larger.

Soybeans

Broadcast Application

- Use 10-25 gallons of water per acre.
- Use flat fan nozzles at 25-60 psi.
- Under heavy weed pressure or dense crop foliage, increase minimum spray volume to 15-25 gal per acre.
- Do not use flood, hollow cone, rain drop, whirl chamber, or controlled droplet applicator (CDA) type nozzles. Unacceptable crop injury, excessive spray drift, or poor weed control may result.
- For proper spray coverage, adjust the boom and nozzle height according to the specifications listed by the nozzle manufacturer.

Band Application

- For band application, use proportionately less spray mixture.
- To avoid crop injury, carefully calibrate the band applicator not to exceed the labeled rate.
- Carefully follow the manufacturer's instructions for nozzle types (flat fan nozzles preferred), nozzle orientation, distance of the nozzles from the crop and weeds, spray volumes, calibration, and spray pressure.
- For additional information on row banders, see DuPont bulletin, "Application Accuracy Row Banders."

AERIAL APPLICATION

Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage.

Wheat, Barley, Oat, and Fallow

- use 2 to 5 GPA
- Use at least 3 GPA in Idaho, Oregon, or Utah

Soybeans

- Use a minimum of 5 GPA.

When applying HARMONY® GT by air in areas adjacent to sensitive crops, use solid stream nozzles oriented straight back. Adjust the swath to avoid spray drift damage to sensitive crops downwind and/or use ground equipment to treat the border edge of fields. See the Spray Drift Management section of this label.

ENVIRONMENTAL CONDITIONS

If rain is expected within 1 hour, do not apply HARMONY® GT, or weed control may be decreased.

CROP ROTATION

Wheat, barley, oat, soybeans and field corn may be replanted anytime after the application of HARMONY® GT. Any other crop may be planted 45 days after the application of HARMONY® GT.

GRAZING

Cereals and Soybeans

Do not graze or feed forage or hay from treated areas to livestock (harvested straw may be used for bedding and/or feed).

MIXING INSTRUCTIONS

Do not use with spray additives that alter the pH of the spray solution below pH 5.0 or above pH 9.0, as rapid product degradation can occur. Spray solutions of pH 6.0 - 8.0 allow for optimum stability of HARMONY® GT.

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of HARMONY® GT.
3. Continue agitation until the HARMONY® GT is fully dispersed, at least 5 minutes.
4. Once the HARMONY® GT is fully dispersed, maintain agitation and continue filling tank with water. HARMONY® GT should be thoroughly mixed with water before adding any other material.
5. As the tank is filling, add tank mix partners (if desired) then add the required volume of nonionic surfactant. Always add surfactant last.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Apply HARMONY® GT spray mixture within 24 hours of mixing to avoid product degradation.
8. If HARMONY® GT and a tank mix partner are to be applied in multiple loads, pre-slurry the HARMONY® GT in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the HARMONY® GT.

SPRAY EQUIPMENT

For specific application equipment, refer to the manufacturer's recommendations for additional information on GPA, pressure, speed, nozzle types and arrangements, nozzle heights above the target canopy, etc.

Be sure to calibrate air or ground equipment properly before application. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern with minimum drift. Use higher spray volumes to obtain better coverage when crop canopy is dense. Avoid swath overlapping, and shut off spray booms while starting, turning, slowing, or stopping, to avoid injury to the crop.

12/15

Do not make applications using equipment and/or spray volumes or during weather conditions that might cause spray to drift onto nontarget sites. For additional information on spray drift refer to Spray Drift Management section of label. Continuous agitation is required to keep HARMONY® GT in suspension.

SPRAYER CLEANUP

The spray equipment must be cleaned before HARMONY® GT is sprayed. Follow the cleanup procedures specified on the labels of the previously applied products. If no directions are provided, follow the six steps outlined in After Spraying HARMONY® GT.

AT THE END OF THE DAY

It is recommended that during periods when multiple loads of HARMONY® GT herbicide are applied, at the end of each day of spraying the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits which can accumulate in the application equipment.

AFTER SPRAYING HARMONY® GT AND BEFORE SPRAYING CROPS OTHER THAN WHEAT, BARLEY, OAT AND SOYBEANS

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of HARMONY® GT as follows:

1. Drain tank; thoroughly rinse spray tanks, boom, and hoses with clean water. Loosen and physically remove any visible deposits.
2. Fill the tank with clean water and 1 gal of household ammonia* (contains 3% active) for every 100 gal of water. Flush the hoses, boom, and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 min. Flush the hoses, boom, and nozzles again with the cleaning solution, and then drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat step 2.
5. Rinse the tank, boom, and hoses with clean water.
6. If only Ammonia is used as a cleaner, the rinsate solution may be applied back to the crop(s) recommended on this label. Do not exceed the maximum labeled use rate. If other cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

* Equivalent amounts of an alternate-strength ammonia solution or a DuPont-approved cleaner can be used in the cleanout procedure. Carefully read and follow the individual cleaner instructions. Consult your Ag dealer, applicator, or DuPont representative for a listing of approved cleaners.

Notes:

1. **CAUTION:** Do not use chlorine bleach with ammonia as dangerous gases will form. Do not clean equipment in an enclosed area.

2. Steam-cleaning aerial spray tanks is recommended prior to performing the above cleanout procedure to facilitate the removal of any caked deposits.
3. When HARMONY® GT is tank mixed with other pesticides, all cleanout procedures should be examined and the most rigorous procedure should be followed.
4. In addition to this cleanout procedure, all precleanout guidelines on subsequently applied products should be followed as per the individual labels.
5. Where routine spraying practices include shared equipment frequently being switched between applications of HARMONY® GT and applications of other pesticides to HARMONY® GT -sensitive crops during the same spray season, it is recommended that a sprayer be dedicated to HARMONY® GT to further reduce the chance of crop injury.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150 - 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See Wind, Temperature and Humidity, and Temperature Inversions sections of this label.

Controlling Droplet Size - General Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - The boom length should not exceed 3/4 of the wing or rotor length - longer booms increase drift potential.
- **Application Height** - Application more than 10 ft above the canopy increases the potential for spray drift.

BOOM HEIGHT

Setting the boom at the lowest referenced height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID GUSTY AND WINDLESS CONDITIONS.**

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring.

Note: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the spray equipment section of this label to determine if use of an air assist sprayer is recommended.

RESISTANCE

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes. If applicable, see the Weeds Controlled section of this label for additional information on managing herbicide resistant weed biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

Naturally occurring weed biotypes that are resistant to "Amber" herbicide, "Pursuit" herbicide, DuPont™ ALLY® herbicide, DuPont™ GLEAN® FC herbicide, DuPont™ EXPRESS® herbicide, or DuPont™ HARMONY® EXTRA herbicide will also be resistant to HARMONY® GT.

14/15

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

PRECAUTIONS

- Injury to or loss of desirable trees or vegetation may result from failure to observe the following:
 - Do not apply, drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
 - Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants
- Injury to or loss of adjacent sensitive crops and vegetation may result from failure to observe the following:
 - Take all necessary precautions to avoid all direct or indirect contact (such as spray drift) with non-target plants or areas.
 - Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than wheat, barley or oat.
- Wheat, barley, oat and soybean varieties may differ in their response to various herbicides. DuPont recommends that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use of HARMONY® GT to a small area.
- Do not apply to wheat, barley, oat or soybeans that are stressed by severe weather, drought (including low levels of subsoil moisture), water-saturated soil, disease or insect damage as crop injury may result. Under certain conditions such as prolonged cool weather (daily high temperature less than 50 Deg.F.) or wide fluctuations in day/night temperatures just prior to, during or soon after treatment, temporary yellowing and/or crop stunting may occur. Risk of injury is greatest when the cereal crop is in the 2 to 5-leaf stage.
- Do not apply to wheat, barley or oat crops underseeded with another crop.
- For ground applications applied to weeds when dry, dusty field conditions exist, control of weeds in wheel track areas may be reduced.

STORAGE AND DISPOSAL

Storage: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a cool, dry place.

Product Disposal: Do not contaminate water, food, or feed by disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: For Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **For Fiber Sacks:** Completely empty fiber sack by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into manufacturing or application equipment. Then dispose of sack in a sanitary landfill or by incineration if allowed by State and local authorities. **For Fiber Drums With Liners:** Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner. **For Bags Containing Water Soluble Packets:** Do not reuse the outer box or the resealable plastic bag. When all water-soluble packets are used, the outer packaging should be clean and may be disposed of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by open burning. If burned, stay out of smoke. If the resealable plastic bag contacts the formulated product in any way, the bag must be triple-rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer wrap as described above. **For Metal Containers (non aerosol):** Triple rinse (or equivalent) the container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. **For Paper and Plastic Bags:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval, DuPont™, HARMONY®, EXPRESS®, ALLY®, GLEAN®, PINNACLE® and ASSURE® are trademarks or registered trademarks of DuPont or its affiliates

"Banvel", "Clarity", "Dash", "Poast Plus", "Basagran" and "Galaxy" are registered trademarks of BASF Corporation.

"Buctril" and "Bronate" are registered trademarks of Rhonc-Poulenc Ag Company.

"Amber" is a registered trademark of Novartis.

"RA" is a registered trademark of Delavan Corporation.

"Hoelon", "Puma" and "Tiller" are registered trademarks of AgrEvo USA Company.

"Assert", "Avenge", and "Pursuit" are registered trademarks of American Cyanamid Company.

"Landmaster", "Fallow Master", and "Roundup" are registered trademarks of Monsanto.

"Di-Syston" is a registered trademark of Bayer Corporation.

"Malathion" is a registered trademark of United Ag Products.

"Achieve", "Dyfonate", and "Force" are registered trademarks of Zeneca Ag Products

"Lorsban" is a registered trademark of Dow AgroScience

D - 911 080103

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants.

DuPont does not agree to be an insurer of these risks. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT

Internet address: <http://cropprotection.dupont.com/>

© 2000-2003 E. I. du Pont de Nemours and Company, Wilmington, Delaware 19898.

All rights reserved.