



AGRICULTURAL BULLETIN

1053
250007
DUPONT

BENLATE® 50 DF™ FUNGICIDE
CARROTS

SUPPLEMENTAL LABELING
EPA REG. NO. 352-447

BENLATE® 50 DF™ FUNGICIDE
FOR CONTROL OF
WHITE MOLD ON CARROTS

ACCEPTED
MAY 12 1987
U.S. DEPARTMENT OF AGRICULTURE
Pest Control Administration
Registration Division
EPA Reg. No. 352-447

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Du Pont "Benlate" 50 DF™ Fungicide is recommended for use on carrots as a foliar spray to control white mold (Sclerotinia).

HOW TO USE

Apply 1/4 to 1 lb. "Benlate" 50 DF™ per acre using sufficient water to obtain thorough coverage (3 to 10 gallons per acre for aerial application). Begin applications when disease first appears and repeat at 7 to 10 day intervals. Under severe disease conditions use the higher rate and shorter interval.

Do not apply within 4 days of harvest.

Note: Do not tank mix "Benlate" 50 DF™ with lime or alkaline pesticides such as Bordeaux mixture or lime sulfur.

IMPORTANT

BEFORE USING "BENLATE" 50 DF™, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on the package label. Follow the instructions carefully.

This application must be in the possession of the user at the time of pesticide application.

012787

1/82



AGRICULTURAL BULLETIN



BENLATE® 50 DF™ FUNGICIDE
WHEAT

ACCEPTED
MAY 12 1987
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 352-447

SUPPLEMENTAL LABELING
EPA REG. NO. 352-447

BENLATE® 50 DF™ FUNGICIDE
FOR CONTROL OF
CERTAIN DISEASES OF WHEAT

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Du Pont "Benlate" 50 DF™ Fungicide is recommended for use as a foliar spray for the control of Cercospora Foot Rot, Powdery Mildew and Leaf and Glume Blotch of wheat.

HOW TO USE

Apply "Benlate" 50 DF™ at the following rates, using sufficient water to provide thorough and uniform coverage; for aerial application, use 5 to 10 gals. per acre. Use the higher dosage rate under conditions of severe disease pressure. Do not apply within 21 days of harvest. Do not allow livestock to forage in treated fields.

Cercospora Foot Rot - (Strawbreaker Foot Rot) - For early fall seeded winter wheat, make a single application of 1 to 2 lbs. per acre in the fall or spring when wheat is fully tillered but before elongation has occurred.

Powdery Mildew - Apply 1/2 to 1 lb. per acre at the boot stage; repeat 14 to 21 days later. A nonphytotoxic superior-type spray oil (60 to 70 second viscosity) may be added at the rate of 1 qt. per acre to improve spray coverage; add as last ingredient to spray tank.

Leaf and Glume Blotch - Apply 1/2 to 1 lb. of "Benlate" 50 DF™ per acre before low levels of disease spread to cause serious crop damage. Repeat 14 days later. Applications should be made at late boot to early heading stage to keep the flag leaf and emerging seedhead free of disease. Do not make more than a total of three applications. Treatments made after the disease is severe or after the flag leaf becomes infected may not provide economical disease control. A nonphytotoxic, superior-type spray oil (60 to 70 second viscosity) at the rate of 1 quart per acre, or a "spreader-sticker" spray adjuvant may be added to improve spray coverage; add oil or adjuvant as last ingredient to spray tank.

2

IMPORTANT

BEFORE USING "BENIATE" 50 DFTM, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental information for use of this product which may not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

120886

50 DF
BENIATE