DRAFT, NOT APPROVED

"LOROX" PLUS HEPBICIDE EPA REG.NO. 354-451 ACCEPTED

APR 13 1989

The fact the Federal Insection for the period on the period for the period for the period for the period on the period of the period

SUPPLEMENTAL LABELING

"LOROX" PLUS HERBICIDE

REDUCED RATE FOLIOWED BY "CLASSIC" HERBICIDE

FOR BROADLEAF WEED CONTROL IN SOYBEANS

Du Pont "Lorox" Plus Herbicide is recommended preplant incomorated or preemergence at reduced rates followed by "Classic" Herbici e postemergence for broadleaf weed control in soybeans. This treatment will control those broadleaf weeds listed on the "Lorox" Plus and "Classic" Herbicide product labels.

DIRECTIONS FOR USE

Apply "Lorox" Plus preplant incorporated or preemergence at 12 ounces product per acre to any soil texture with 1/2 to 3% organic matter as directed on the product label. Follow the "Lorox" Plus preplant incorporated or preemergence reduced rate with a postemergence application of "Classic" Herbicide at its labeled rate when weeds are small and actively growing. (Usually 14-28 days after preplant incorporated or preemergence application of "Lorox" Plus reduced rate.)

For grass weed control "Lorox" Plus can be tank mixed with Dual[2], Lasso[3], Surflan[4], Treflan[4] and Sonalan[5]. Consult labels for rates and use instructions of these products.

IMPORTANT

Before using "Lorox" Plus and "Classic" herbicide, read and carefully note the cautionary statements and all other information appearing on the product labels.

This bulletin contains new or supplemental information for use of theesproducts which does not appear on the product label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

[4] Registered trademark of Elanco Products Company

E. I. Du Pont de Nemours & Co. (Inc.) Agricultural Products Department Wilmington, Delaware 19880-0038