




DuPont Agricultural Products

"..... A Growing Partnership With Nature"



EXTRAZINE® II 4L HERBICIDE PLUS "MARKSMAN" OR "BANVEL" HERBICIDE TANKMIXES FOR POSTEMERGE USE ON FIELD CORN IN THE STATES OF INDIANA AND OHIO

EXTRAZINE® II 4L HERBICIDE

(EPA Reg. No. 352-500)

POSTEMERGE TANKMIXES WITH "MARKSMAN" OR "BANVEL" HERBICIDES FOR USE ON FIELD CORN IN THE STATES OF INDIANA AND OHIO

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont EXTRAZINE II 4L herbicide may be applied in a tankmix combination with "Marksman" or "Banvel" for control of certain grasses and broadleaf weeds in field corn. Addition of "Marksman" or "Banvel" will improve control of difficult broadleaf weeds including triazine-resistant lambsquarters.

WHEN TO APPLY

Under warm, sunny, dry conditions of low humidity, an EXTRAZINE II 4L + "Marksman" or "Banvel" tankmix may be applied Postemergence in field corn.


Apply from crop emergence through the three-leaf stage of corn growth, but before weeds exceed about 1.5 inches in height. Do not apply over the top of corn if the fourth leaf is visible.

HOW TO USE

The use rate of this tank mix is 2 qts/acre of EXTRAZINE II 4L plus 3.5 pts/acre of "Marksman" or 1 pt/acre of "Banvel". To load, mix, and apply these tankmixes, follow all instructions in the federal EXTRAZINE II 4L label under the "General Mixing and Spraying Directions" section.

- Apply these tankmixes in water only
• Do not use a surfactant, crop oil or any other adjuvant.
• Do not spray emerged corn plants using a liquid nitrogen carrier

Do not use on sands or loamy sands as unacceptable injury may occur.



APPLICATION INFORMATION

Refer to the respective labels for application information on application methods, gallonage, nozzles, pressures, and equipment. The most restrictive labeling shall apply.

All federal guidelines for protective clothing, precautions, and use requirements for EXTRAZINE II 4L must be followed.

PRECAUTIONS

Refer to "Marksman" label for atrazine rate limits and precautions on rotation crops.

When applied as a postemergence herbicide, EXTRAZINE II 4L is active through foliage as well as through roots. Yellowing and/or stunting of the corn may result from this treatment, particularly if cold, adverse growing conditions occur after application. Do not apply this EXTRAZINE II 4L tankmix under cold (daytime high <55 degrees F), wet weather conditions or to corn growing under stress caused by weather, insects, diseases etc. Extended or extreme cold and wet conditions may reduce stands.

IMPORTANT

BEFORE USING "EXTRAZINE" II 4L, MARKSMAN, OR BANVEL, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABELS.

This bulletin contains new or supplemental instructions for use of these products which do not appear on the EPA-registered package labels. Follow the instructions carefully.

This bulletin must be in the possession of the user at the time of pesticide application.

"Marksman" and "Banvel" are registered trademarks of Sandoz Ltd.

013196