

lot 2  
1/10/86  
1000

ACCEPTED  
DEC 19 1986  
U.S. DEPT. OF COMMERCE  
FEDERAL BUREAU OF INVESTIGATION  
WASHINGTON, D.C. 20535  
EPA Reg. No. 352-507

SPECIMEN LABEL  
86% OF ACTUAL SIZE



Krovar® I D F™

Directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

**IMPORTANT**—Injury to or loss of desirable trees or other plants may result from failure to observe the following.

Do not apply (except as recommended for crop use), or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend or in locations where their material may be washed or moved into contact with their roots. Do not use on lawns, walks, drive ways, tennis courts or similar areas. Do not use in home fruit plantings nor in citrus orchards interplanted to other trees or desirable plants. Prevent drift of dry powder or spray to desirable plants. Do not contaminate any body of water. Keep from contact with fertilizers, insecticides, fungicides and seeds.

Thoroughly clean all traces of "Krovar" I DF from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

**GENERAL INFORMATION**

Du Pont "Krovar" I DF Herbicide is a dispersible granule to be mixed in water and applied as a spray for selective control of weeds in citrus and for non-crop weed control. "Krovar" I DF controls annual weeds such as barnyardgrass (watergrass), chickweed, crabgrass, finger fleabane, Florida pusley, foxtail, groundsel, horseweed, johnsongrass seedlings, jungerice, lambsquarters, natagrass (redtop), nightshade (annual), pigweed, pineappleweed, puncturevine, purslane, ragweed, sandbur (sandspur), shepherdspurse, smoothstille (annual), spanishneedles, wild lettuce, and wild mustard. At higher use rates, treatment controls certain perennial weeds such as balsamapple vine seedlings, bermudagrass, heart leaf, drymary, guineagrass and milkweed (strangler), vine seedlings, and provides suppression of established balsamapple vine and milkweed (strangler) vine. Effects on perennial weeds are slow to appear, usually progressing over a period of several months. Moisture is necessary to activate the herbicide, best results are obtained if treatment is made to moist soil, and moisture is supplied by rainfall or sprinkler irrigation within two weeks after application. The degree and duration of control will vary with the amount applied, soil texture, rainfall, and other conditions.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Du Pont "Krovar" I DF should be used only in accordance with recommendations on this label or in separate published Du Pont recommendations available through local dealers.

Du Pont will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by Du Pont. User assumes all risk associated with such non-recommended use.

Farm workers performing hand labor operations should not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops not treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to farm workers performing hand labor operations who are expected to be in a treated cropland area or in a cropland area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Warnings must include the following information: "CAUTION! Area treated with 'Krovar' I DF (on date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of contact, flush eyes with plenty of water; wash skin with plenty of soap and water. Get medical attention if irritation persists."

**CITRUS**

Apply any time of year provided overhead moisture (rainfall or sprinkler irrigation) is available to activate the herbicide, preferably just before or just after weeds have germinated.

Use only where trees have been established in the orchard for at least three years (except in Florida and Texas). Apply with a properly calibrated fixed-boom power sprayer as a band or broadcast treatment beneath and/or between trees. Use sufficient water (min. 40 gals per acre) to obtain uniform coverage, avoid contact of foliage and fruit with spray or mist. Avoid overlapping and shut off spray boom when starting, turning, slowing or stopping or injury to trees may result. Continuous agitation in the spray tank is required to keep the material in suspension. Best results are obtained if application is made shortly before or shortly after acid growth begins; if dense growth is present, remove tops and spray ground. Do not graze cattle in treated areas. Dosages are expressed as broadcast rates, for band treatment use proportionately less.

© 1977 E. I. du Pont de Nemours & Co.

4-6 lbs per acre on finer soils (silt loams, clay loams, or soils with organic matter of 2% or more); use the higher rates for maximum suppression of perennials. Alternately, make two applications per year at rates of 2 lbs per acre on coarser soils and 3 lbs per acre on finer soils. Make the second application when needed to maintain weed control.

**Florida and Texas**

**Trees Established Less Than One Year:**  
Apply 2-4 lbs "Krovar" I DF per acre as needed to maintain weed control. Do not apply at less than 60-day intervals. Do not apply more than 6 lbs per acre during any six-month period or more than 8 lbs per acre during the first year.

**Trees Established One or Two Years:**  
Apply 2-4 lbs "Krovar" I DF per acre. A second application may be made when needed to maintain weed control, but do not exceed 8 lbs per acre.

**Trees Established Three or More Years:**  
Make one or two applications of 4-8 lbs per acre. Make the second application when needed to maintain weed control. Do not apply more than 16 lbs per acre per year.

**Tank Mixtures:**

Tank mixtures of "Krovar" I DF and paraquat are recommended for weed control in citrus. For rates of individual products, refer to the respective product label. Do not apply tank mixture to orange or green wood of citrus, as injury may result.

\*Texas—Use 2-4 lbs per acre on coarser soils (sands, loamy sands, sandy loams) and 4-6 lbs per acre on finer soils (silt loams, clay loams, or soils with organic matter of 2% or more). Use the higher rate for maximum suppression of perennials.

Note: Temporary yellowing of citrus leaves may occur following treatment. Because injury to citrus trees may result, do not use on soils with less than 1% organic matter (1% in Florida), poorly drained soils, gravelly soils, nor thinly covered or exposed subsoils, do not treat trees planted in irrigation furrows, do not treat diseased trees such as those with foot rot.

Do not re-plant treated areas to any crop within two years after last application as injury to subsequent crops may result, except that citrus trees may be planted one year after the last application.

**E. I. du Pont de Nemours & Co. (Inc.), Agricultural Products Department, Wilmington, DE**