

Transferred from
352-502

352-505

PM 25

1 of 2

ACCEPTED

BEST AVAILABLE COPY

SPECIMEN LABEL
86% OF ACTUAL SIZE


DEC 19 1986

352-502



Krovar® I DF™

HERBICIDE





KEEP OUT OF REACH OF CHILDREN
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS

DANGER! CAUSES SKIN DAMAGE

Do not get in eyes, dusting on clothing. Harmful if swallowed. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling.

STATEMENT OF PRACTICAL TREATMENT

If on skin: Wash with plenty of soap and water. Get medical attention.

If in eye: Flush with plenty of water. Get medical attention if irritation persists.

If swallowed: Call your doctor or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

For medical emergencies involving this product, call toll free 1-800-441-3637.

ENVIRONMENTAL HAZARDS

(Use only if directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of waste.)

IMPORTANT--Injury to or loss of desirable trees or other plants may result from failure to observe the following

Do not apply (except as directed) to crops used, or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend. Do not apply to lawns, lawns, walks, driveways, tennis courts, swimming pools, or to areas where people or animals may be present. Do not use in home lawns, gardens, or on citrus orchards or plants interplanted to other trees or desirable plants. Prevent drift of dry powder or spray, to desirable plants. Do not contaminate any food, feed, or feed ingredients from contact with fertilizers, insecticides, fungicides and seeds.

Thoroughly clean all parts of sprayer. I/D of from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle(s) and screens (clean these parts separately).

GENERAL INFORMATION

[illegible]

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont™ Kroyal™ 100 should be used only in accordance with recommendations on this label, or in separate published DuPont recommendations available through local dealers.

DuPont will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by DuPont. User assumes all risk associated with use for non-recommended use.

Farm workers performing plant and other operations should not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to farm workers performing hand labor of operations who are expected to be in a treated crop/land area or in a crop/land area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Such warnings must be given if there is a reasonable risk that written warnings cannot be understood by workers. Warnings must include the following information: (A) If the Area is treated with "Krovel" (1-4% rate of application). Do not enter. Without appropriate protection (such as clothing and sprays) have died. In case of only 1" of water with plenty of water, wash skin with plenty of soap and water. Get medical attention if irritation occurs.

CITRUS

Apply any time of year, but overhead moisture (rainfall or sprinkler irrigation) is available to activate the herbicide, preferably just before or just after annuals have germinated.

Use only when the tree is at least 18 inches in diameter and has at least three vigorous sprouts per foot of trunk (1 inch to 1 foot). Apply with a properly calibrated backpack sprayer or hand-held sprayer. For best results, highest benefit and/or best control, use 49 gallons of water per acre to obtain uniform coverage. Avoid contact with leaves and fruit with the spray. Avoid overlapping or runoff of the spray when watering, burning, blowing or shipping or injury to the trees may result. Continue agitation in this spray. The tree is required to keep the material in suspension. Test results are obtained and application is made shortly before or shortly after a hard frost begins. If dense conditions prevail, remove tops and spray ground. Do not graze cattle in treated areas. Dosages are expressed as broadcast rates. For hand treatment, use proportionately less.

1. *Chrysomelids*

SPECIMEN
73% OF ACTUAL

BEST AVAILABLE COPY

Copyright Clearance Center, Inc.

Annual Weeds

California, Arizona—Best results occur when applied in late fall or early winter but before winter annuals become well established. Application should be made after the first fall or early winter rains have settled the soil.

For the initial treatment, apply 4-5 lbs per acre on coarse soils containing 1-2% organic matter and 5-6 lbs per acre on fine-textured soils with organic matter of 2% or more and repeat annually as needed. Alternatively, apply 3-4 lbs/acre in the fall and repeat at 2-4 lbs per acre in the spring. Do not exceed 6 lbs per acre per year. Use the higher range of rates where ground-sel or puncture-weeds are known to be a problem. These higher rates will also suppress low density stands of bermudagrass and yellow nuttall. Repeat annually.

Louisiana—Make a single application of 2-4 lbs per acre on coarser soils (sands, loamy sands, sandy loams) and 4-6 lbs per acre on finer soils (silt loams, clay loams, or soils with organic matter of 2% or more); use the higher rates for maximum suppression of perennials. Alternatively, make two applications per year at rates of 2 lbs/acre on coarser soils and 3 lbs per acre on finer soils. Make the second application when needed to maintain weed control.

Florida and Texas

Trees Established Less Than One Year:

Apply 2.4 lbs. Krovar[®] 1 DF per acre as needed to maintain weed control. Do not apply at less than 60 day intervals. Do not apply more than 6 lbs per acre during any six month period or more than 8 lbs per acre during the full year.

Trees Established One or Two Years:

Apply 2 lbs. Krazar 1 DF per acre. A second application may be made when needed to maintain weed control, but do not exceed 8 lbs. per acre.

Trees Established Three or More Years:

Make one or two applications of 4-8 lbs per acre. Make the second application when needed to maintain weed control. Do not apply more than 16 lbs per acre in a year.

Isaiah 61:1-3

Tank mixtures of Kroxat 1 DF and paraquat are recommended for weed control in citrus. For rates of individual products, refer to the respective product label. Do not apply tank mixture to foliage or green wood of citrus, as injury may result.

Texas. Use 2-4 lb. per acre on coarser soils (sands, sandy loams, etc.), and 4-6 lbs. per acre on finer soils (silt loams, clay loams, or soils with organic matter of 2% or more). Use the higher rate for maximum and best control of weeds.

Note: Temporary yellowing of citrus leaves may occur following treatment. Because injury to citrus trees may result, do not use on soils with less than 1% organic matter (1% in Florida), poorly drained soils, gravelly soils, not thinly covered or exposed subsoils, do not treat trees planted in irrigation furrows, do not treat diseased trees such as those with foot rot.

Do not re-treat treated areas to any crop within two years after last application as injury to subsequent crops may result except that citrus trees may be replanted one year after the last application.

E. I. du Pont de Nemours & Co. (Inc.), Agricultural Products Department, Wilmington, DE