

352-505

12/8/2005

Page 17 13



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

AND

OFFICE OF  
PREVENTION, PESTICIDES  
TOXIC SUBSTANCES

DEC - 8 2005

Mr. J. H. Cain  
E.I. DuPont de Nemours & Company, Inc  
DuPont Crop Protection  
Stine-Haskell Research Center  
P. O. Box 30  
Newark, DE 19714-0030

Dear Mr. Cain:

Subject: DuPont Krovar 1 DF Herbicide (Amend Label Based on Diuron RED)  
EPA Registration No. 352-505  
Application Dated October 17, 2005

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable provided you make the following change before you release the product for shipment.

--In your Limitation of Warranty and Liability revise the last sentence of your second paragraph to read "**To the fullest extent permitted by law**, when you buy or use this product, you accept these risks.--

Submit one copy pf your final printed labeling incorporating the above changes before you release the product for shipment. Amended labeling supercedes all previously accepted ones. A stamped copy of labeling is enclosed for your records.

Sincerely,

*Dickie K Walters for*  
James A. Tompkins  
Product Manager 25  
Herbicide Branch  
Registration Division (7505C)

28 H-13




# DuPont™ Krovar® I DF

herbicide

---

## DRAFT LABEL



*“..... A Growing Partnership With Nature”*

**TABLE OF CONTENTS**

ACTIVE INGREDIENT .....1

PRECAUTIONARY STATEMENTS .....1

    First Aid .....1

    Hazards to Humans and Domestic Animals .....1

    Personal Protective Equipment .....1

    Engineering Control Statement .....1

    User Safety Recommendations .....1

    Environmental Hazards .....1

GENERAL INFORMATION .....2

    Use Precautions and Restrictions .....2

    Crop Rotation Bioassay .....2

    Resistance .....2

    Integrated Pest Management .....2

DIRECTIONS FOR USE .....2

AGRICULTURAL USES .....3

    Agricultural Use Requirements .....3

APPLICATION INFORMATION .....3

TANK MIXTURES .....3

WEEDS CONTROLLED .....3

CITRUS .....3

    California, Arizona .....4

    Florida .....4

    Louisiana .....4

    Texas .....4

SPRAY PREPARATION .....4

NON-AGRICULTURAL USES .....4

    Non-Agricultural Use Requirements .....4

USE RESTRICTIONS - STATE OF FLORIDA .....4

APPLICATION INFORMATION .....4

NON-CROP WEED CONTROL .....5

    Application Timing .....5

    Application Rates .....5

    Weeds Controlled .....5

    Retreating .....6

    Small Areas .....6

    Tank Mixtures .....6

SPECIAL USES .....6

    Under Asphalt and Concrete Pavement .....6

        Application Information .....6

        Application Timing .....6

        Application Rates .....6

        Tank Mixtures .....6

    Important Precautions - Under Asphalt Only .....6

SPRAY PREPARATION .....6

    Mixing in Water .....6

    Mixing with Other Herbicides .....6

SPRAY DRIFT MANAGEMENT .....6

SPRAY TANK CLEAN OUT .....7

STORAGE AND DISPOSAL .....9

NOTICE TO BUYER .....9

LIMITATION OF WARRANTY AND LIABILITY .....10



# DuPont™ Krovar® I DF

herbicide **ACCEPTED**  
with **COMMENTS**  
In EPA Letter Dated:  
**DEC - 8 2005**

Dispersible Granules Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 352-505

Active Ingredients	PBY Weight
Bromacil [5-bromo-3-sec-butyl-6-methyluracil]	40%
Diuron [3-(3,4-dichlorophenyl)-1,1-dimethylurea]	40%
Inert Ingredients	20%
<b>TOTAL</b>	<b>100%</b>

EPA Reg. No. 352-505

## KEEP OUT OF REACH OF CHILDREN CAUTION

### FIRST AID

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

For medical emergencies involving this product, call toll free 1-800-441-3637.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed or if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

### PERSONAL PROTECTIVE EQUIPMENT

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistant category selection chart.

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinylchloride
- Dust mist respirator
- Shoes plus socks

#### Mixers and loaders must also wear:

- Apron

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### ENGINEERING CONTROL STATEMENT

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

**Users should:** Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Bromacil is known to leach through soil and has been found in ground water as a result of normal field use. Users are advised not to apply in areas where soils are permeable, particularly where ground water is used for drinking water. Consult with the pesticide state lead agency for information regarding soil permeability and aquifer vulnerability in your area.

5 8 13

## GENERAL INFORMATION

DuPont™ KROVAR® I DF is a dispersible granule to be mixed in water and applied as a spray for selective control of weeds in citrus and for non-crop weed control.

KROVAR® I DF controls many annual weeds and, at the highest rates allowed by this label, it controls certain perennial weeds.

Moisture is necessary to move the herbicide into the root zone of weeds. Best results are obtained if treatment is made to moist soil, and moisture is supplied by rainfall or sprinkler irrigation within two weeks after application. Weed control symptoms are slow to appear and may not become apparent until the herbicide has been carried into the root zone of the weeds by moisture. The degree and duration of control will vary with the amount of herbicide applied, soil texture, rainfall, and other soil and water management practices.

## USE PRECAUTIONS AND RESTRICTIONS

KROVAR® I DF is not to be used in any recreational areas or around homes.

Injury to or loss of desirable trees or other plants may result from failure to observe the following:

Do not apply (except as recommended for crop use), or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts or similar areas. Do not use in home fruit plantings nor in citrus orchards interplanted to other trees or desirable plants. Prevent drift of dry powder or spray to desirable plants. Keep from contact with fertilizers, insecticides, fungicides and seeds.

Do not apply this product through any type of irrigation system.

Do not graze cattle in treated areas.

Treated areas may be planted to citrus trees one year after last application. Do not replant to other crops within two years after last application as injury may result.

### NOTE: Additional Precautions

Avoid storage of pesticides near well sites.

Calibrate sprayers only with clean water away from the well site.

Measure out only enough KROVAR® I DF for the job at hand.

Avoid over-filling the spray tank.

Do not discharge excess material as a point source.

Dilute and agitate excess spray solution and apply at labeled rates.

## CROP ROTATION BIOASSAY

In arid climates (10 inches of rainfall or less) or areas where drought conditions have prevailed for one or more years, a field bioassay should be completed prior to planting any desired crop(s). The results from this bioassay may require the two-year crop rotation interval to be extended. A successful field bioassay means growing to maturity a test strip of the crop(s) intended for production. The test strip should cross the entire field including knolls and low areas.

## RESISTANCE

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

## INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. KROVAR® I DF herbicide should only be used in accordance with recommendations on this label. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

## AGRICULTURAL USES

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Chemical Resistant Gloves made of any Waterproof material.
- Shoes plus socks.

## APPLICATION INFORMATION

Apply DuPont™ KROVAR® I DF with a properly calibrated fixed-boom power sprayer as a band or broadcast treatment. Apply any time of year provided overhead moisture (rainfall or sprinkler irrigation) is available to activate the herbicide, preferably just before or just after weeds have germinated.

All use rates of KROVAR® I DF are expressed for broadcast treatments. For band treatments, use proportionately less.

Use sufficient spray volume (minimum of 10 gallons per acre) to provide uniform coverage of the treated area and to allow proper dispersion and suspension of the product in the spray tank. Continuous agitation in the spray tank is required to keep the product in suspension. Agitate spray tank contents by mechanical or hydraulic means. If a by-pass or return line is used, it should terminate at a bottom of tank to minimize foaming. Do not use air agitation.

Best results are obtained if KROVAR® I DF is applied to bare ground. If weeds are present at application, tank mixtures with foliar active herbicides are recommended (see Tank Mixtures section of label). If dense populations of hard-to-kill weed species are present, control of these weeds prior to application of KROVAR® I DF is recommended.

### TANK MIXTURES

KROVAR® I DF may be tank mixed with other suitable herbicides registered for use in citrus. Refer to the tank mixture partner label(s) for any additional use information or restrictions. Follow the label guidelines that are the most restrictive.

KROVAR® I DF may also be tank mixed with appropriate adjuvants used with herbicides in citrus.

NOTE: If there is no prior use experience with the tank mixture combination, a compatibility test should be performed prior to adding the products into the spray tank. See SPRAY PREPARATION section of the label for further information.

When using KROVAR® I DF alone or in combination, thoroughly re-agitate the spray tank contents if allowed to settle.

## WEEDS CONTROLLED

### Annuals

- |                              |                                 |
|------------------------------|---------------------------------|
| Barnyardgrass                | <i>Echinochloa crus-galli</i>   |
| Brome, downy (cheatgrass)    | <i>Bromus tectorum</i>          |
| Chickweed, common            | <i>Stellaria media</i>          |
| Chickweed, mouseear          | <i>Cerastium vulgatum</i>       |
| Clovers (annual)             | <i>Trifolium spp.</i>           |
| Filaree                      | <i>Erodium spp.</i>             |
| Fleabane, flaxleaved (hairy) | <i>Conyza bonariensis</i>       |
| Foxtail                      | <i>Setaria spp.</i>             |
| Goatweed                     | <i>Scoparia dulcis</i>          |
| Groundsel                    | <i>Senecio spp.</i>             |
| Horseweed (marestail)        | <i>Conyza canadensis</i>        |
| Johnsongrass                 | <i>Sorghum halepense</i>        |
| Junglerice                   | <i>Echinochloa colona</i>       |
| Kochia                       | <i>Kochia scoparia</i>          |
| Lambsquarter                 | <i>Chenopodium album</i>        |
| Lettuce, wild                | <i>Lactuca serriola</i>         |
| Mustard, wild                | <i>Brassica kaber</i>           |
| Natalgrass (red top)         | <i>Rhynchelytrum repens</i>     |
| Nightshade (annual)          | <i>Solanum spp.</i>             |
| Pigweed                      | <i>Amaranthus spp.</i>          |
| Pineappleweed                | <i>Matricaria matricariodes</i> |
| Puncturevine, common         | <i>Tribulus terrestris</i>      |
| Purslane, common             | <i>Portulaca oleracea</i>       |
| Pusley, Florida              | <i>Richardia scraba</i>         |
| Ragweed, common              | <i>Ambrosia artemisifolia</i>   |
| Sandbur (sandspur)           | <i>Cenchrus spp.</i>            |
| Shepherdspurse               | <i>Capsella bursa-pastoris</i>  |
| Sowthistle, annual           | <i>Sonchus oleraceus</i>        |
| Spanishneedles               | <i>Bidens pilosa</i>            |
| Thistle, Russian             | <i>Salsola australis</i>        |

### Perennials (At maximum rates and repeat treatments)

- |                             |                            |
|-----------------------------|----------------------------|
| Balsamapple vine (seedling) | <i>Momordica charantia</i> |
| Bermudagrass                | <i>Cynodon dactylon</i>    |
| Drymary                     | <i>Drymaria spp.</i>       |
| Guineagrass                 | <i>Panicum maximum</i>     |
| Milkweed vine (strangler)   | <i>Morrenia odorata</i>    |
| Quackgrass                  | <i>Agropyron repens</i>    |
| Vines (seedlings)           |                            |

## CITRUS

Apply KROVAR® I DF as a band or broadcast treatment beneath and/or between trees.

Avoid contact of citrus foliage and fruit with spray or mist. Avoid overlapping and shut off spray boom while starting, turning, slowing or stopping as injury to trees may result.

Temporary yellowing of citrus leaves may occur following treatment. As injury to citrus trees may result, do not use on soils with less than 1% organic matter. Do not use on poorly drained soils, gravelly soils or thinly covered or exposed subsoils.

Do not treat trees planted in irrigation furrows. Do not treat diseased or stressed citrus trees.

Do not use in citrus groves inter-planted with other trees or desirable plants or in areas where roots of desirable plants or trees may extend, as injury may result. Do not use in home citrus plantings.

Do not graze cattle in treated areas.

When making multiple applications, do not apply at less than 60 day intervals.

**CALIFORNIA, ARIZONA**

**Trees Established for at least Three Years:** Best results occur when applied in late fall or early winter, but before winter annuals become well established. Application should be made after the first fall or early winter rains have settled the soil.

For the initial treatment, apply 4-5 pounds DuPont™ KROVAR® I DF per acre on coarse soils containing 1-2% organic matter and 5-6 pounds per acre on fine soils, or soils with organic matter of 2 1/2% or more and repeat as needed. Alternatively, apply 3-4 pounds per acre in the fall and repeat at 2-4 pounds per acre in the spring. Do not exceed 6 pounds per acre per year.

Use the highest rate allowed by this label where groundsel or puncturevine are known to be a problem. These rates will also suppress low density stands of bermudagrass and yellow nutsedge. Repeat annually for best treatment effect.

**FLORIDA**

For applications in the state of Florida refer to the additional instructions at the end of this label.

**LOUISIANA**

**Trees Established for at least Three Years:** Make a single application of 2 to 4 pounds per acre on coarser soils (sands, loamy sands, sandy loams) and 4 to 6 pounds per acre on finer soils (silt loams, clay loams, or soils with organic matter of 2 1/2% or more); use the highest rate allowed by this label for maximum suppression of perennials. Alternatively, make two applications per year at rates of 2 pounds per acre on coarser soils and 3 pounds per acre on finer soils; make the second application when needed to maintain weed control. Do not apply more than 6 pounds per acre per year.

**TEXAS**

**Trees Established Less than One Year:** Apply 2-4 pounds KROVAR® I DF per acre as needed to maintain weed control. Do not apply at less than 60-day intervals. Do not apply more than 6 pounds per acre per year.

**Trees Established One or Two Years:** Apply 2-4 pounds KROVAR® I DF per acre. A second application may be made when needed to maintain weed control, but do not exceed 6 pounds per acre per year.

**Trees Established Three or More Years:** Make one to two applications per year as needed to maintain weed control. Use 2-4 pounds per acre on coarser soils (sands, loamy sands, sandy loams) and 4-6 pounds per acre on finer soils (silt loams, clay loams, or soils with organic matter of 2 1/2 % or more). Use the higher rate for maximum suppression of perennials. Do not use more than 6 pounds per acre per year.

**SPRAY PREPARATION**

**Mixing in water** - Fill tank 1/2 full with water. Start agitation system, add KROVAR® I DF and continue adding water. Add separately each additional component of any tank-mix while adding water. Continue agitation throughout.

**Mixing in liquid fertilizer** - A fertilizer solution may be used in the spray mixture. Small quantities should be tested

for compatibility by the following procedures before full scale mixing.

1. Put 1 pint fertilizer solution in a quart jar.
2. Mix 2 teaspoonfuls KROVAR® I DF with 2 tablespoons of water; mix thoroughly and add to fertilizer solution.
3. Close jar and shake well.
4. If other herbicides are used in the mixture, premix 2 teaspoons of dry materials or 1 teaspoon of liquids with 2 tablespoons of water; add to KROVAR® I DF-fertilizer solution mixture.
5. Close jar and shake well.
6. Watch mixture for several seconds; check again in 30 minutes.
7. If mixture does not separate, foam, gel or become lumpy, it may be used.

Provided the above procedure shows the mixture to be compatible, prepare the tank mixture as follows: Add the fertilizer solution to the spray tank first; with agitator running, add the required amount of KROVAR® I DF and thoroughly mix.

**Mixing with other herbicides** - Determine the tank mixture partner(s) compatibility with KROVAR® I DF by following the directions above. For Step 1 above, use 1 pint of water instead of the liquid fertilizer. Provided the above procedure shows the mixture to be compatible, KROVAR® I DF may be used in this tank mixture.

**NON-AGRICULTURAL USES**

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Noncrop weed control is not within the scope of the Worker Protection Standard.

Do not enter or allow others to enter the treated area until sprays have dried.

**USE RESTRICTIONS - STATE OF FLORIDA**

In the state of Florida the use of KROVAR® I DF (bromacil + diuron) is prohibited in the counties of Hardee, Highland, Polk, Orange and Lake. For Non-Agricultural Usage in all other areas of the state, do not apply more than 16 pounds per acre per year of KROVAR® I DF. This amount corresponds to 6.4 pounds of bromacil and 6.4 pounds of diuron, the active ingredients in KROVAR® I DF. The maximum allowable use rate for bromacil is 6.4 pounds per acre per year inclusive of all bromacil formulations.

**APPLICATION INFORMATION**

KROVAR® I DF is recommended for general weed control as follows: uncultivated non-agricultural areas (such as, airports, highway, railroad and utility rights-of-way, sewage disposal areas); uncultivated agricultural areas (non-crop producing, which includes: farmyards, fuel storage areas,

fence rows, barrier strips); industrial sites (outdoor, such as, lumberyards, pipeline and tank farms).

Apply DuPont™ KROVAR® I DF using a properly calibrated fixed-boom power sprayer. Use sufficient spray volume (minimum of 10 gallons per acre) to provide uniform coverage of the treated area and to allow proper dispersion and suspension of the product in the spray tank. All rates of KROVAR® I DF are expressed for broadcast treatments. For band treatments, use proportionately less.

Combination with other herbicides broadens the spectrum of weeds controlled. In addition, total vegetation control can be achieved with higher rates of KROVAR® I DF plus residual-type companion herbicides. To improve the control of emerged weeds, add surfactant at 0.25% by volume.

**Note:** Applications may also be made using a handgun sprayer. Use a spray volume of at least 40 gallons per acre to insure uniform coverage. For small areas, a hand sprayer or sprinkling may be used.

**NON-CROP WEED CONTROL**

**APPLICATION TIMING**

Apply KROVAR® I DF as a preemergence spray prior to or during the rainy season when weeds are actively germinating or growing. Moisture is required to activate and move KROVAR® I DF into the root zone of weeds for preemergence control. For best preemergence control, apply prior to rainfall and weed germination.

In arid regions of the Western U.S., to insure adequate moisture for activation and even dispersion of the herbicide in the soil profile, KROVAR® I DF should be applied several weeks prior to the Fall freeze or shortly after Spring thaw to coincide with periods of higher seasonal moisture probability. Do not treat frozen or saturated soils, or soils that are non-receptive to percolation.

Do not apply to sites which have roots of desirable plants growing into the treatment zone as plant injury or death may occur. Do not apply to hard or impervious soils, water saturated soils or to any surface that does not allow the herbicide to be moved into the soil horizon with moisture. Unusually heavy rainfall shortly after application may move the product off-target to the lowest surrounding point and cause plant injury or death.

If herbicide treated soil is disturbed by any physical or mechanical means, the herbicide barrier is disrupted and the likelihood of non-performance may increase. For best performance results, make sure the treatment area is stable after the application for the desired weed control period.

**APPLICATION RATES**

Apply KROVAR® I DF at the rates indicated by weed type. When applied at lower rates, KROVAR® I DF provides short-term control of weeds listed; when applied at higher rates, weed control is extended.

**WEEDS CONTROLLED**

KROVAR® I DF effectively controls the following broadleaf weeds and grasses when applied at the rates shown.

**Broadleaf Weeds--6-8 pounds per acre**

Clovers (annual)	<i>Trifolium spp.</i>
Fiddleneck	<i>Amsinckia intermedia</i>
Filaree	<i>Erodium spp.</i>
Knapweed, diffuse	<i>Centaurea diffusa</i>
Lambsquarter, common	<i>Chenopodium album</i>
Lettuce, prickly	<i>Lactuca serriola</i>
Mustards	<i>Brassica spp.</i>
Pigweed	<i>Amaranthus spp.</i>
Ragweed	<i>Ambrosia spp.</i>
Sunflower, common	<i>Helianthus annuus</i>
Thistle, Russian	<i>Salsola iberica</i>

**Broadleaf Weeds--8-12 pounds per acre**

Carrot, wild	<i>Daucus carota</i>
Dandelion, common	<i>Taraxacum officinale</i>
Dock, curly	<i>Rumex crispus</i>
Knapweed, spotted	<i>Centaurea maculosa</i>
Knotweed, prostrate	<i>Polygonum aviculare</i>
Kochia	<i>Kochia scoparia</i>
Marestail, common (horseweed)	<i>Conyza canadensis</i>
Parsnip, wild	<i>Pastinaca sativa</i>
Plantain	<i>Plantago spp.</i>
Puncturevine	<i>Tribulus terrestris</i>
Spurge	<i>Euphorbia spp.</i>
Thistle, milk	<i>Silybum marianum</i>
Yarrow, common	<i>Achillea millefolium</i>

**Broadleaf Weeds--12-16 pounds per acre**

Cinquefoil, common	<i>Potentilla canadensis</i>
Goldenrod	<i>Solidago spp.</i>
Milkweed, common	<i>Asclepias syriaca</i>

**Grasses--6-8 pounds per acre**

Barley, foxtail	<i>Hordeum jubatum</i>
Brome	<i>Bromus spp.</i>
Cheat	<i>Bromus secalinus</i>
Cupgrass, Prairie	<i>Eriochloa contracta</i>
Foxtail	<i>Setaria spp.</i>
Oat, wild	<i>Avena fatua</i>
Ryegrass, Italian	<i>Lolium multiflorum</i>
Quackgrass	<i>Agropyron repens</i>
Wheatgrass, intermediate	<i>Agropyron intermedium</i>

**Grasses--8-12 pounds per acre**

Bahiagrass	<i>Paspalum notatum</i>
Crabgrass	<i>Digitaria spp.</i>
Goosegrass	<i>Eleusine indica</i>
Rye	<i>Secale cereale</i>
Vaseygrass	<i>Paspalum urvillei</i>

**Grasses--12-16 pounds per acre**

Bluegrass	<i>Poa spp.</i>
Dropseed, sand *	<i>Sporobolus cryptandrus</i>
Fescue	<i>Festuca spp.</i>
Saltgrass*	<i>Distichlis spp.</i>

\*Note: Best control of Saltgrass and Sand Dropseed is achieved from a Spring application prior to plant green-up.

For control of hard-to-kill perennials such as bermudagrass (*Cynodon dactylon*), bouncingbet (*Saporaria officinalis*), dogbane (*Apocynum spp.*), Johnsongrass (*Sorghum halepense*), and nutsedge (*Cyperus spp.*) apply 19 - 30 pounds per acre (except Florida).

For extended control of annual weeds and partial control of perennials such as bermudagrass and nutsedge, apply 10-18 pounds\* per acre. Use the higher KROVAR® I DF rates on adsorptive soils (high in organic matter or carbon). Best results occur when application is made just before weed emergence or in the early stages of weed growth.

**Retreating:** Apply 4 to 6 pounds per acre when annual weeds and grasses reappear on sites where weed growth has been controlled.

**Small Areas:** 1/4 cupful of DuPont™ KROVAR® I DF per 200 sq. ft. is approximately 15 pounds per acre.

### TANK MIXTURES

KROVAR® I DF may be tank mixed with other suitable herbicides registered for non-agricultural use. Refer to the tank mixture partner label(s) for any additional use information or restrictions. Follow the label guidelines that are the most restrictive.

KROVAR® I DF may also be tank mixed with appropriate adjuvants used with herbicides for non-agricultural use.

**NOTE:** If there is no prior use experience with the tank mixture combination, a compatibility test should be performed prior to adding the products into the spray tank. See **SPRAY PREPARATION** section of the label for further information.

When using KROVAR® I DF alone or in combination, thoroughly re-agitate the spray tank contents if allowed to settle.

### SPECIAL USES

#### UNDER ASPHALT AND CONCRETE PAVEMENT

##### APPLICATION INFORMATION

KROVAR® I DF can be used to control weeds under asphalt and concrete pavement, such as that used in parking lots, highway shoulders, median strips, roadways, and other industrial sites.

KROVAR® I DF should only be used in an area that has been prepared according to good construction practices. Use sufficient water to insure uniform coverage, generally 100 gal per acre. Agitate the tank continuously to keep KROVAR® I DF in suspension.

##### APPLICATION TIMING

KROVAR® I DF should be applied immediately before paving to avoid lateral movement of the herbicide as a result of soil movement due to rainfall or mechanical means.

##### APPLICATION RATES

Apply KROVAR® I DF at 17 to 30 pounds per acre. Use a higher rate for hard to control weeds and/or for longer term weed control.

### TANK MIXTURES

To control a broader spectrum of weeds, or for an extended period of weed control, a tank mixture of KROVAR® I DF at 7 to 15 pounds per acre plus DuPont™ OUST® XP at 4 to 8 ounces per acre may be used.

#### Important Precautions-Under Asphalt Only

- Do not use KROVAR® I DF under pavement in residential properties such as driveways, or in recreational areas, including jogging or bike paths, tennis courts, or golf cart paths.
- Desirable plants may be injured if their roots extend into treated areas or if planted in treated areas.

### SPRAY PREPARATION

**Mixing in water** - Fill tank 1/2 full with water. Start agitation system, add KROVAR® I DF and continue adding water. Add separately each additional component of any tank-mix while adding water. Continue agitation throughout.

**Mixing with other herbicides** - Determine the tank mixture partner(s) compatibility with KROVAR® I DF by following the directions below. Provided the procedure below shows the mixture to be compatible, KROVAR® I DF may be used in this tank mixture.

1. Put 1 pint of water in a quart jar.
2. Add 2 teaspoons of KROVAR® I DF and mix thoroughly.
3. For other herbicides used in the mixture, premix 2 teaspoons of dry materials or 1 teaspoon of liquids with 2 tablespoons of water; add to KROVAR® I DF mixture.
4. Close jar and shake well.
5. Watch mixture for several seconds; check again in 30 minutes.
6. If the mixture does not separate, foam excessively, gel or become lumpy, it may be used.

### SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

**AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.**

#### IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150 - 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **Wind, Temperature and Humidity, and Surface Temperature Inversions** sections of this label.

#### CONTROLLING DROPLET SIZE - GENERAL TECHNIQUES

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

#### CONTROLLING DROPLET SIZE - AIRCRAFT

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is

emitted backwards, parallel to the airstream will produce larger droplets than other orientations.

- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.

**BOOM LENGTH AND HEIGHT**

- **Boom Length (aircraft)** - The boom length should not exceed 3/4 of the wing length, using shorter booms decreases drift potential. For helicopters use a boom length and position that prevents droplets from entering the rotor vortices.
- **Boom Height (aircraft)** - Application more than 10 ft above the canopy increases the potential for spray drift.
- **Boom Height (ground)** Setting the boom at the lowest height which provides uniform coverage reduces the exposure of droplets to evaporation and wind. The boom should remain level with the crop and have minimal bounce.

**WIND**

Drift potential increases at wind speeds of less than 3 mph (due to variable direction and inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. AVOID APPLICATIONS DURING GUSTY OR WINDLESS CONDITIONS.

**NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they effect spray drift.

**TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

**SURFACE TEMPERATURE INVERSIONS**

Drift potential is high during a surface temperature inversion. Surface inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Surface inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates a surface inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

**SPRAY TANK CLEAN OUT**

Thoroughly clean all traces of DuPont™ KROVAR® I DF from application equipment immediately after use. Flush the tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts

separately). Dispose of the equipment wash water by applying it to a use-site listed on this label.

**FOR USE IN CONTROLLING WEEDS IN CITRUS IN THE STATE OF FLORIDA ONLY**

KROVAR® I DF controls many annual weeds, and at the highest rates allowed by this label, it controls certain perennial weeds.

Best results are obtained if KROVAR® I DF is applied to bare ground. If dense populations of hard-to-kill weed species are present, control of these weeds prior to application of KROVAR® I DF is recommended. If weeds are present at application, tank mixtures with foliar active herbicides is recommended to control existing weeds. (See Tank Mixtures section of Label). Apply any time of year provided overhead moisture (rainfall or sprinkler irrigation) is available to activate the herbicide, preferably just before or just after weeds have germinated. Weed control symptoms are slow to appear and may not become apparent until the chemical has been carried into the root zone of the weeds by moisture. Best results are obtained if moisture is supplied by rainfall or sprinkler irrigation within two weeks after application. The degree and duration of control will vary with the amount of herbicide applied, soil texture, rainfall, and other soil and water management practices.

**GENERAL PRECAUTIONS AND USE RESTRICTIONS**

The use of KROVAR® I DF (bromacil + diuron) is prohibited for weed control in non-bedded citrus groves located on any permeable, better drained soil identified in the intended site of application. Permeable, better drained soils which occur in citrus producing areas of the state include soils unnamed and characteristic of quartzipsamments, and the following soil series classifications:

Adamsville	Dade	Orsino
Archbold	Florahome	Palm Beach
Astatula	Fort Meade	Paola
Bahiahonda	Gainesville	Satellite
Broward	Lake	St. Augustine
Canaveral	Lakewood	St. Lucie
Candler	Neilhurst	Tavares
Cocoa	Orlando	

Do not use DuPont™ KROVAR® I DF in recreational areas or around homes. Prevent drift of dry powder or spray to desirable plants.

Do not use on lawns, walks, driveways, tennis courts, or similar areas. Do not apply directly to water, or to areas where surface water is present, or to inter-tidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash water. Keep from contact with fertilizer, insecticides, fungicides, and seeds. Thoroughly clean all traces of KROVAR® I DF from application equipment immediately after use. Flush tank, pump, hoses and boom with several changes of water after removing nozzle tips and screens (clean these parts separately.)

Temporary yellowing of citrus leaves may occur following treatment. Because injury to citrus trees may result, do not use on soils with less than 1.0% organic matter, or on poorly drained soils.

Avoid contact of foliage or fruit with spray or mist. Avoid

overlapping treated areas and shut off spray boom while starting, turning, slowing or stopping, or injury to trees may result.

Injury to or loss of desirable trees or other plants may result from failure to observe the following:

Do not apply (except as recommended for crop use), drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.

Do not treat trees planted in irrigation furrows.

Do not treat diseased or stressed trees.

Do not use in citrus groves inter-planted with other desirable trees or plants, or in areas where roots of desirable trees or plants may extend, as injury to desirable trees or plants may result.

Do not use in home citrus plantings.

Do not apply this product through any type of irrigation system. Do not apply this product in a way that will contact workers or other persons, either directly or through drift.

Only protected handlers may be in the area during application.

Citrus trees may be planted one year after the last application of KROVAR® I DF.

Do not replant areas to any crop within two years after the last application of KROVAR® I DF as injury to subsequent crop(s) may result.

Allow an interval of 60 days between applications of KROVAR® I DF.

Do not graze cattle in treated areas.

In areas where drought conditions have prevailed for one or more years, a field bioassay should be completed prior to planting any desired crop(s). The results from this bioassay may require the two year crop rotation interval to be extended. A successful field bioassay means growing to maturity a test strip of the crop(s) intended for production. The test strip should cross the entire field including knolls and low areas.

**APPLICATION INSTRUCTIONS**

Apply KROVAR® I DF as a band treatment only using a properly calibrated fixed-boom power sprayer. Do not use Trunk to Trunk. All use rates of KROVAR® I DF are expressed for broadcast treatments. For band treatments, use proportionately less.

Use sufficient spray volume (minimum of 10 gallons per acre) to provide uniform coverage of the treated area and to allow proper dispersion and suspension of the product in the spray tank. Continuous agitation in the spray tank is required to keep the product in suspension.

Do not apply more than 16 pounds of KROVAR® I DF per treated acre per year. This amount corresponds to 6.4 pounds of bromacil and 6.4 pounds of diuron, the active ingredients in KROVAR® I DF.

The maximum allowable use rate for bromacil is 6.4 pounds per treated acre per year inclusive of all bromacil formulations. The maximum allowable used rate for diuron is 9.6 pounds per treated acre per year inclusive of all diuron formulations.

Multiple applications may improve control of "hard-to-kill" weed species.

**Trees Established Less Than One Year:** For control of annual weeds, apply 2-4 pounds of KROVAR® I DF per treated acre as needed to maintain weed control. Do not apply more than 6 pounds per treated acre during any 6 month period nor more than 8 pounds per treated acre during the first year.

**Trees Established One to Three Years:** For control of annual weeds, apply 2-4 pounds of KROVAR® I DF per treated acre. A second application may be made when needed to maintain weed control, but do not exceed 8 pounds per treated acre per year.

**Trees Established Three or More Years:** Apply 4-8 pounds per treated acre as needed to maintain weed control. Do not apply more than 16 pounds of KROVAR® I DF per treated acre per year.

**Tank Mixtures:** KROVAR® I DF may be tank mixed with other suitable herbicides registered for use in citrus. Refer to the tank mixture partner label(s) for any additional use information or restrictions. Follow the label guidelines that are the most restrictive.

**NOTE:** If there is no prior use experience with the tank mixture combination, a compatibility test should be performed prior to adding the products into the spray tank. See **SPRAY PREPARATION** section of the label for further information.

When using KROVAR® I DF alone or in combination, thoroughly re-agitate the spray tank contents if allowed to settle.

**WEEDS CONTROLLED (Annuals)**

- |                         |                      |
|-------------------------|----------------------|
| Annual clovers          | Lambsquarter         |
| Barnyardgrass           | Natalgrass (red top) |
| Chickweed               | Nightshade (annual)  |
| Cheatgrass              | Pigweed              |
| Crabgrass               | Pineappleweed        |
| Filaree                 | Puncturevine         |
| Fleabane                | Purslane             |
| Florida pusley          | Ragweed              |
| Foxtail                 | Russian thistle      |
| Goatweed                | Sandbur (sandspur)   |
| Groundsel               | Shepherdspurse       |
| Horseweed               | Sowthistle (annual)  |
| Johnsongrass (seedling) | Spanish needles      |
| Junglerice              | Wild lettuce         |
| Kochia                  | Wild mustard         |

**Perennial Weeds**

- Balsamapple vine (seedling)
- Bermudagrass
- Heartleaf
- Drymary
- Guineagrass
- Milkweed (strangler)
- Quackgrass
- Vine (seedling)

**NOTE:** Use the highest rates allowed by this label for best control of perennial weeds listed on this label. Partial control of perennial weeds can result with only a single treatment of KROVAR® I DF. Repeat applications are required (in season and/or annually) for best control of the perennial weeds on this label. Control of perennials may be improved by cultivation prior to treatment, otherwise, avoid working the soil as long as weed control continues or else effectiveness of the treatment may be reduced.

12 7 13

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store product in original container only. Store in a cool, dry place.

**Pesticide Disposal:** Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal: For Plastic Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **For Fiber Sacks:** Completely empty fiber sack by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into manufacturing or application equipment. Then dispose of sack in a sanitary landfill or by incineration if allowed by State and local authorities. **For Fiber Drums With Liners:** Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner. **For Paper and Plastic Bags:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Container Refilling and Disposal (For Containers up to 250 gal):** This is a refillable container. If the container is to be refilled, do not rinse with any material or introduce any pesticide other than DuPont™ KROVAR® I DF. Reseal and return the container to any authorized DuPont refilling facility. If the container is not to be refilled, triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-441-3637 day or night.

**Container Disposal for Bulk Containers:** When this container is empty, replace the cap and seal all openings that have been opened during use, and return the container to the point of purchase or to a designated location named at time of purchase of this product. The container must only be refilled with this pesticide product. **DO NOT REUSE THE CONTAINER FOR ANY OTHER PURPOSE.** Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transporting. Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, contact DuPont at 1-800-441-3637. If not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling. Disposal of this container must be in compliance with state and local regulations.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-441-3637 day or night.

**NOTICE TO BUYER:** Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval, DuPont™, KROVAR® and OUST® are trademarks or registered trademarks of E. I. duPont de Nemours and Company

D - 1063 092905

### **LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read This Limitation Of Warranty And Liability Before Buying Or Using This Product. If The Terms Are Not Acceptable, Return The Product At Once, Unopened, And The Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product, crop injury, or injury to non-target crops or plants. DuPont does not agree to be an insurer of these risks. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

**DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE FULLEST EXTENT PERMITTED BY LAW, IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict Liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

**For product information call: 1-888-6-DUPONT**

**Internet address: [www.dupont.com/ag/us](http://www.dupont.com/ag/us)**

**© 1999-2005 E. I. du Pont de Nemours and Company, Wilmington, Delaware 19898.**

**All rights reserved.**