

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division

OCT 22 1993

Ian Wellings
E. I. DU PONT DE NEMOURS AND CO, INC.
BARLEY MILL PLAZA, WALKER'S MILL
WILMINGTON DE 19880

Subject: Label Amendment Submission of 06/24/93
to Comply with WPS Labeling Requirements
EPA Reg Nr. 352-508
DUPONT KARMEX DF HERBICIDE

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is accepted subject to the comments listed below.

Based on your certification, the Agency has accepted only those changes to your labeling which are necessary to comply with PR Notices 93-7 and 93-11, which reflect the WPS labeling requirements of 40 CFR part 156, subpart K. Any other labeling changes submitted in connection with this amendment application and not directly related to compliance with PR Notice 93-7 or 93-11 have neither been reviewed nor accepted by the Agency. If you wish to make any such changes, you must submit a separate amendment application proposing them. If your product registration is currently suspended, acceptance of this labeling amendment does not affect the suspension in any way.

A copy of your proposed labeling stamped, "Accepted with comments", is attached. Make any required changes described below and send three copies of final labeling as soon as it is available to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (H-7504C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, DC 20460-0001

BEST AVAILABLE COPY

2 of 13

E. I. DU PONT DE NEMOURS AND CO, INC.
EPA Reg. Number: 352-508
DUPONT KARMEX DF HERBICIDE
Application date: 06/24/93
Original Submission


Hand or courier deliveries of final labels may be made to:

Document Processing Desk (FIN-LAB:L)
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

The Nonagricultural Use Requirements section is not required as no special requirements are identified.

Please correct the typographical errors circled on the draft before printing final labeling.

In your final labeling, the "Agricultural Use Requirements" text must be contained in a clearly separate box. This box may be set apart by a line, by another graphical device, by a different color background, or in any other way that clearly distinguishes it from surrounding text.

Sincerely,

Registration Support Team
Registration Division

BEST AVAILABLE COPY

3 of 13

ACCEPTED
with COMMENTS
in EPA Letter Dated

OCT 22 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

352-508



KARMEX® DF

HERBICIDE

DISPERSIBLE GRANULES

ACTIVE INGREDIENT

Diuron [3-(3,4-dichlorophenyl)-1,1-dimethylurea] 80%

INERT INGREDIENTS 20%

TOTAL 100%

EPA Reg. No. 352-508

KEEP OUT OF REACH OF CHILDREN

WARNING

AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

If on skin: Wash with plenty of soap and water; get medical attention if irritation persists.

If in eyes: Flush with plenty of water; get medical attention if irritation persists.

For medical emergencies involving this product, call toll free 1-800-441-3637.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING! CAUSES EYE IRRITATION. MAY IRRITATE NOSE, THROAT, AND SKIN.

Avoid contact with spray mist. Avoid contact with skin, eyes, and clothing.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

Long-sleeve shirt and long pants.

Waterproof gloves.

Protective eyewear.

Shoes plus socks.

Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Cover or incorporate spills.

IMPORTANT

Injury to or loss of desirable trees or other plants may result from failure to observe the following:

Do not apply (except as recommended for crop use), or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on home plantings of trees, shrubs or herbaceous plants, nor on lawns, walks, driveways, tennis courts or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water. Do not mix/load, or use near wells including abandoned wells, drainage wells, and sink holes. Avoid storage of pesticides near well sites. Keep from contact with fertilizers, insecticides, fungicides and seeds. Calibrate sprayers only with clean water away from the well site. Do not apply this product through any type of irrigation system.

Thoroughly clean all traces of "Karmex" DF from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

GENERAL INFORMATION

Du Pont "Karmex" DF Herbicide is a dispersible granule to be mixed in water and applied as a spray for selective control of weeds in certain crops and for weed control on non-cropland areas. It is non-corrosive to equipment, non-flammable and non-volatile.

"Karmex" DF may be applied to soil prior to emergence of weeds to control susceptible weed seedlings for an extended period of time; the degree of control and duration of effect will vary with the amount of chemical applied, soil texture, rainfall and other conditions. Soils high in clay or organic matter require higher dosages than soil low in clay or organic matter, for equivalent herbicide performance. Moisture is required to activate the herbicide; best results occur if rainfall (or sprinkler irrigation) occurs within 2 weeks of application.

"Karmex" DF applied before emergence of crop and weeds, is an effective procedure because susceptible weeds are controlled in an early, vulnerable seedling stage before they compete with the crop. With favorable moisture conditions, "Karmex" DF continues to control weeds for some time as the crop becomes better able to compete. Should weed seedlings begin to break through the preemergence treatment in significant numbers, secondary weed control procedures should be implemented; these include cultivation and postemergence herbicide application.

"Karmex" DF may also be used to control emerged weeds. Results vary with the amount applied and environmental conditions; best results are obtained on succulent weeds growing under conditions of high humidity and temperatures of 70 degrees F or higher. Addition of a surfactant to the spray (where recommended) increases contact effects of "Karmex" DF.

"Karmex" DF may be used as a directed postemergence application. Contact of crop foliage and/or fruit with spray or mist must be avoided on the following crops: artichoke, corn (field), cotton, sorghum (grain), sugar cane and established plantings of apples, bananas, plantains, blueberries, caneberries, gooseberries, citrus, grapes, macadamia nuts, olives, papayas, peaches, pears, pecans, walnuts and certain tree plantings.

Under specified conditions (see Directions for Use), "Karmex" DF without surfactant may be applied over the top of alfalfa (established, dormant or semidormant), asparagus (established), birdsfoot trefoil (established, dormant), grass seed crops (established), oats, red clover (established, dormant), sugar cane, wheat, and pineapple.

Weed species vary in susceptibility to "Karmex" DF and they may be more difficult to control when under stress. Combinations of "Karmex" DF with other herbicides (as registered) increase the number of weed species controlled; consult labels of the companion product for this and other information. Observe all cautions on labeling of all products used in mixtures.

Since the effect of "Karmex" DF varies with soils, uniformity of application, and environmental conditions, it is suggested that growers limit their first use to small areas.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Waterproof gloves.
- Protective eyewear.
- Shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Non-crop weed control is not within the scope of the Worker Protection Standard.

Du Pont "Karmex" DF Herbicide should be used only in accordance with recommendations on this label, or in separate published Du Pont recommendations.

Du Pont will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by Du Pont. User assumes all risk associated with such non-recommended use.

BEST AVAILABLE COPY

SELECTIVE USE IN CROPS

PREEMERGENCE USE (Germinating Weeds)

"Karmex" DF, at recommended rates, controls annual weeds such as:

Broadleaves Controlled

<u>3/4 to 1 Lb/Acre</u>	<u>1 1/2 to 2 Lbs/Acre</u>	<u>2 to 6 Lbs/Acre</u>
Lambsquarters	Chickweed	Ageratum
Pigweed	Corn spurry	Corn speedwell
Purslane	Dogfennel	Dayflower
Ragweed	Fiddleneck	Flora's
	(amsinckia)	paintbrush
	Gromwell	Hawksbeard
	Groundcherry, annual	Horseweed
	Knawel	Kochia
	Morningglory, annual	Marigold
	Pennycress	Mexican clover
	Shepherd's- purse	Pineappleweed
	Tansymustard	Pokeweed
	Wild buckwheat	Rabbit tobacco
	Wild lettuce	Smartweed, annual
	Wild mustard	Sowthistle, annual
		Spanishneedles
		Velvetleaf (buttonweed)
		Wild radish

Partial control

<u>1 Lb/Acre</u>	<u>4 Lbs/Acre</u>
Cocklebur	Horsenettle
Morningglory, annual	
Prickly sida (teaweed)	
Sesbania	
Sicklepod	

Grasses Controlled

<u>3/4 to 1 Lb/Acre</u>	<u>1 1/2 to 2 Lb/Acre</u>	<u>2 to 6 Lb/Acre</u>
Barnyardgrass (watergrass)	Bluegrass, annual	Lovegrass, annual
Crabgrass	Foxtail	Ryegrass, annual
	Rattail fescue	Kyllinga
	Red sprangletop	Orchardgrass
	Velvetgrass	Peppergrass
	Vernalgrass, sweet, annual	Ricegrass
		Sandbur
		Johnsongrass, seedling

Partial control

<u>4 Lb/Acre</u>	<u>8 to 10 Lb/Acre</u>
Quackgrass	Guineagrass
	Maidencane
	Pangolagrass

POSTEMERGENCE USE

"Karmex" DF at recommended rates, controls seedling annual weeds such as morningglory, barnyardgrass (watergrass), crabgrass, crowfoot, goosegrass, pigweed and purslane. Addition of a surfactant to the spray (where recommended) increases contact effects of "Karmex" DF. Best results are obtained on succulent weeds growing under conditions of high humidity and temperatures of 70 degrees F or higher.

EQUIPMENT

Use a boom power sprayer properly calibrated to a constant speed and rate of delivery. Openings in screens should be 50 mesh or larger. Continuous agitation in the spray tank is required to keep the material in suspension. Agitate by mechanical or hydraulic means; if by-pass or return line is used, it should terminate at bottom of tank. Avoid overlapping, and shut off spray booms while starting, turning, slowing or stopping, or injury to the crop may result.

SPRAY VOLUME AND PRESSURES

For preemergence application, use 25 to 40 gals. per acre and spray pressure of 30 to 40 psi. For postemergence application, use sufficient volume (min. 25 gals. per acre) for thorough coverage of weed foliage; use spray pressure of 20 to 25 psi to keep spray drift to a minimum.

Aerial: For alfalfa, asparagus, barley (winter), cotton (preplant or preemergence only), grass seed crops, pineapple, sugar cane and wheat (winter), application may be made by aircraft at 5 to 10 gals. per acre. Avoid overlapping of spray swath and avoid application under conditions where excessive drift may occur. Where land is bedded, make application parallel to rows.

SPRAY PREPARATION

Mix proper amount of "Karmex" DF into necessary volume of water; where use of surfactant is recommended, dilute with 10 parts of water and add as last ingredient to nearly full tank.

USE RATES

All dosages of "Karmex" DF are expressed as broadcast rates; for band treatment, use proportionately less. For example, use 1/3 of the broadcast rate when treating a 14" band where row spacing is 42". Where a range of dosages is given, use the lower rate on coarse textured soils low in clay or organic matter and the higher rate on the fine textured soils high in clay or organic matter. For postemergence application, use the lower rate on smaller weeds and the higher rate on larger weeds.

BEST AVAILABLE COPY

6 of 13

SOIL LIMITATIONS

Crop injury may result from failure to observe the following:

Unless otherwise directed, do not use on sand, loamy sand, gravelly soils or exposed subsoils; nor on pecans where organic matter is less than 1/2%; nor on alfalfa, apples, artichoke, barley (winter), bermudagrass pasture, citrus, cotton, grapes, oats, olives, papayas, peaches, pears, sorghum, sugar cane, walnuts and wheat (winter) where organic matter is less than 1%; nor on blueberries, birdsfoot trefoil, caneberries, gooseberries, macadamia nuts and peppermint where organic matter is less than 2%.

Preemergence weed control will be reduced on high organic matter soils such as peat or muck.

REPLANTING

Unless otherwise directed, do not replant treated areas to any crop within 2 years after last application as injury may result.

FIELD CROPS

A good seedbed must be prepared before preemergence use of "Karmex" DF, as crop injury may result if application is made to ground which is cloddy or compacted resulting in improperly planted seed. Plant seed to depth specified.

Unless otherwise directed, the surface of the soil should not be cultivated or disturbed after application of "Karmex" DF and before emergence of the crop as weed control may be reduced and crop injury may result. However, if moisture is insufficient to activate the herbicide, a shallow cultivation or rotary hoe preferred should be made after emergence of crops while weeds are small enough to be controlled by mechanical means. See "Soil Limitations" for crops listed below.

ALFALFA

Treat only stands established for 1 year or more. Do not apply to seedling alfalfa nor to alfalfa/grass mixtures; do not apply to alfalfa under stress from disease, insect damage, shallow root penetration (such as on shallow hard pans), alkali spots; nor to flooded fields as crop injury may result. Do not spray on snow-covered or frozen ground.

Idaho, Oregon, Washington: Use 1 1/2 to 3 lbs per acre. For control of volunteer alfalfa, use 4 lbs per acre. Apply in fall after alfalfa becomes dormant but no later than mid-December.

California (Dormant and Semi-Dormant Varieties): Use 1 1/2 to 3 lbs per acre. For control of volunteer alfalfa, use 4 lbs per acre. Apply in fall or winter after alfalfa becomes dormant or semi-dormant, but before growth begins in the spring. Crop injury may result if application is made to actively growing alfalfa. For best results, apply before weeds have emerged or become established (2" in height or diameter). Control of established weeds is improved by applying "Karmex" DF with a suitable contact herbicide registered for such use. Sufficient rainfall for soil activation of "Karmex" DF is unlikely in California after February 1. Treated areas may be replanted to any crop after one year from last application if rate does not exceed 2 lbs per acre.

Arizona, Nevada: Use 1 1/2 to 3 lbs per acre. Apply in fall after alfalfa becomes dormant but no later than January.

Eastern Colorado, Kansas: For control of tansymustard, apply 1 lb per acre shortly after emergence of mustard in the fall or winter. Use 2 lbs per acre if weeds are 2" to 4" in height. Alternatively, if other annual weeds are present, apply 2 to 3 lbs per acre in February or March.

Other Areas Where Alfalfa Becomes Winter Dormant: Use 1 1/2 to 3 lbs per acre (1 1/2 to 2 lbs per acre East of Appalachian Mountains). Apply in March or early April, but before spring growth begins.

ARTICHOKE

California: Apply 2 to 4 lbs per acre in late fall or early winter after the last cultivation. Apply before weeds germinate or to emerging seedlings. Direct spray to cover the area between the rows and at the base of artichoke plants, keeping contact with crop plants at a minimum.

ASPARAGUS

Apply as a band or broadcast treatment. Do not apply to young plants during the first growing season (except as noted below), nor to newly seeded asparagus, nor on plants with exposed roots as severe injury may result. Preemergence weed control will be reduced on soils with greater than 5% organic matter.

Established Plantings: On light sandy soils and other soils low in clay or organic matter, apply 1 to 2 lbs per acre. On soils high in clay or organic matter, use 2 to 4 lbs per acre. Two applications may be used; the first application should be made before weeds become established but no earlier than 4 weeks before spring emergence and no later than the early cutting period. If weeds are controlled into the cutting period by cultural practices, application may be delayed until immediately after the last cultivation. A second application may be made immediately following completion of harvest provided rainfall is expected. When two applications are used in one season, do not exceed 3 lbs per acre per application. In Washington (irrigated crop), apply a single treatment of 4 lbs per acre.

If treatment is delayed until late winter or early spring, incorporation of the chemical in the top 1" to 2" of soil may substitute for lack of rain to activate the herbicide.

California (San Joaquin Delta)

Newly Planted Crowns: Make a single application of 2 to 4 lbs per acre on soils high in clay or organic matter. Use the lower rate on clay loams and the higher rate on peat soils. Do not use on soils containing less than 2% organic matter. Soil must be settled by rainfall or irrigation prior to treatment. Do not treat crowns planted to a depth of less than 2".

BARLEY, Winter

Western Oregon and Western Washington: For drill planted only, make a single application of 1 1/2 to 2 lbs per acre as soon as possible after planting but before emergence of barley. Do not replant treated areas to any crop within 1 year after last application as injury to subsequent crops may result.

BERMUDAGRASS PASTURES

For newly sprigged only, apply 1 to 3 lbs after planting and before emergence of bermudagrass or weeds. Alternatively, for control of emerged annual weeds up to 4" in height, apply 1/2 to 1 lb per acre with surfactant. If bermudagrass has emerged at time of treatment, temporary burn may occur.

Plant sprigs (stolons) 2" deep in a well-prepared seedbed. Do not treat areas where sprigs are planted less than 2" deep as crop injury may result. Do not graze or feed foliage from treated areas to livestock within 70 days after application.

BIRDSFOOT TREFOIL (Lotus)

Western Oregon: Treat only stands established for at least 1 year. Do not apply to seedling trefoil as injury may result. Make a single application of 2 lbs per acre when trefoil is dormant (October 15 to December 15). Do not replant treated areas to any crop within 1 year after last application as injury may result.

CORN, Field

Postemergence - Make a single application of 3/4 lb per acre in combination with non-pressure nitrogen solution. If nitrogen solution is not used, apply 1 lb per acre with surfactant. Apply as a directed spray when corn is at least 20" high and weeds are no taller than 3".

DO NOT APPLY OVER TOP OF CORN. Do not replant to any crop within 1 year, except that cotton, corn and grain sorghum may be planted the spring following treatment.

Alabama, Arkansas, Mississippi and Tennessee:

Preemergence - Make a single application of 2/3 to 1 lb per acre as a broadcast or band treatment after planting but before corn emerges. Plant corn at least 1 1/2" deep. Do not replant treated areas to crops other than corn or cotton within 4 months following band treatment and 6 months following broadcast treatment as injury may result.

COTTON

During a single crop season, do not exceed the following amounts of "Karmex" DF per acre as injury to subsequent crops may result: 1 lb on loamy sand; 1 1/2 lbs on sandy loam; 2 lbs on clay loam; 2 3/4 lbs on clay. Injury may occur if "Karmex" DF is used in conjunction with soil-applied organic phosphate pesticides. Do not allow livestock to graze treated cotton plants.

Preplant

Arizona and California: Use "Karmex" DF alone, or apply as a separate operation following preplant broadcast treatment with 'Treflan' (incorporated according to directions on "Treflan" label). Apply "Karmex" DF as a broadcast spray after beds are formed, preirrigated, and final seedbeds prepared. Prior to planting, drag-off the tops of the beds and plant in moist soil not treated with "Karmex" DF. Treated soil is returned to the bed after planting when irrigation furrows are reformed after cotton has emerged. If more than two furrowing-out operations

are made prior to lay-by, or deep furrows are made early, weed control may be reduced in furrow bottoms. Use at the following rates:

"Karmex" DF Alone

Apply 1 to 2 1/2 lbs per acre.

"Karmex" DF Following "Treflan" (preplant)

Apply as follows:

Soil Texture	Rate Per Acre	
	"Treflan" Preplant	"Karmex" DF Preemergence
Sandy loam, loam, silt loam, silt	1 pt	2/3 to 1 lb
Sandy clay loam, clay loam, silty clay loam, sandy clay, clay	1 1/2 pts	1 to 1 1/4 lbs

Seedling disease may weaken plants and increase the possibility of injury from the use of "Treflan" followed by "Karmex" DF. These treatments should be used only in conjunction with a standard fungicide seed treatment plus a good supplemental soil fungicide program such as captan-PCNB mixture.

Preemergence

United States, except Arizona, California, and areas west of Interstate 35 or 35W in Texas and Oklahoma: Use "Karmex" DF alone or apply as a separate operation following preplant treatment with "Treflan". Apply "Karmex" DF after planting but before cotton emerges. Do not treat cotton in deep furrows as crop injury may result. Use only where cotton is planted on flat or raised seedbeds. Shallow incorporation (no deeper than 1/4") with a rotary hoe or similar equipment following planting usually improves results especially during dry weather. A wide press wheel should be used on the planter to provide a level seedbed for subsequent early season postemergence treatments. If moisture is insufficient to activate "Karmex" DF or if soil becomes crusted before crop emerges, a shallow rotary hoeing (no deeper than 1/4") should be made before weeds become established.

"Karmex" DF Alone:

Make a single application as a broadcast or band spray, using the following broadcast rates; for band treatment, use proportionately less.

Soil Texture*	Lbs Per Acre
Loamy sand	2/3
Sandy loam, loam, silt loam, silt	1
Sandy clay loam, clay loam, silty clay loam, sandy clay	1 1/4
Silty clay, clay	2

*Do not use on soils with less than 1% organic matter as crop injury may result.

"Karmex" DF following "Treflan" (preplant):

Apply "Treflan" prior to planting as a broadcast or band treatment; incorporate according to directions on "Treflan" label. As a separate operation, apply "Karmex" DF as a band treatment (14" to 20" wide) after planting but before cotton emerges. Use at the following broadcast rates; for band treatment, use proportionately less. See "Note" under preplant Table above.

Soil Texture*	Rate Per Acre	
	"Treflan" Preplant	"Karmex" DF Preemergence
Loamy sand	1/2 pt	2/3 lb
Sandy loam, loam, silt loam, silt	1 pt	1 lb
Sandy clay loam, clay loam, silty clay loam, sandy clay, silty clay, clay	1 1/2 pts	1 1/4 - 2 lbs

*Do not use on soils with less than 1% organic matter as injury may result.

Postemergence Applications

United States

Apply only as a directed spray to cover weed foliage; adjust nozzles to minimize contact of cotton leaves with spray or drift. Injury may result.

DO NOT SPRAY OVER TOP OF COTTON PLANTS.

Early Season

Apply when cotton is at least 6" tall and when weeds are actively growing and do not exceed 2" in height. Apply as a band treatment at following rates. Two applications may be made if needed.

Weeds	lbs "Karmex" DF Per Acre Plus Surfactant	
	Broadcast	Band(1/3 area)
Up to 2" Tall		
Annual grasses	1/2	1/6
Pigweed	1/4	1/12

For control of seedling perennial grasses such as johnsongrass and partial control of nutsedge or when weed growth is under drought stress or as high as 4", add 2 to 3 1/2 lbs disodium methylarsonate (DSMA; 63% anhydrous or equivalent) to above spray mixture. If DSMA is used, do not apply after first bloom.

Late Season (Lay-By)

Apply 1 to 1 1/2 lbs per acre (1 to 2 lbs in Arizona and California) when cotton is at least 12" tall (at least 20" tall for Pima S-2). For control of germinating weed seedlings, apply to soil beneath cotton plants and between rows immediately after last cultivation. In irrigated cotton, best weed control is obtained if the field is irrigated within 3 to 4 days after application; thoroughly wet the surface of the ground over the row to carry the herbicide into the root

zone of germinating weeds. Alternatively, for control of emerged annual weeds (up to 4" in height) at lay-by time, make a single application in combination with surfactant, or use 1/2 to 3/4 lb "Karmex" DF per acre plus surfactant and repeat if needed.

Replanting Cotton

If initial seeding fails to produce a stand, cotton may be replanted in soil treated preemergence with "Karmex" DF, alone or following "Treflan"(preplant). Wherever possible, avoid disturbing original bed. If necessary to rework soil before replanting, use shallow cultivation such as disking. Do not relist nor move soil into the original drill area. Plant seed at least 1" deep. Do not retreat field with a second preplant or preemergence application during the same crop year as injury may result.

Replanting to Subsequent Crops

For:

Band preemergence or postemergence-

Any crop 4 months after last application.

Band preemergence plus postemergence -or-

Broadcast preemergence/preplant* -or-

Broadcast preemergence plus band postemergence-

Cotton, soybeans, corn or grain sorghums (not sorghos or forage sorghums nor grass sorghums) the next spring. Do not replant treated areas to any other crop within one year after last application as injury may result.

Broadcast preemergence (lay-by)-

Cotton, corn, grain sorghums (not sorghos or forage sorghums nor grass sorghums) the next spring. Do not replant treated areas to any other crop within one year after last application as injury may result.

* For subsequent crops in fields where "Treflan" is used, follow instructions on "Treflan" label.

GRASS SEED CROPS

Perennial Plantings At Least 1 Year Old.

Colorado, Kansas, New Mexico and Oklahoma: On sand bluestem, side oats grama and switchgrass, apply 2 to 3 lbs per acre during the dormant period shortly before weed seedlings emerge. Do not apply after crop begins growth in the spring as crop injury may result. In fields where ash residues have accumulated from burning straw, use 3 lbs per acre; spread unburned chaff or straw with a harrow or chopper before application.

Western Oregon: On alta fescue, Astoria bentgrass, Highland bentgrass, Kentucky bluegrass (Merion bluegrass) and orchardgrass, apply 2 to 4 lbs per acre between October 1 and November 15. In fields where ash residues have accumulated from burning straw, use 3 to 4 lbs per acre; spread unburned chaff or straw with a harrow or chopper before application. If perennial velvetgrass, *Poa lanata*, is a problem, use 4 lbs per acre. For best results, apply as soon as possible after fall rains start. Established weeds beyond 2 to 4 leaf stage should be removed prior to treatment.

Well established vigorous stands of spring-planted tall fescue, Kentucky bluegrass and orchardgrass may be treated the following fall provided the crop is planted before April 1 and treatment is not applied before October 15, use 2 lbs per acre.

New Plantings

Oregon and Washington: For use in newly planted bentgrass, chewing fescue, Kentucky bluegrass, perennial ryegrass, orchardgrass and tall fescue, spray Aqua Nu-Char² or Gro-Safe¹ or other brands of activated charcoal during planting as a 1" band on soil surface at 15 lbs per acre of crop where row spacing is 20" (300 lbs per acre broadcast basis). Mount nozzles to apply directly over seed rows to prevent crop injury. Follow with "Karmex" DF as a single broadcast spray at a rate of 2 1/2 to 3 lbs per acre. Apply as soon as possible after planting but before crop or weeds emerge and before rains or sprinkler irrigation. Fall or spring plantings may be treated. Best results usually occur with early fall plantings. Treatment will not control downy brome or wild oats.

OATS

Do not replant treated areas to any crop within one year after last application as injury may result.

Drill-planted Spring oats

Idaho, Eastern Oregon, Eastern Washington: Use in areas where average annual rainfall exceeds 16". Make a single application of 1 to 1 1/2 lbs per acre after planting, either before or after oats emerge but within 6 weeks of planting. Best results are usually obtained when application is made 3 to 4 weeks after planting. Apply before weeds are 3" to 4" tall.

Drill-planted Winter Oats and Mixtures with Peas or Vetch

Western Oregon and Western Washington: Make a single application of 1 1/2 to 2 lbs per acre as soon as possible after planting but before crop emergence.

PEPPERMINT

Pacific Northwest: Apply 3 lbs per acre after the last cultivation in the spring prior to emergence of peppermint. Do not apply to plantings less than 1 year nor to emerged peppermint as injury may result.

RED CLOVER

Western Oregon: Make a single application of 2 lbs per acre on established red clover stands at least 9 months old. Apply "Karmex" DF when red clover is dormant from October 15 to December 15. Do not apply to seedling red clover. Do not replant treated area to any crop within one year after last application.

Treatment will control annual weeds such as bluegrass, chickweed, hawkbeard, rattail fescue, rye grass and velvetgrass.

SORGHUM, Grain

Southwestern States: Apply 1/4 to 1/2 lb per acre plus surfactant. Apply as a directed postemergence spray after sorghum is 15" tall to control weeds 2" to 4". DO NOT SPRAY OVER TOP OF SORGHUM. Use the lower rate on broadleaf weeds up to 2" tall. Use the higher rate on

grasses up to 2" and broadleaf weeds up to 4" tall. When the lower rate is used, a second application may be made if needed. Do not exceed 1/2 lb per acre. Treatment of weeds under drought stress is usually ineffective.

Do not replant treated areas to crops other than cotton or corn within 4 months following band treatment and 6 months following broadcast treatment as injury may result.

SUGAR CANE

To prevent crop injury on new cane varieties, test tolerance to "Karmex" DF prior to adoption as field practice. Do not treat cane growing on thinly covered subsoils or rocky areas as injury may result. Temporary chlorosis may result from application over emerged cane; to minimize chlorosis, use directed postemergence sprays.

Florida: Preemergence--For high organic soils, apply 2 to 4 lbs per acre as a broadcast or band spray prior to weed emergence after planting or after harvesting plant crop (for ratoon crop).

Postemergence--Make 1 or 2 applications of 2 lbs per acre as needed by directed spray inter-row. Alternatively, for panicum control, make up to 3 applications of 1/2 to 1 lb per acre plus surfactant as a directed spray after cane has emerged but before panicum exceeds 2" in height. Adjust nozzles to spray beneath cane plants and between rows to cover weed foliage and to minimize contact of cane leaves with spray or drift. Do not apply more than 6 lbs total per acre between planting (or ratooning) and harvest.

Hawaii and Puerto Rico: Apply 4 to 8 lbs per acre as a broadcast spray prior to weed emergence after planting or after harvesting plant crop (for ratoon crop). A second and third application of 2 to 4 lbs per acre may be made as a broadcast spray over emerged cane or by directed spray inter-row.

If weeds have emerged, add a surfactant and apply as a directed spray. DO NOT SPRAY OVER TOP OF CANE.

Do not apply more than 3 treatments nor more than 10 lbs per acre in Puerto Rico or 12 lbs per acre in Hawaii between planting (or ratooning) and harvest. Treated areas may be planted to sugar cane or pineapple one year after last application.

Louisiana: Use on plant cane seeded on fallowed ground. Make a single application of 3 to 3 3/4 lbs per acre at either of the following times. August through October--Treat a 2 ft. band over the row after planting of cane, but before weeds or cane emerge. January through April--if shaving and off-barring are practiced, treat a 2 ft. band over the row before weeds or cane emerge.

WHEAT, Winter

Crop injury may result where severe winter stress, disease or insect damage follows application. Winter-sensitive varieties may be less tolerant to "Karmex" DF than winter-hardy varieties. Crop injury may result from failure to observe the following: Do not use on sand or loamy sand soils, nor on gravelly or sandy loams with less than 1% organic matter, nor on thinly covered or exposed subsoil areas (clay knobs). Do not treat wheat planted less than 1" deep. Do not treat wheat where winter climatic conditions

have caused "heaving" of plants. Do not treat wheat plants lacking in vigor due to poor emergence, insect damage, disease, high alkalinity or other causes; do not apply after wheat has reached the "boot" stage. Unless specified otherwise, do not use with surfactants, or nitrogen solution. Do not replant treated areas to any other crop within 1 year after last treatment (except as noted) as injury may result.

Idaho, Oregon and Washington--East of Cascade

Range: Where average annual rainfall exceeds 16 inches, make a single application of 1 to 1 1/2 lbs per acre. For early fall-planted wheat seeded before September 10, apply 3 to 6 weeks after planting but before weeds are 3" to 4" tall. Treatment after October 1 has generally given best results. Application should not be made after soil freezes in the fall. Wheat planted in late October should not be treated until the following spring. For spring treatment, apply as soon as wheat starts to grow. Treatment made prior to April 10 will usually give good results provided weed growth is less than 4" tall. Application later than May 1 may give poor results.

Alternatively, make a single application of 1/2 to 1 lb "Karmex" DF plus 1/4 lb bromoxynil per acre as a tank mix, either in the fall after wheat has emerged but before soil freezes or in the spring as soon as soil thaws; apply before weeds are 2" tall or across.

Where average annual rainfall is 10 to 16 inches following fall planting, make a single application of 1 to 1 1/2 lbs per acre when moisture is available to germinate wheat seed.

Apply before soil freezes and before weeds are 2" tall.

Application later than March 1 may give poor results.

If fall-planted wheat fails to grow due to winter kill or adverse growing conditions after fall treatment, only fields treated before November 1 may be replanted to spring wheat. Spring wheat should not be planted before April 1, and only after deep discing and plowing to a depth of 4" to 6" prior to planting. Do not make a second application during the same crop year as injury may result.

Oregon and Washington--West of Cascade Range:

Make a single application of 1 1/2 to 2 lbs per acre as soon as possible after planting. If wheat and weeds have emerged, apply before weeds are 3" to 4" tall.

Alternatively, apply a tank mix of "Karmex" DF plus bromoxynil as detailed above for "East of Cascade Range".

Other Areas of Oregon and Washington: Make a single application in the spring as soon as wheat (fall-planted) starts to grow and before weeds are 2" tall. Application later than May 1 may give poor results.

Central Plains and Midwest: Use 1 to 2 lbs per acre.

Kansas, Oklahoma and Texas: Do not use on sand or sandy loam soils. Use 1 lb per acre on silt and silt loam soils and 1 1/2 to 2 lbs per acre on clay, clay loam, and silty clay loam soils.

Northeast: Use 1 to 1 1/2 lbs per acre.

FRUIT AND NUT CROPS

Unless otherwise directed, make a single application per year as a directed spray, avoiding contact of foliage and fruit with spray or drift. See "Soil Limitations" section for restrictions. Do not graze livestock in treated orchards or groves.

APPLES

United States

"Karmex" DF Alone--Use only under trees established in the orchard for at least 1 year. Do not treat varieties grafted on full-dwarf root stocks. Apply 4 lbs per acre from March through May. In the Far West, apply 4 lbs per acre to small weeds less than 2" in height or diameter under dormant trees. Alternatively, treatments to small weeds may be applied at 2 lbs per acre postharvest followed by 2 lbs per acre prior to budbreak.

Georgia - Apply 2 to 3 lbs per acre in the spring.

Repeat application in the fall but do not use more than 4 lbs per acre per year. Add surfactant to improve control of small, emerged weeds.

"Karmex" DF + Sinbar® Herbicide--Use only under trees established in the orchard for at least 2 years. Apply either in the spring or after harvest in the fall before weeds emerge or during early seedling stage of weed growth.

"Karmex" DF + "Sinbar" Lbs Per Acre

Soil Texture	1 to 2% Organic Matter	More Than 2% Organic Matter
Sandy loam	1 + 1	1 1/2 + 1 1/2
Loam, silt loam, silt	1 1/2 + 1 1/2	2 + 2
Clay loam, clay	2 + 2	2 + 2

Where crop is grown under furrow irrigation or under raised-berm flood irrigation (trees 4" to 6" above waterline), apply only as a band treatment. Do not treat trees planted in the bottom of irrigation furrows, nor trees grown under flat flood or basin irrigation, as injury may result. Where complete weed control to harvest is desired, additional weed control measures may be required during the growing season.

BANANAS AND PLANTAINS

New Plantings:To control annual weeds, apply 1 1/2 to 3 lbs per acre after planting but before weeds emerge. Do not apply to loose soil directly over the planting material.

Established Plantings: For control of annuals and for top-kill of perennials such as bermudagrass, birdseed grass and guineagrass, apply 3 to 6 lbs per acre plus surfactant. Avoid contact of plants with spray or drift as injury may result. When tall, dense weed growth is present, remove weed growth before application. If application is made to soil free of weeds, omit surfactant from the spray. Repeat treatment as needed. Apply at 6-week intervals or longer for a maximum of 12 lbs per acre (broadcast) in 12-months.

Do not replant treated areas to any crop within 2 years after last application as injury to subsequent crops may result, except that sugar cane or pineapple may be planted after one year.

BLUEBERRIES, CANEBERRIES AND GOOSEBERRIES

Use only in fields that have been established for 1 year or more. Do not apply to berries interplanted with fruit trees. Do not apply to plants where roots are exposed as injury may result. Apply as a band treatment at the base of canes or bushes. For spring application, apply before germination of annual weeds.

Georgia--Blueberries: Apply 1 1/2 to 2 lbs per acre in the spring and repeat treatment after harvest in the fall. Add surfactant to improve control of small, emerged weeds.

Indiana, Michigan and Ohio--Blueberries: Apply 2 to 4 lbs per acre in late spring. Alternatively, apply 2 lbs per acre in the fall and repeat in the spring. **Raspberries:** Apply 3 lbs per acre in the spring.

Massachusetts--Blueberries: Apply 2 lbs per acre in late spring.

New Jersey--Blueberries: For control of winter annuals, apply 2 lbs per acre from October - December, or make a single application of 2 1/2 lbs per acre in early to mid-Spring.

California--Raspberries, Blackberries, Boysenberries, Dewberries and Loganberries: For control of winter annuals, apply 2 lbs per acre in October or November. Repeat at same rate in late spring to control annuals. A single application of 3 lbs per acre in January or February will control annuals in some areas, but the separate fall and spring schedule is preferred.

Western Oregon and Western Washington--Blueberries, Caneberries and Gooseberries: Use California recommendations.

CITRUS

Use only under trees established in the grove for at least 1 year. Time application as indicated for specific areas, except application may be made any time of the year where sprinkler or flood irrigation can be timed to activate the herbicide. Established perennial weeds require other special control procedures. Do not apply under citrus trees that have been subjected to freezing within 6 months.

Arizona (except Yuma area) and California (except Imperial and Coachella Valleys): Apply 3 to 4 lbs per acre shortly after grove has been laid up in final form (nontillage program) in late fall or early winter. Alternatively, apply 2 lbs per acre in October or November and repeat at the same rate in March or April. Subsequent annual applications of 2 to 3 lbs per acre will usually give adequate weed control.

Florida and Puerto Rico: Make a single application of 4 to 8 lbs per acre, or apply 3 to 4 lbs per acre followed by the same rate 4 to 6 months later. On bearing citrus, apply any time when seasonal rains are expected. On nonbearing trees, apply when winter banks are pulled down.

Texas: Apply 2 to 4 lbs per acre for annual weeds. Use 4 to 6 lbs per acre for control of johnsongrass seedlings. Spring treatments give best results. Well established weeds should be eliminated by cultivation prior to treatment.

GRAPES

Apply only as a band treatment to established vineyards at least 3 years old. On soils low in clay or organic matter (1 to 2%), severe plant injury may result if heavy rainfall or more than one inch of irrigation occurs soon after treatment. This risk must be assumed by the user.

East of the Rocky Mountains: On soils low in clay or organic matter (1 to 2%), apply 2 to 3 lbs per acre. On soils high in clay or organic matter, apply 3 to 6 lbs per acre. Apply in the spring just prior to germination of annual weeds.

West of the Rocky Mountains: For best results, apply during the winter months when weeds are less than 2" in height or diameter. Rainfall or overhead sprinkler irrigation sufficient to wet the soil to a depth of 2" is necessary to activate the herbicide. Abnormally heavy rainfall following application just before spring growth may move the herbicide into the root zone of grapes which could result in injury. For initial treatment, apply 3 to 4 lbs per acre. Subsequent annual applications of 2 lbs per acre will usually give adequate weed control. Do not apply to vines with trunks less than 1 1/2" in diameter as injury may result.

New York and Pennsylvania: Use only in established vineyards (at least 4 years old) for spot control of perennial grasses such as orchardgrass, quackgrass and ryegrass. Apply in the spring as a band treatment to ridged soil (2" to 4" high) under the trellis at the rate of 8 to 12 lbs per acre. Band width should not exceed 30". Do not apply more than once every 4 years. Use only on heavy soils such as loams, silt loams, and clay loams. Do not use in areas where grape roots are shallow or exposed because of high bedrock, poor drainage, or erosion as injury may result.

MACADAMIA NUTS

Hawaii: Use only under trees established in the orchard for at least 1 year. Apply 2 to 6 lbs per acre immediately after harvest, preferably before weeds emerge. If weeds have emerged, add surfactant. Retreat as needed but do not exceed 10 lbs per acre per year.

OLIVES

California: Use only under trees established in the grove for at least 1 year. Apply 2 lbs per acre after grove has been laid up in final form in late October or November. Repeat at same rate in March or April. Remove weed growth prior to treatment.

PAPAYAS

Use only under trees established in the orchard for at least 1 year. Apply 2 1/2 to 5 lbs per acre, preferably before weeds emerge. If weeds have emerged, add surfactant.

PEACHES

United States

"Karmex" DF Alone--Use only under trees established in the orchard for at least 3 years. Apply 2 to 5 lbs per acre in the early spring before weeds emerge or during the early seedling stage. Do not apply within 3 months of harvest. In the Far West, do not apply within 8 months of harvest.

Georgia--On trees established for 2 years or more, apply 2 to 3 lbs per acre in the spring. Repeat application in the fall but do not exceed 5 lbs per acre per year. Add surfactant to improve control of small, emerged weeds.

Where crop is grown under furrow irrigation or under raised-berm flood irrigation (trees 4" to 6" above waterline), apply only as a band treatment. Do not treat trees planted in the bottom of irrigation furrows, nor trees grown under flat flood or basin irrigation, as injury may result. Where weed control to harvest is desired, additional measures may be required.

"Karmex" DF + "Sinbar" - United States - See combination recommendation for Apples.

PEARS

Use only under trees established in the orchard for at least 1 year. Do not treat varieties grafted on full-dwarf root stocks. Apply 4 lbs per acre from March through May. In the Far West, apply 4 lbs per acre to weeds less than 2" in height or diameter under dormant trees. Alternatively, apply to small weeds at 2 lbs per acre postharvest followed by 2 lbs per acre prior to budbreak.

PECANS

Use "Karmex" DF alone or apply as a tank mix with "Sinbar". Make a single band or broadcast application as a directed spray using a minimum of 30 gals. of water per acre. Apply in the spring before weeds emerge or during the early seedling stage.

Soil Texture	Lbs Per Acre		Tank Mixture
	"Karmex" DF Alone*	or	"Karmex" DF + "Sinbar" **
Sandy loam	2	or	1 1/2 + 1 1/2
Loam, silt loam, silt	3	or	1 3/4 + 1 3/4
Clay loam, clay	4	or	2 + 2

*Use only on trees established in grove for at least 3 yrs. and on soils with at least 1/2% organic matter.

**Use on trees established in the grove for at least 1 yr. and on soils with at least 1% organic matter.

Do not use on eroded areas where subsoil or roots are exposed, nor on trees that are diseased or lacking in vigor or on trees planted in irrigation furrows as injury to the trees may result.

PINEAPPLE

Hawaii and Florida: Apply 4 to 8 lbs per acre as a broadcast spray just before or immediately after planting but prior to weed emergence. For ratoon crop use 4 lbs per acre after harvesting plant crop. For plant crop only, a second and third broadcast or interspace application may be made prior to differentiation at the rate of 2 lbs per acre at intervals of not less than 2 months. Additional applications to plant crop may be made as needed to interspace only using 2 lbs per acre. Do not apply more than 3 broadcast sprays (maximum 12 lbs per acre) prior to differentiation nor more than 16 lbs total per acre per plant crop. Treated areas may be planted to pineapple or sugar cane 1 year after last application.

Puerto Rico: Apply 3 3/4 to 6 1/4 lbs per acre as a broadcast spray before or immediately after planting but prior to weed emergence. Application controls weeds such as pigweed, crotalaria, morning glory, purslane, crabgrass, foxtail, goosegrass, fall panicum and sourgrass.

WALNUTS, English

California: Use only under trees established in the orchard for at least 1 year. As an initial treatment, apply 3 to 5 lbs per acre after the orchard has been laid up in final form (nontillage program) in late fall or early winter. Retreat annually with 2 to 3 lbs per acre. Alternatively, apply 2 lbs per acre in October or November and repeat in March or April.

NON-CROP WEED CONTROL

"Karmex" DF is an effective herbicide for control of many weeds. The degree of control and duration of effect will vary with the amount of chemical applied, soil texture, rainfall and other conditions.

"Karmex" DF may be used as a preemergence treatment at any time of the year except when ground is frozen, provided adequate moisture is supplied by rainfall or artificial means to activate the herbicide. Best results are obtained if application is made to the soil shortly before weed growth begins. If dense growth is present, remove tops and spray the ground.

Increased contact activity on established weeds may be obtained using a non-ionic surfactant. Apply as a drenching spray to actively growing weeds during warm weather when daily temperature will exceed 70 degrees F.

Use a fixed-boom power sprayer properly calibrated to insure a constant rate of application. Mix proper amount of "Karmex" DF into volume of water necessary to obtain uniform coverage. If surfactant is used, dilute with 10 parts of water and add as last ingredient to nearly full tank.

"Karmex" DF must be kept in suspension at all times.

Agitate by mechanical or hydraulic means in the spray tank. If bypass or return line is used, it should terminate at bottom of tank to minimize foaming. Use 50 mesh or larger screens.

GENERAL WEED CONTROL

To control weeds for an extended period of time on non-cropland such as utility, highway, pipeline and railroad right of ways, petroleum tank farms, lumberyards, storage areas, industrial plant sites, around farm buildings, and similar areas--apply 5 to 15 lbs per acre to control annual weeds including:

Broadleaves

5 to 15 Lbs/Acre

Ageratum	Knawel	Sesbania
Chickweed	Kochia	Shepherd's-purse
Cocklebur	Lambsquarters	Sicklepod
Corn speedwell	Marigold	Smartweed, annual
Corn spurry	Mexican clover	Sowthistle, annual
Dayflower	Morningglory, annual	Spanishneedles
Dogfennel	Pennycress	Tansymustard
Fiddleneck (amsinckia)	Pigweed	Velvetleaf (buttonweed)
Flora's paintbrush	Pincappleweed	Wild buckwheat
Gromwell	Pokeweed	Wild lettuce
Groundcherry, annual	Prickly lettuce	Wild mustard
Hawksbeard	Prickly sida (teaweed)	Wild radish
Horsenettle	Purslane	
Horseweed	Rabbit tobacco	
	Ragweed	

Grasses

5 to 8 Lbs/Acre

Barnyardgrass (watergrass)	Orchardgrass	Seedling johnsongrass
Bluegrass, annual	Peppergrass	Velvetgrass
Crabgrass	Quackgrass	Vernalgrass, sweet, annual
Foxtail	Rattail fescue	
Kyllinga	Red sprangletop	
Lovegrass, annual	Ricegrass	
	Ryegrass, annual	
	Sandbur	

8 to 15 Lbs/Acre

Guineagrass	Maidencane	Pangolagrass
-------------	------------	--------------

Irrigation and Drainage Ditches: Apply 5 to 15 lbs per acre to control annual weeds as shown above. Apply only when water is not in the ditch. For irrigation ditches, apply during the non-crop season, and when ditch is not in use. To avoid crop injury, it is essential to minimize movement of "Karmex" DF in irrigation water. The herbicide must be fixed in the soil by moisture. Apply before expected seasonal rainfall, if possible when soil in the ditch is still moist. Following treatment, if rainfall has not totaled at least 4 inches, fill ditch with water and allow to stand for 72 hours. Drain off and waste remaining water before using ditch. Do not treat any ditch into which roots of trees or other desirable plants may extend as injury may result.

TREE PLANTINGS

Colorado, Montana, Nebraska, North Dakota, South Dakota, Wyoming: Use only under plantings of American elm, caragana, cottonwood, Douglas fir, green ash, honeysuckle, Ponderosa pine, redcedar, Russian olive and Siberian elm, of 1 year or older. Use 2 1/2 to 5 lbs per acre. Apply as a band 4 ft. wide in the tree row (2 ft. on

each side of row). For example, 1 oz. "Karmex" DF treats 135 ft. of tree row (2 ft. on each side of row) at the rate of 5 lbs per acre. Apply as a directed spray in early spring before weeds emerge and before trees leaf out. Do not apply to foliage of trees, nor under trees growing in low areas as injury may result.

STORAGE AND DISPOSAL

STORAGE: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage.

PRODUCT DISPOSAL: Do not contaminate water, food, or feed by disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE TO BUYER - Purchase of this material does not confer any rights under patents of countries outside of the United States.

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. **DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

- 1 Reg. trademark of Elanco Products Co.
 - 2 Reg. trademark of Westvaco Corp.
 - 3 Reg. trademark of ICI United States Inc.
- AG-7654 9043 4/19/93

BEST AVAILABLE COPY