

PLEASE NOTE

This image contains more than one label approved for this product on this date.

SUPPLEMENTAL LABELING

DuPont Crop
Protection

DUPONT™ EXPRESS® XP

HERBICIDE

EXPRESS® HERBICIDE

FOR USE ONLY ON IMAZETHAPYR
TOLERANT CANOLA

DUPONT™ EXPRESS® XP HERBICIDE

EXPRESS® HERBICIDE

EPA REG. NO. 352-509

FOR USE ONLY ON IMAZETHAPYR TOLERANT CANOLA
(CANOLA VARIETIES WITH THE SMART™ TRAIT)

FOR USE IN THE STATES OF MN, ND AND MT

ACCEPTED

NOV 18 2004

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended for the pesticides
registered under
EPA Reg. No. 352-509

EXPRESS® Herbicide or DuPont™ EXPRESS® XP Herbicide (EXPRESS®) can be applied on Imazethapyr Tolerant Canola (eg. canola varieties with the SMART™ trait).

DIRECTIONS FOR USE

Apply EXPRESS® at 1/10 ounce per acre. Add a DuPont recommended, nonionic surfactant having at least 80% active ingredient strength at 0.25% v/v (1 qt per 100 gal of spray solution).

Apply EXPRESS® when the Imazethapyr Tolerant Canola is in the 3 to 6 leaf stage of growth but prior to beginning of bolting. Temporary crop yellowing may be observed shortly after application of EXPRESS®, especially when applied to crops growing under environmentally stressful conditions (See PRECAUTIONS).

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TANK MIXTURES

For the control of annual grasses as well as broadleaf weeds, apply a tank mixture of EXPRESS® at 1/10 ounce per acre plus ASSURE® II Herbicide (See the EPA approved ASSURE II label for use rates, weed size, precautions, and restrictions). Add a DuPont recommended, nonionic surfactant having at least 80% active ingredient strength at 0.25% v/v (1 qt per 100 gal of spray solution).

PRECAUTIONS

- Application of EXPRESS® prior to the 3 leaf stage of Imazethapyr Tolerant Canola could result in crop injury.
- Avoid application to Imazethapyr Tolerant Canola fields in which germination is uneven (some plants outside the recommended leaf stage for application), as crop injury may result.
- Under certain environmentally stressful conditions (such as heavy rainfall, prolonged cool or hot weather, frost conditions, wide fluctuations in day/night temperatures), temporary lightening in crop color and occasionally a slight reduction in crop height may occur.
- Applications to Imazethapyr Tolerant Canola that are or have been stressed prior to application by severe weather conditions, frost, low fertility, drought, water saturated soil, disease or insect damage, may result in crop injury. Applications to Imazethapyr Tolerant Canola if the above stress conditions are expected within 3 days after application, may also result in crop injury.
- Application of the EXPRESS® or EXPRESS® plus ASSURE II Herbicide tank mixture on Imazethapyr Tolerant Canola may delay flowering.

RESTRICTIONS

- DO NOT apply to NON-Imazethapyr Tolerant Canola (i.e. Canola varieties that DO NOT contain the SMART™ trait) as severe crop injury or death of the plant may occur.

For product information call 1-888-6-DUPONT

© 2004 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898

282

IMPORTANT

**BEFORE USING EXPRESS® OR ASSURE II
HERBICIDE, READ AND FOLLOW ALL
APPLICABLE DIRECTIONS, RESTRICTIONS
AND PRECAUTIONS ON THE EPA-REGIS-
TERED LABEL.**

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

DR-406 081004

For product information call 1-888-6-DUPONT
© 2004 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898

NEXT

LABEL

SUPPLEMENTAL LABELING

**DuPont Crop
Protection**

**DUPONT™ EXPRESS® XP
HERBICIDE
EXPRESS® HERBICIDE
BURNDOWN IN COTTON**

ACCEPTED

NOV 18 2004

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended for the pesticide
registered under
EPA Reg. No. 352-509

**DUPONT™ EXPRESS® XP HERBICIDE
EXPRESS® HERBICIDE**

EPA Reg. No. 352-509

PRE-PLANT BURNDOWN IN COTTON

Dupont™ EXPRESS® Herbicide or DuPont™ EXPRESS® XP Herbicide (EXPRESS®) may be applied for burndown of emerged weeds prior to the emergence of cotton.

DIRECTIONS FOR USE

EXPRESS® may be used as part of a pre-plant burndown treatment, in combination with other suitable registered pre-plant herbicides.

Read and follow all manufacturers label recommendations for the companion herbicide. If those recommendations conflict with this label, do not tank mix the herbicide with EXPRESS®. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply EXPRESS® at 0.2 ounce per acre for control or partial control of the weeds listed on the EPA registered label. Allow at least 14 days between application of EXPRESS® and planting of cotton. Include a nonionic surfactant, petroleum based crop oil concentrate, or vegetable-seed oil-based product (methylated seed oils are considered a vegetable seed-based oil).

- If another herbicide is tank mixed with EXPRESS® to increase the broadleaf weed spectrum, select adjuvants based on the adjuvant limitations of the companion herbicide.

SPRAY ADJUVANTS

Nonionic Surfactant (NIS)

Apply at a rate (concentration) of 0.25-0.5% v/v (1-2 qt per 100 gal spray solution). Use the higher rate in hot and dry conditions to enhance control.

Crop Oil Concentrate

Under dry conditions or during cool weather, a petroleum based crop oil concentrate, or vegetable-seed oil-based product may be used in place of a nonionic surfactant at 1-2 gallon/100 gal of spray solution (1-2% v/v) to enhance weed control. Use a petroleum-based crop oil concentrate with at least 14% emulsifiers/surfactant and 80% oil.

Ammonium Nitrogen Fertilizer

An ammonium nitrogen fertilizer can be added to a surfactant or a crop oil concentrate to enhance control. Alternatively, a high-quality, sprayable grade of ammonium sulfate (21-0-0) may be used.

IMPORTANT PRECAUTIONS

Seedling disease, nematodes, cold weather, deep planting (more than 2"), excessive moisture, high salt concentration, and/or drought may weaken cotton seedlings and increase the possibility of crop injury. Cotton resumes normal growth once favorable growing conditions return.

RESTRICTION

- DO NOT apply later than 14 days before planting cotton.

IMPORTANT

BEFORE USING EXPRESS®, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

DR-415 082604

For product information all 1-888-6-DUPONT

© 2004 E. I. du Pont de Nemours and Company, DuPont Crop Protection, Wilmington, Delaware 19898

NEXT

LABEL

SUPPLEMENTAL LABELING

**DuPont Crop
Protection**

**DUPONT™ EXPRESS®
HERBICIDE
EXPRESS® HERBICIDE
FOR USE ONLY ON
CDC TRIFFID FLAX**

ACCEPTED

NOV 18 2004

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended for the pesticide
registered under
EPA Reg. No. 352-509

**DUPONT™ EXPRESS® XP HERBICIDE
EXPRESS® HERBICIDE**

EPA REG. NO. 352-509

**FOR USE ONLY ON CDC TRIFFID FLAX
IN THE STATES OF MN, ND AND MT**

EXPRESS® Herbicide or DuPont™ EXPRESS® XP Herbicide (EXPRESS®) can be applied on flax that contains the "Freedom™" trait (CDC Triffid Flax).

DIRECTIONS FOR USE

Apply EXPRESS® at 1/10 ounce per acre. Add a DuPont recommended, nonionic surfactant having at least 80% active ingredient strength at 0.25% v/v (1 qt per 100 gal of spray solution).

Apply EXPRESS® from the time the CDC Triffid Flax is 1/2 inch in crop height to the pre-bud stage. Temporary crop yellowing may be observed shortly after application of EXPRESS® Herbicide, especially when applied to crops growing under environmentally stressful conditions (See PRECAUTIONS).

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS

- Application of EXPRESS® prior to CDC Triffid Flax crop height reaching 1/2 inch tall could result in crop injury.
- Avoid application to CDC Triffid Flax fields in which germination is uneven (some plants outside the recommended leaf stage for application), as crop injury may result.
- Under certain environmentally stressful conditions (such as heavy rainfall, prolonged cool or hot weather, frost conditions, wide fluctuations in day/night temperatures), temporary lightening in crop color and occasionally a slight reduction in crop height may occur.

- Applications to CDC Triffid Flax that is or has been stressed prior to application by severe weather conditions, frost, low fertility, drought, water saturated soil, disease or insect damage, may result in crop injury. Applications to CDC Triffid Flax if the above stress conditions are expected within 3 days after application may also result in crop injury.

RESTRICTIONS

- Only apply to flax that contains the "Freedom™" trait. EXPRESS® will severely damage flax varieties that do not contain the "Freedom™" trait.

IMPORTANT

BEFORE USING EXPRESS®, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

DR-405 081004

For product information call 1-888-6-DUPONT

© 2004 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898