

352-509

01/26/2006

1/19



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES  
TOXIC SUBSTANCES

JAN 26 2006

AND

Mr. Jaccob J. Vukich  
E.I. DuPont de Nemours & Company  
DuPont Crop Protection  
Stine-Haskell Research Center  
P. O. Box 30  
Newark, DE 19714-0030

Dear Mr. Vukich:

Subject: DuPont Express XP Herbicide (Reformat Label & Incorporate Supplementals)  
EPA Registration No. 352-509  
Letter Dated December 9, 2004

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable provided you make the following changes before you release the product for shipment.

1. Delete the statement "Wear long-sleeved shirt and long pants, socks and shoes" from your Precautionary Statements, since the information appears under Personal Protective Equipment (PPE).
2. On page 12 revise the heading "Pesticide Storage and Disposal" to read "Storage and Disposal".

Submit one copy of your final printed labeling incorporating the above changes before you release the product for shipment. Amended labeling supercedes all previously accepted ones. A stamped copy of labeling is enclosed for your records.

Sincerely,

*Vickie K Walters for*  
James A. Tompkins  
Product Manager 25  
Herbicide Branch  
Registration Division (7505C)

H-  
2/19




# DuPont™ Express® XP

herbicide

---

## DRAFT LABEL



*“..... A Growing Partnership With Nature”*

# DUPONT™ EXPRESS® XP HIGHLIGHTS

- *For selective postemergence broadleaf weed control in Wheat, Barley, Triticale, Post Harvest in Cereals, Fallow and Pre-plant Burndown.*
- *Apply at the rate of 1/6 to 1/3 ounce per acre (see Application Information).*
- *In wheat, barley and triticale apply after the crop is in the 2-leaf stage, but before the flag leaf is visible.*
- *In fallow apply when the majority of weeds have emerged and are actively growing.*
- *As a burndown treatment to wheat (including durum), barley or triticale to control emerged weeds prior to, or shortly after planting (prior to emergence). Make applications when the majority of weeds have emerged and are actively growing.*
- *As a Post Harvest burndown treatment to crop stubble apply when the majority of weeds have emerged and are actively growing.*
- *May be applied by ground or by air.*
- *Use in tank mixtures with other registered herbicides for broader spectrum weed control (see Tank Mixtures).*
- *Can rotate to any crop 45 - 60 days after last application.*
- *Consult label text for complete instructions. Always read and follow label "Directions For Use".*

# TABLE OF CONTENTS

PRECAUTIONARY STATEMENTS .....1  
 DIRECTIONS FOR USE .....2  
 GENERAL INFORMATION .....2  
   Environmental Conditions and Biological Activity ... 2  
 FALLOW .....  
   Use Rate ..... 2  
   Application Timing .....2  
   Tank Mixtures .....2  
 PREPLANT BURNDOWN .....  
   Application Timing .....3  
   Rates and Adjuvants .....3  
   Tank Mixtures .....3  
   Application Equipment and Volumes .....3  
   Weeds Controlled .....3  
 WHEAT, BARLEY AND TRITICALE .....  
   Use Rate ..... 3  
   Application Timing .....3  
   Weeds Controlled .....4  
   Tank Mixtures .....4  
   Liquid Nitrogen Fertilizer Solutions .....5  
   Specific Weed Problems .....6  
 GRASS GROWN FOR SEED .....6  
 SPRINKLER CHEMIGATION .....7  
 POST HARVEST .....8  
 GENERAL INFORMATION .....8  
   Spray Adjuvants .....8  
   Ground Application .....8  
   Aerial Application .....8  
   Product Measurement .....8  
 CROP ROTATION .....8  
 GRAZING .....9  
 MIXING INSTRUCTIONS .....9  
 SPRAY EQUIPMENT .....9  
 SPRAYER CLEANUP .....9  
 SPRAY DRIFT MANAGEMENT .....10  
 RESISTANCE ..... 10  
 INTEGRATED PEST MANAGEMENT .....11  
 PRECAUTIONS .....11  
 STORAGE AND DISPOSAL .....11  
 NOTICE OF WARRANTY .....12

4/19



# DuPont™ Express® XP

herbicide

Dry Flowable

For Use on Wheat, Barley, Triticale, Fallow  
and Pre-plant burndown

Active Ingredient	By Weight
Tribenuron methyl:	
Methyl 2-[[[(4-methoxy-6-methyl -1,3,5-triazin-2-yl)methylamino]carbonyl] amino]sulfonyl]benzoate	75%
Inert Ingredients	25%
TOTAL	100%

EPA Reg. No. 352-509

EPA Est. No.

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:**

**JAN 26 2006**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.

352-509

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

### FIRST AID

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for emergency medical treatment information.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION!** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long-sleeved shirt and long pants, socks and shoes. Wash thoroughly with soap and water after handling.

For medical emergencies involving this product, call toll-free 1-800-441-3637.

### PERSONAL PROTECTIVE EQUIPMENT

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical-resistance category selection chart.

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride.
- Shoes plus socks.

Discard clothing and other absorbent materials that have been drenched, or heavily contaminated with this product. Follow manufacturer's instructions for cleaning / maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Control Statement:** When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR Part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

**USERS SHOULD:** Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposing of equipment washwaters or wastes.

## PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Ensure that all operation employees accurately measure pesticides.
- Mix only enough product for the job at hand.
- Avoid overfilling of spray tank.
- Do not discharge excess material on the soil at a single spot in the field, grove, or mixing/loading station.
- Dilute and agitate excess solution and apply at labeled rates or uses.
- Avoid storage of pesticides near well sites.
- When triple-rinsing the pesticide container, be sure to add the rinsate to the spray mix.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Chemical Resistant Gloves made of any waterproof material such as polyethylene or polyvinyl chloride.
- Shoes plus socks.

DuPont™ EXPRESS® XP should be used only in accordance with recommendations on this label or in separately published DuPont recommendations.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by DuPont.

EXPRESS® XP is recommended for use on wheat, barley, triticale, post-harvest burndown, fallow and pre-plant burndown in most states. Check with your state extension service or Department of Agriculture before use, to be certain EXPRESS® XP is registered in your state.

## GENERAL INFORMATION

EXPRESS® XP is a dry flowable granule that is used for selective postemergence weed control in wheat (including durum), barley, triticale, post-harvest burndown, fallow and pre-plant burndown. The best control is obtained when EXPRESS® XP is applied to young, actively growing weeds. The use rate will depend on weed spectrum and size of weed at time of application. The degree and duration of control may depend on the following:

- weed spectrum and infestation intensity
- weed size at application
- environmental conditions at and following treatment

EXPRESS® XP is noncorrosive, nonflammable, nonvolatile, and does not freeze. EXPRESS® XP should be mixed in water and applied as a uniform broadcast spray.

## USE RATE

Apply 1/6 to 1/3 oz EXPRESS® XP per acre to wheat (including durum), barley, triticale, fallow and pre-plant burndown. Two applications of EXPRESS® XP may be made per season provided the total amount applied does not exceed 1/3 oz per acre.

## BIOLOGICAL ACTIVITY AND ENVIRONMENTAL CONDITIONS

EXPRESS® XP is absorbed through the foliage of broadleaf weeds, rapidly inhibiting their growth. Leaves of susceptible plants appear chlorotic from 1 to 3 weeks after application and the growing point subsequently dies.

EXPRESS® XP provides the best control in vigorously growing crops that shade competitive weeds. Weed control in areas of thin crop stand or seeding skips may not be as satisfactory. However, a crop canopy that is too dense at application can intercept spray and reduce weed control.

EXPRESS® XP may injure crops that are stressed from adverse environmental conditions (such as extreme temperatures or moisture), abnormal soil conditions, or cultural practices. In addition, different varieties of the crop may have differing levels of sensitivity to treatment with EXPRESS® XP under otherwise normal conditions.

Treatment of sensitive crop varieties may injure crops. To reduce the potential of crop injury, tank mix EXPRESS® XP with 2,4-D (ester formulations perform best—see the Tank Mixtures section of this label) and apply after the crop is in the tillering stage of growth.

In warm, moist conditions, the expression of herbicide symptoms is accelerated in weeds; in cold, dry conditions, expression of herbicide symptoms is delayed. In addition, weeds hardened-off by drought stress are less susceptible to EXPRESS® XP.

Weed control may be reduced if rainfall or snowfall occurs soon after application. Several hours of dry weather are needed to allow EXPRESS® XP to be sufficiently absorbed by weed foliage.

## FALLOW

### FALLOW USE RATE

Apply 1/6 to 1/3 oz DuPont™ EXPRESS® XP per acre to fallow fields. Two applications of EXPRESS® XP may be made per crop season provided the total amount applied does not exceed 1/3 oz per acre.

EXPRESS® XP should be applied in combination with other suitable registered fallow herbicides (See the "TANK MIXTURES" section of this label for additional information)

### APPLICATION TIMING

EXPRESS® XP may be used as a fallow treatment when the majority of weeds have emerged and are actively growing.

### TANK MIXTURES IN FALLOW

EXPRESS® XP may be used as a fallow treatment, and should be tank mixed with other herbicides that are registered for use in fallow. Read and follow all manufacturers' label recommendations for the companion herbicide. If those recommendations conflict with this label, do not tank mix the herbicide with EXPRESS® XP.

## PREPLANT BURNDOWN

### RATES AND ADJUVANTS

Apply EXPRESS® XP at 1/6 - 1/3 oz per acre. Use the higher rate for denser weed populations or where weeds are approaching the maximum size. Also use the higher rate when the weed infestation predominantly consists of those weeds listed in the "Weeds Partially Controlled" section below, or when application timing and environmental conditions are marginal.

Add a spray adjuvant to the tank. Crop Oil Concentrate is preferred at 1% v/v (1 gallon per 100 gallons of final spray volume). Refer to the "Spray Adjuvants" section of this label.

Sequential treatments of EXPRESS® XP may be made provided the total amount of EXPRESS® XP applied during one fallow/pre-plant cropland season does not exceed 1/3 ounce per acre.

Cotton Pre-plant Burndown: Apply 1/5 oz per acre. Allow at least 14 days from time of application to planting cotton.

### APPLICATION TIMING

Apply EXPRESS® XP as a burndown treatment to control emerged weeds in the fall or spring. Make applications when the majority of weeds have emerged and are actively growing.

Applications can be made:

- prior to wheat. Apply before planting or shortly after planting (and prior to emergence).
- prior to cotton. Allow at least 14 days between application and planting.
- prior to sugarbeets, winter rape and canola. Allow at least 60 days between application and planting.
- prior to any other crops (such as corn, grain sorghum, rice and soybeans). Allow at least 45 days between application and planting.

### TANK MIXTURES

Addition of a minimum of 1/2 lb active ingredient per acre of 2,4 D LVE ester (e.g. 1 pt of a 4 lb/gal 2,4-D LVE formulation) is recommended for best results, and required for burndown of some weeds.

EXPRESS® XP may be mixed with one or more other suitably registered herbicides for expanded weed size, or weed spectrum, and/or to add residual control. Read and follow all manufacturers label recommendations for the companion herbicide. If those recommendations conflict with this label, do not tank mix the herbicide with EXPRESS® XP.

### APPLICATION EQUIPMENT AND SPRAY VOLUMES

Apply uniformly by ground equipment using a properly calibrated fixed boom sprayer. For burndown applications of existing vegetation, use with spray nozzles that provide thorough coverage of the weeds.

## WEEDS CONTROLLED - UP TO 3" IN HEIGHT OR DIAMETER - FALLOW AND PRE-PLANT BURNDOWN

Black mustard	Hairy buttercup
Blue/purple mustard	Kochia
Bushy wallflower/Treacle mustard	Marestail *
Canada thistle	Mayweed chamomile
Coast fiddleneck	Miners lettuce
Corn spurry	Pineappleweed
Common chickweed	Poison hemlock*
Common groundsel	Prickly lettuce **
Common lambsquarters	Purslane speedwell
Common purslane	Russian thistle
Cressleaf groundsel (butterweed)	Shepherd's-purse**
Dandelion *	Slimleaf lambsquarters
Deadnettle **	Small-flower buttercup
Early whitlowgrass	Smallseed falseflax
False chamomile	Tarweed fiddleneck
Field pennycress	Tumble / Jim hill mustard*
Flixweed	Wild chamomile
Henbit**	Wild mustard
	Wild parsnip *

\* 2,4-D LVE addition required

\*\* 2,4-D LVE addition recommended

## WEEDS PARTIALLY CONTROLLED\*\*\* UP TO 3" IN HEIGHT OR DIAMETER - FALLOW AND PRE-PLANT BURNDOWN

Annual sowthistle	Prostrate knotweed
Common sunflower (volunteer)	Redroot pigweed
Common vetch	Redmaids
Hairy vetch	Tansymustard
Hairy nightshade	Wild buckwheat
Pennsylvania smartweed	Wild garlic
	Wild radish

\*\*\* partially controlled weeds exhibit a visual reduction in numbers as well as a significant loss of vigor.

## WHEAT, BARLEY AND TRITICALE

### USE RATE

Use 1/3 oz EXPRESS® XP per acre for heavy infestation of those weeds listed in the "WEEDS PARTIALLY CONTROLLED" section of this label and/or when application timing and environmental conditions are marginal (refer to "BIOLOGICAL ACTIVITY AND ENVIRONMENTAL CONDITIONS" section of this label for best performance).

Use 1/6 to 1/4 oz EXPRESS® XP per acre for light infestation of weeds listed in the "WEEDS CONTROLLED" section of this label. Conditions at application should be optimum for effective treatment of these weeds.

**APPLICATION TIMING**

Apply DuPont™ EXPRESS® XP after the crop is in the 2-leaf stage, but before the flag leaf is visible. Do not harvest within 45 days of the last application.

Since EXPRESS® XP has very little or no soil activity, it controls only those weeds that have germinated; therefore, apply EXPRESS® XP when all or most of the weeds have germinated. For best results, annual broadleaf weeds should be past the cotyledon stage, actively growing, and less than 4" tall or wide. See the "SPECIFIC WEED PROBLEMS" section of this label for more information.

Rainfall immediately after treatment can wash EXPRESS® XP off of weed foliage, resulting in reduced weed control. Several hours of dry weather are needed to allow EXPRESS® XP to be sufficiently absorbed by weed foliage.

**WEEDS CONTROLLED**

EXPRESS® XP effectively controls the following weeds when used according to label directions:

- |  |                                    |
|--|------------------------------------|
| Black mustard                                | Hairy buttercup                    |
| Blue/Purple mustard                          | Kochia ** ‡                        |
| Bushy wallflower/Treacle mustard             | London Rocket                      |
| Canada thistle **                            | Mayweed chamomile /                |
| Coast fiddleneck                             | Stinking Chamomile /               |
| Common Chickweed                             | dog fennel (Anthemis cotula L.) ** |
| Common Groundsel                             | Miners lettuce                     |
| Common Lambsquarters                         | Pineappleweed                      |
| Common Purslane                              | Prickly lettuce ** ‡               |
| Corn Gromwell **                             | Redroot pigweed                    |
| Corn spurry                                  | Russian thistle ** ‡               |
| Cowcockle                                    | Shepherd's-purse                   |
| Curly Dock **                                | Slimleaf lambsquarters             |
| False chamomile / Wild chamomile /           | Smallseed falseflax                |
| Scentless chamomile (Matricaria maritima L.) | Tansymustard                       |
| Field pennycress                             | Tarweed fiddleneck                 |
| Flixweed                                     | Tumble/Jim Hill mustard **         |
|  | Wild mustard                       |

**WEEDS PARTIALLY CONTROLLED\***

EXPRESS® XP partially controls the following weeds when used according to label directions:

- |                                 |                        |
|---------------------------------|------------------------|
| Annual sowthistle               | Henbit                 |
| Common cocklebur                | Pennsylvania smartweed |
| Common sunflower (volunteer) ** | Prostrate knotweed     |
| Common Vetch **                 | Redmaids               |
| Hairy nightshade                | Wild buckwheat         |
| Hairy Vetch **                  | Wild garlic            |
|                                 | Wild radish**          |

\* Partially controlled weeds exhibit a visual reduction in numbers as well as a significant loss of vigor. For better results, use 1/4 to 1/3 oz EXPRESS® XP per acre and include a tankmix partner such as 2,4-D, MCP, bromoxynil (such as "Buctril", "Bison", "Bronate" or "Bronate Advanced"), or dicamba (such as "Banvel"/ "Clarity"). Refer to the "TANK MIXTURES" section of this label.

\*\* See the Specific Weed Problems section of this label for more information.

‡ Naturally occurring resistant biotypes of kochia, prickly lettuce and Russian thistle are known to occur. See the "TANK MIXTURES" and "SPECIFIC WEED PROBLEMS" sections of this label for additional details.

**TANK MIXTURES**

EXPRESS® XP may be tank mixed with other suitable registered herbicides to control weeds listed as partially controlled, weeds resistant to EXPRESS® XP, or weeds not listed under Weeds Controlled. Read and follow all

manufacturers label recommendations for the companion herbicide. If those recommendations conflict with this label, do not tank mix the herbicide with EXPRESS® XP.

**With 2,4-D (amine or ester) or MCP (amine or ester)**

EXPRESS® XP may be tank mixed with 2,4-D and MCP (preferably ester formulations) herbicides for use on wheat, barley and triticale. For best results, add 2,4-D or MCP herbicides to the tank at 1/8 to 3/8 lb active ingredient per acre. In tank mixes containing 1/8 lb active ingredient 2,4-D or MCP per acre, add 1 to 2 pt of nonionic surfactant; in tank mixes containing 1/4 to 3/8 lb active ingredient 2,4-D or MCP per acre, add 1 pt of nonionic surfactant.

Higher rates of 2,4-D or MCP may be used, but do not exceed the highest rate allowed by those respective labels. When using rates of 3/8 lb ai per acre or higher, use of additional nonionic surfactant may not be needed, unless specified otherwise in the 2,4-D or MCP label, or local recommendations.

Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using these tank mixtures.

**With 2,4-D or MCP (amine or ester) and Dicamba (such as "Banvel"/"Clarity")**

EXPRESS® XP may be applied in a 3-way tank mix with formulations of dicamba (such as "Banvel"/"Clarity") and 2,4-D or MCP. Observe all applicable directions, restrictions and precautions on labels of all products used.

Make applications at 1/8 - 1/3 oz of EXPRESS® XP + 1-1.5 oz active ingredient dicamba (such as "Banvel"/ "Clarity") + 1/4 to 3/8 lb active ingredient of 2,4-D or MCP (ester or amine) per acre. Use higher rates when weed infestation is heavy. Add 1-2 pt of nonionic surfactant to the 3 way mixture, where necessary, as deemed by local recommendations. Use of additional nonionic surfactant may not be needed with the higher phenoxy rates and ester phenoxy formulations. Consult the specific 2,4-D or MCP and dicamba labels, or local recommendations for more information.

Apply this 3-way combination to winter wheat after the crop is tillering and prior to jointing (first node). In Spring Wheat (including Durum), apply after the crop is tillering and before it exceeds the 5-leaf stage.

Tank mixtures of EXPRESS® XP plus dicamba (such as "Banvel"/ "Clarity") may result in reduced control of some broadleaf weeds.

Do not apply this 3-way mixture at high rates more than once a year, or more than twice per year at the low rates.

**With Bromoxynil containing products (such as "Buctril", "Bison", "Bronate", "Bronate Advanced" or "Rhino")**

EXPRESS® XP may be tank mixed with bromoxynil containing herbicides registered for use on wheat, barley or triticale. For best results, add bromoxynil-containing herbicides to the tank at 3 to 6 oz active ingredient per acre (such as "Bronate" or "Bison" at 3/4 - 1 1/2 pt per acre). Note that tank mixtures of EXPRESS® XP plus bromoxynil may result in reduced control of Canada thistle.

Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using these tank mixtures. Follow the most restrictive labeling.

**With "Starane", "Starane + Salvo" or "Starane + Sword"**

For improved control of Kochia (2-4" tall), DuPont™ EXPRESS® XP may be tank mixed with 1/3 to 2/3 pints per acre of Starane, 2/3 to 1 1/3 pints per acre of Starane + Salvo, or 3/4 to 1 1/2 pints per acre of Starane + Sword. Refer to the DuPont herbicide label, and the Starane, Starane + Salvo, and Starane + Sword labels for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on any of the labels will apply. Do not use the tank mix if any restriction on the labels conflict with recommendations on the DuPont herbicide label.

2,4-D and MCP herbicides (preferably ester formulations) may be tank mixed with EXPRESS® XP plus Starane. Consult local recommendations and the Tank Mixtures section of this label for additional information.

**With "Maverick"**

EXPRESS® XP can be tank mixed with "Maverick" herbicide for improved control of weeds in wheat.

Refer to the "Maverick" label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. Do not use the tank mix if any restrictions on the "Maverick" label conflict with recommendations on the DuPont herbicide label.

**With "Aim"**

EXPRESS® XP can be tank mixed with "Aim" herbicide for improved control of weeds in wheat and barley.

Refer to the "Aim" label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. Do not use the tank mix if any restrictions on the "Aim" label conflict with recommendations on the DuPont herbicide label.

**With "Stinger", or "Curtail", or "Curtail M", or "WideMatch"**

EXPRESS® XP can be tank mixed with "Stinger", "Curtail", "Curtail M", or "Widematch" herbicides for improved control of weeds in wheat and barley.

Refer to the "Stinger", "Curtail", "Curtail M", or "Widematch" label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. Do not use the tank mix if any restrictions on the "Stinger", "Curtail", "Curtail M", or "Widematch" label conflict with recommendations on the DuPont herbicide label.

**With "Assert" Herbicide**

EXPRESS® XP can be tank mixed with "Assert". When tank mixing EXPRESS® XP with "Assert", always include another broadleaf weed herbicide with a different mode of action - for example 2,4-D ester, MCP ester, or bromoxynil (such as "Buctril", "Bison", "Bronate", "Bronate Advanced" or "Rhino"). Applications of EXPRESS® XP plus "Assert" may cause temporary crop discoloration, stunting, or injury when heavy rainfall occurs shortly after application.

**With "Puma"**

EXPRESS® XP can be tank mixed with "Puma" herbicide for improved control of weeds in wheat and barley.

Refer to the "Puma" label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. Do not use the tank mix if any restrictions on the "Puma" label conflict with recommendations on the DuPont herbicide label.

**With "Discover" or "Discover" NG**

EXPRESS® XP can be tank mixed with "Discover" or "Discover" NG herbicide for improved control of weeds in spring wheat.

Refer to the "Discover" or "Discover" NG label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. Do not use the tank mix if any restrictions on the "Discover" or "Discover" NG label conflict with recommendations on the DuPont herbicide label.

**With "Everest"**

EXPRESS® XP can be tank mixed with "Everest" herbicide for improved control of weeds in spring wheat.

Refer to the "Everest" label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. Do not use the tank mix if any restrictions on the "Everest" label conflict with recommendations on the DuPont herbicide label.

**With Other Herbicides**

Tank mixtures of EXPRESS® XP plus metribuzin may result in reduced control of wild garlic.

Tank mixtures of EXPRESS® XP with "Hoelon 3EC" may result in reduced grass control.

**With Fungicides**

EXPRESS® XP may be tank mixed or used sequentially with fungicides registered for use on cereal crops.

**With Insecticides**

EXPRESS® XP may be tank mixed or used sequentially with insecticides registered for use on cereal crops. However, under certain conditions (drought stress, or if the crop is in the 2 to 4 leaf stage), tank mixtures or sequential applications of EXPRESS® XP with organophosphate insecticides (such as "Lorsban") may produce temporary crop yellowing or, in severe cases, crop injury. The potential for crop injury is greatest when wide fluctuations in day/night temperatures occur just prior to or soon after application.

Test these mixtures in a small area before treating large areas.

Do not apply EXPRESS® XP within 60 days of crop emergence where an organophosphate insecticide has been applied as an in-furrow treatment since crop injury may result.

Do not use EXPRESS® XP plus Malathion since crop injury may result.

**With Liquid Nitrogen Solution Fertilizer**

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing



DuPont™ EXPRESS® XP in fertilizer solution. EXPRESS® XP must first be pre-slurried with water and then added to liquid nitrogen solutions (e.g., 28-0-0, 32-0-0). Ensure that the agitator is running while the EXPRESS® XP is added. Use of this mixture may result in temporary crop yellowing and stunting.

If using low rates of liquid nitrogen fertilizer in the spray solution (less than 50% of the spray solution volume), the addition of surfactant is necessary. Add surfactant at 1/2 pt – to 1 qt per 100 gal of spray solution (0.06 – to 0.25% v/v) based on local recommendations.

When using high rates of liquid nitrogen fertilizer solution in the spray solution, adding surfactant increases the risk of crop injury. If 2,4-D or MCP is included with EXPRESS® XP and fertilizer mixture, ester formulations tend to be more compatible (see manufacturer's label). Additional surfactant may not be needed when using EXPRESS® XP in tank mix with 2,4-D ester or MCP ester and liquid nitrogen fertilizer solutions. Consult your agricultural dealer, consultant, field advisor, or DuPont representative for a specific recommendation before adding an adjuvant to these tank mixtures.

Note: In certain areas east of the Mississippi river unacceptable crop response may occur with use of straight or dilute nitrogen fertilizer carrier solutions where cold temperatures or widely fluctuating day/night temperatures exist. In these areas consult your agricultural dealer, consultant, field advisor, or DuPont representative for a specific recommendation before using nitrogen fertilizer carrier solutions.

Do not use low rates of liquid nitrogen fertilizer solution as a substitute for a surfactant.

Liquid nitrogen fertilizer solutions that contain sulfur can increase crop response.

Do not use with liquid fertilizer solutions with a pH less than 3.0.

### SPECIFIC WEED PROBLEMS

**Canada thistle:** For best results, apply 1/3 oz of EXPRESS® XP per acre when all thistles are 4" to 8" tall with 2" to 6" of new growth. Make the application in the spring.

**Corn Gromwell :** For best results, apply 1/3 oz of EXPRESS® XP per acre in combination with 2,4-D or MCP (refer to the Tank Mixtures section of this label).

**Curly Dock :** For best results, apply 1/4 to 1/3 oz of EXPRESS® XP per acre in combination with 2,4-D or MCP (refer to the Tank Mixtures section of this label).

**Kochia:** Naturally occurring biotypes resistant to EXPRESS® XP are known to occur. For best results, use EXPRESS® XP in a tank mixture with Starane, Starane + Salvo, Starane + Sword, dicamba (such as "Banvel"/ "Clarity") and 2,4-D or MCP (ester or amine), or bromoxynil containing products (such as "Buctril", "Bison", "Bronate", "Bronate Advanced" or "Rhino").

EXPRESS® XP should be applied in the spring when kochia are less than 2" tall and are actively growing (refer to the Tank Mixtures section of this label for additional details on rates and restrictions).

**Mayweed chamomile / Stinking Chamomile / dog fennel:** For best results, apply 1/4 to 1/3 oz of EXPRESS® XP per acre.

**Russian thistle, Prickly lettuce:** Naturally occurring biotypes resistant to EXPRESS® XP of these weeds are known to occur. For best results, use EXPRESS® XP in a tank mixture with dicamba (such as "Banvel"/ "Clarity") and 2,4-D or MCP (ester or amine), or bromoxynil containing products (such as "Buctril", "Bison", "Bronate", "Bronate Advanced" or "Rhino").

EXPRESS® XP should be applied in the spring when Russian thistle, and prickly lettuce are less than 2" tall or 2" across and are actively growing (refer to the Tank Mixtures section of this label for additional details on rates and restrictions).

**Tumble/Jim Hill mustard:** For best results, apply 1/3 oz of EXPRESS® XP per acre in combination with 2,4-D or MCP (refer to the Tank Mixtures section of this label).

**Vetch (common and hairy):** For best results, apply 1/4 to 1/3 oz of EXPRESS® XP per acre when vetch is less than 6" in length. For severe infestations of vetch, or when vetch is greater than 6" in length, apply EXPRESS® XP in combination with 2,4-D or MCP (refer to the Tank Mixtures section of this label).

**Wild radish:** For best results, apply 1/6 - 1/3 oz EXPRESS® XP per acre, plus 1/4 – to 3/8 lb active ingredient per acre MCP, plus 0.25% v/v nonionic surfactant (1 qt per 100 gal of spray solution) to wild radish rosettes less than 6" diameter. Make the application either in the fall or spring. Applications made later than 30 days after weed emergence will result in partial control. Fall applications should be made before plants harden-off.

**SU / IMI Tolerant Volunteer Sunflowers:** Varieties resistant to SU and IMI products (like EXPRESS® XP, "Beyond", "Pursuit", "Raptor") are under development. For best results, use EXPRESS® XP in a tank mix with Starane, Starane + Salvo, Starane + Sword, dicamba (such as "Banvel"/ "Clarity") and 2,4-D or MCP (ester or amine), or bromoxynil containing products (such as "Buctril", "Bison", "Bronate", "Bronate Advanced" or "Rhino").

### GRASS GROWN FOR SEED ONLY IN THE STATES OF IDAHO, OREGON, AND WASHINGTON

#### GENERAL INFORMATION

EXPRESS® XP Herbicide is recommended for selective postemergence control/suppression of certain broadleaf weeds in seedling and established stands of bentgrass, bluegrass, annual ryegrass, orchardgrass, tall fescue, fine fescue and Timothy grown for seed.

EXPRESS® XP may be used on seedling and established perennial ryegrass providing user accepts all risk of possible crop injury and/or reduced seed yield. See the "USE RATES" section of this label.

EXPRESS® XP may cause temporary yellowing and stunting of grass. Best results are obtained when EXPRESS® XP is applied to young, actively growing weeds. The degree of control and duration of effect are dependent on the rate used, sensitivity and size of target weeds and environmental conditions at the time of and following application.

Note: Certain varieties of grass may be sensitive to EXPRESS® XP. When using EXPRESS® XP for the first time on a particular variety, limit use to one small container.

**USE RATES AND TANK MIXES WITH OTHER HERBICIDES**

**Bentgrass, Bluegrass, Annual Ryegrass, Orchardgrass, Fine Fescue, Tall Fescue and Timothy**

**Seedling Stands:** For best results apply DuPont™ EXPRESS® XP in a tank mixture with another suitable broadleaf herbicide.

For use on annual ryegrass, orchardgrass, tall fescue and fine fescue, apply at 1/6 oz/A after stand is in 4-leaf stage.

For use on timothy, apply at 1/6 oz/A after stand is in the 4-5 leaf stage. Always use in a tank-mix with 2,4-D at 1/2 lb A/A.

For use on bentgrass, apply at 1/6 oz/A after stolens are 3 to 5 inches across.

For use on bluegrass, apply at 1/6 to 1/3 oz/A after stand is in 4-leaf stage.

**Established Stands:** For stands that have been established for at least one growing season (fall or spring), apply EXPRESS® XP at 1/6 to 1/3 oz/A in a tank mixture with another suitable broadleaf herbicide. Use the higher rate for larger weeds and hard to control weeds like wild carrot. Apply prior to jointing. For application on Timothy, limit maximum use rate to 1/4 oz per acre, and always use in a tank mix with 2,4-D at 1/2 pound active ingredient per acre (1 pint of a 4 pound per gallon product).

**Perennial Ryegrass**

Perennial ryegrass is more sensitive to EXPRESS® XP than other grass species. Crop injury in the form of stunting and possible reduced seed yield may occur. To minimize the risk of crop injury, use the 1/6 oz/A rate and always use either 2,4-D or dicamba and liquid nitrogen with EXPRESS® XP.

**Seedling Stands:** Apply EXPRESS® XP at 1/6 oz/A in a tank mixture with another suitable broadleaf herbicide after grass is in 5- to 6-leaf stage.

**Established Stands:** For stands that have been established for one growing season (fall or spring), apply EXPRESS® XP at 1/6 oz/A to 1/3 oz/A in a tank mixture with another suitable broadleaf herbicide. Apply prior to jointing.

**Note:** The 1/3 oz rate of EXPRESS® XP should be used only for the control or suppression of problem weeds like wild carrot where the benefit of weed control can be offset by possible crop injury including possible yield reduction.

**TANK MIXTURES**

Always use EXPRESS® XP in a tank mixture with another broadleaf herbicide such as 2,4-D, MCP or dicamba as these herbicides safen the effects of EXPRESS® XP on grasses while improving weed control performance on most broadleaf weeds. Testing has shown that 2,4-D and dicamba provide the best overall weed control in a tank mixture with EXPRESS® XP, however 2,4-D at 1/2 pound a.i. per acre provides the best crop safening effects. The addition of liquid fertilizer is also recommended. See "WITH LIQUID NITROGEN SOLUTION FERTILIZER" section of this label.

Use a minimum of 1/4 to 1/2 lb. ai/acre of 2,4-D or MCP (8 to 16 fluid ounces of 4 lb/gal product).

Use a minimum of 1/8 to 1/4 lb ai/acre of dicamba (such as 4 to 8 fluid ounces of "Banvel" or "Clarity").

**Liquid Fertilizer**

EXPRESS® XP can be applied with liquid fertilizers. Liquid fertilizers (20%, 28%, 32% N at a minimum of 4 gallons/100

gallons of spray solution) enhance the performance of EXPRESS® XP and improve crop safety. Always use a surfactant and another broadleaf herbicide when using liquid fertilizer with EXPRESS® XP.

**WEEDS CONTROLLED**

The following weeds are controlled or suppressed in addition to the weeds listed on the EPA registered package label.

- Red deadnettle
- Red sorrel

**WEEDS PARTIALLY CONTROLLED OR SUPPRESSED\***

- Dovefoot geranium
- Redstem filaree
- Spotted catsear
- Wild carrot

\*Partial Control or Suppression: A visual reduction in weed competition (reduced stand and/or vigor) compared to an untreated area.

**Surfactant**

Always use a nonionic surfactant of at least 80% active ingredient at the rate of 0.25% volume/volume (1 quart per 100 gallon of spray solution).

**Precautions/Restrictions**

The use of methylated seed oil (MSO) or crop oil is not recommended with EXPRESS® XP on grass seed crops as these adjuvants may produce unsatisfactory crop injury.

Do not apply more than 1/3 ounce per acre per growing season.

Do not graze or cut for hay or feed associated by-products for 60 days after application. After harvest, straw and other by-products may be fed to animals. make last application of EXPRESS® at least 60 days prior to harvest of grass seed.

Do not apply EXPRESS® XP in a tank mix with organophosphate insecticides as severe crop injury may occur.

Do not apply to grass that is under stress from severe weather conditions, drought, low fertility, water saturated soil, disease or insect damage, as crop injury may result. Under certain conditions such as prolonged cool weather (daily high temperatures less than 50°F) or wide fluctuations in day/night temperatures just prior to or soon after treatment, temporary yellowing and/or crop stunting may occur.

Do not apply to Bermudagrass.

**SPRINKLER CHEMIGATION WITH EXPRESS® XP AND BROMOXYNIL CONTAINING HERBICIDES (SUCH AS "BISON", "BRONATE", "BRONATE ADVANCED" OR "RHINO") IN WINTER & SPRING WHEAT & SPRING BARLEY IN IDAHO**

**DIRECTIONS FOR USE**

EXPRESS® XP Herbicide is recommended in combination with bromoxynil containing herbicides (such as "Bison", "Bronate", "Bronate Advanced" or "Rhino") for use in fall-seeded wheat, spring seeded barley and spring seeded wheat when applied through sprinkler irrigation systems in the state of Idaho.

## HOW TO USE

Use 1/4 to 1/3 oz DuPont™ EXPRESS® XP per acre in combination with bromoxynil containing herbicides at a rate of 3 to 6 oz active ingredient per acre (such as "Bronate" or "Bison" at 3/4 - 1 1/2 pt per acre). Apply to wheat and barley after the 3-leaf stage but before the flag leaf is visible. Make only one chemigation application of this mixture per crop year.

For best results, apply to broadleaf weeds up to the 4-leaf stage, or 2 inches in height or 1 inch in diameter, whichever comes first. Consult this EXPRESS® XP label and bromoxynil containing herbicides package labels for list of weeds controlled or suppressed.

## SPRINKLER IRRIGATION APPLICATION

Apply this tank mix through sprinkler irrigation systems including center pivot, lateral move, side (wheel) roll, solid set or hand move irrigation systems only. Do not apply these herbicides through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. **Do not connect an irrigation system (including greenhouse systems) used for EXPRESS® XP application to any public water system.** A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The sprinkler chemigation system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

## SPECIFIC REQUIREMENTS FOR APPLICATION THROUGH SPRINKLER IRRIGATION SYSTEMS

1. For use via chemigation only in Idaho.
2. In center pivot and continuous lateral move systems, EXPRESS® XP + bromoxynil-containing herbicides should be applied continuously for the duration of the water application. In solid set systems, application of the tank mix should be made during the last 30 to 45 minutes of the irrigation.

3. Set the sprinkler system to deliver approximately 0.5 inch or less of water per acre for best product performance.

4. Fill the supply tank with half of the water amount desired, add the EXPRESS® XP and agitate it well. Add the bromoxynil containing herbicides and then add the remaining water amount with agitation. Bromoxynil containing herbicides require a dilution with at least 4 parts water to 1 part bromoxynil containing herbicides.

5. Agitation is recommended in the pesticide supply tank when applying this tank mixture.

6. The use of a surfactant is not recommended with this tank mix application.

7. Inject the EXPRESS® XP + bromoxynil containing herbicides solution at least 8 feet ahead of a right angle turn of irrigation pipe to insure adequate mixing. Allow sufficient time for the herbicide mixture to be flushed through the lines before turning off irrigation water.

8. Follow both EXPRESS® XP and bromoxynil containing herbicides label instructions for spray tank cleanout both before and after application. Flush lines with clean water following application.

9. Do not apply when wind speed favors drift beyond the area intended for treatment. Avoiding spray drift is the responsibility of the applicator.

## POST HARVEST

### USE RATE

Apply 1/6 to 1/3 oz EXPRESS® XP per acre to crop stubble after harvest. Use 1/3 ounce per acre rate when weed infestation is heavy and predominantly consists of those weeds listed under the "WEEDS PARTIALLY CONTROLLED" section of this label, or when application timing and environmental conditions are marginal. (See the "APPLICATION TIMING" Section for restriction on planting intervals). EXPRESS® XP should be applied in combination with other suitable registered burndown herbicides (See the "TANK MIXTURES" section of this label for additional information).

Sequential treatments of EXPRESS® XP may also be made provided the total amount of EXPRESS® XP applied during one fallow/pre plant cropland season does not exceed 1/3 ounce per acre.

### APPLICATION TIMING

EXPRESS® XP may be used as a burndown treatment to crop stubble when the majority of weeds have emerged and are actively growing.

### TANK MIXTURES

EXPRESS® XP may be used as a post harvest treatment to crop stubble, and should be tank mixed with other herbicides that are registered for use in fallow. Read and follow all manufacturers' label recommendations for the companion herbicide. If those recommendations conflict with this label, do not tank mix the herbicide with EXPRESS® XP.

## GENERAL INFORMATION - ALL USES

### SPRAY ADJUVANTS

Include a spray adjuvant with applications of EXPRESS® XP. In addition, an ammonium nitrogen fertilizer may be used.

Consult your Ag dealer or applicator, local DuPont fact sheets and technical bulletins prior to using an adjuvant system. If another herbicide is tank mixed with DuPont™ EXPRESS® XP, select adjuvants authorized for use with both products. Products must contain only EPA-exempt ingredients (40 CFR 1001).

**Nonionic Surfactant (NIS)**

- Apply 0.06 to 0.50% volume/volume (1/2 pt to 4 pt per 100 gal of spray solution).
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12. – See the Tank Mixtures section of this label for additional information..

**Petroleum Crop Oil Concentrate (COC) or Modified Seed Oil (MSO)**

- Apply at 1% volume/volume (1 gal per 100 gal spray solution) or 2% volume/volume under arid conditions.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

**Special Adjuvant Types**

- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions.
- In addition to the adjuvants specified above, other adjuvant types may be used if they provide the same functionality and have been evaluated and approved by DuPont product management. Consult separate DuPont technical bulletins for detailed information before using adjuvant types not specified on this label.

**Ammonium Nitrogen Fertilizer**

- Use 2 qt/acre of a high-quality urea ammonium nitrate (UAN), such as 28%N or 32%N, or 2 lb/acre of a spray-grade ammonium sulfate (AMS). Use 4 qt/acre UAN or 4 lb/acre AMS under arid conditions.

**GROUND APPLICATION**

For optimum spray distribution and thorough coverage, use flat-fan or low-volume flood nozzles

For flat-fan nozzles, use a spray volume of at least 5 gal per acre (GPA).

For flood nozzles on 30" spacing, use flood nozzles no larger than TK10 (or the equivalent), a pressure of at least 30 psi and a spray volume of at least 10 GPA only. For 40" nozzle spacing, use at least 13 GPA; for 60" spacing use at least 20 GPA. It is essential to overlap the nozzles 100% for all spacings.

"Raindrop RA" nozzles are not recommended for EXPRESS® XP applications, as weed control performance may be reduced.

Use screens that are 50-mesh or larger.

**AERIAL APPLICATION**

Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage at 1 to 5 GPA. Use at least 3 GPA in Idaho, Oregon, or Utah.

Do not apply EXPRESS® XP by air in the state of New York. See the Spray Drift Management section of this label for additional information.

**PRODUCT MEASUREMENT**

EXPRESS® XP can be measured using the EXPRESS® XP volumetric measuring cylinder included in the case. The degree of accuracy of this cylinder varies by ± 7.5%. For more precise measurement, use scales calibrated in ounces.

**CROP ROTATION**

Wheat, Barley and Triticale may be replanted anytime after the application of EXPRESS® XP. Sugarbeets, Winter Rape, and Canola can be planted at 60 days after the application of EXPRESS® XP. Any other crop may be planted 45 days after the application of EXPRESS® XP.

**GRAZING**

Do not graze livestock in treated areas. In addition, do not feed forage or hay from treated areas to livestock (harvested straw may be used for bedding and/or feed).

**MIXING INSTRUCTIONS**

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of EXPRESS® XP.
3. Continue agitation until the EXPRESS® XP is fully dispersed, at least 5 minutes.
4. Once the EXPRESS® XP is fully dispersed, maintain agitation and continue filling tank with water. EXPRESS® XP should be thoroughly mixed with water before adding any other material.
5. As the tank is filling, add tank mixture partners (if desired) then add the required volume of spray adjuvant. Always add spray adjuvant last. Antifoaming agents may be used. Do not use with spray additives that alter the pH of the spray solution below pH 6.0 as rapid product degradation can occur. Spray solutions of pH 7.0 and higher allow for optimum stability of EXPRESS® XP.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Apply EXPRESS® XP spray mixture within 24 hours of mixing to avoid product degradation.
8. If EXPRESS® XP and a tank mixture partner are to be applied in multiple loads, pre-slurry the EXPRESS® XP in clean water prior to adding to the tank. This will prevent the tank mixture partner from interfering with the dissolution of the EXPRESS® XP.

**SPRAY EQUIPMENT**

For specific application equipment, refer to the manufacturer's recommendations for additional information on GPA, pressure, speed, nozzle types and arrangements, nozzle heights above the target canopy, etc.

Be sure to calibrate air or ground equipment properly before application. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern with minimum drift. Use higher spray volumes to obtain better coverage when crop canopy is dense. Avoid swath overlapping, and shut off spray booms while starting, turning, slowing, or stopping, to avoid injury to the crop.

Do not make applications using equipment and/or spray volumes or during weather conditions that might cause spray to drift onto nontarget sites. For additional information on spray drift refer to Spray Drift Management section of label.

Continuous agitation is required to keep DuPont™ EXPRESS® XP in suspension.

## SPRAYER CLEANUP

The spray equipment must be cleaned before EXPRESS® XP is sprayed. Follow the cleanup procedures specified on the labels of the previously applied products. If no directions are provided, follow the six steps outlined in the After Spraying EXPRESS® XP section of this label.

### AT THE END OF THE DAY

When multiple loads of EXPRESS® XP herbicide are applied, it is recommended that at the end of each day of spraying the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits, which can accumulate in the application equipment.

### AFTER SPRAYING EXPRESS® XP AND BEFORE SPRAYING CROPS OTHER THAN WHEAT, BARLEY AND TRITICALE

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of EXPRESS® XP as follows:

1. Drain tank; thoroughly rinse spray tanks, boom, and hoses with clean water. Physically remove any visible deposits.
2. Fill the tank with clean water and 1 gal of household ammonia\* (contains 3% active ingredient) for every 100 gal of water. Flush the hoses, boom, and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 min. Flush the hoses, boom, and nozzles again with the cleaning solution, and then drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat step 2.
5. Rinse the tank, boom, and hoses with clean water.
6. If only ammonia is used as a cleaner, the rinsate solution may be applied back to the crop(s) recommended on this label. Do not exceed the maximum-labeled use rate. If other cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

\* Equivalent amounts of an alternate-strength ammonia solution or a DuPont-approved cleaner can be used in the cleanout procedure. Carefully read and follow the individual cleaner instructions. Consult your Ag dealer, applicator, or DuPont representative for a listing of approved cleaners.

#### Notes:

**PRECAUTION:** Do not use chlorine bleach with ammonia because dangerous gases will form.

Do not clean equipment in an enclosed area.

1. Steam-cleaning aerial spray tanks is recommended prior to performing the above cleanout procedure to facilitate the removal of any caked deposits.
2. When EXPRESS® XP is tank mixed with other pesticides, cleanout procedures for each product should be examined and the most rigorous procedure should be followed.
3. In addition to this cleanout procedure, all precleanout guidelines on subsequently applied products should be followed

as per the individual product labels.

4. Where routine spraying practices include shared equipment frequently being switched between applications of EXPRESS® XP and applications of other pesticides to EXPRESS® XP sensitive crops during the same spray season, it is recommended that a sprayer be dedicated to EXPRESS® XP to further reduce the chance of crop injury.

## SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

**AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.**

### IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150 - 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **Wind, Temperature and Humidity, and Surface Temperature Inversions** sections of this label.

### Controlling Droplet Size - General Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

### Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.

### BOOM LENGTH AND HEIGHT

- **Boom Length (aircraft)** - The boom length should not exceed 3/4 of the wing length. Using shorter booms decreases drift potential. For helicopters use a boom length and position that prevents droplets from entering the rotor vortices.

13/19

- **Boom Height (aircraft)** - Application more than 10 ft above the canopy increases the potential for spray drift.
- **Boom Height (ground)** Setting the boom at the lowest height which provides uniform coverage reduces the exposure of droplets to evaporation and wind. The boom should remain level with the crop and have minimal bounce.

**WIND**

Drift potential increases at wind speeds of less than 3 mph (due to variable direction and inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. AVOID APPLICATIONS DURING GUSTY OR WINDLESS CONDITIONS.

**Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they effect spray drift.

**TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

**SURFACE TEMPERATURE INVERSIONS**

Drift potential is high during a surface temperature inversion. Surface inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Surface inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates a surface inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

**RESISTANCE**

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will

promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

**INTEGRATED PEST MANAGEMENT**

DuPont recommends the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an Integrated Pest Management (IPM) program which can include biological, cultural, and genetic practices aimed at preventing economic pest damage. Application of this product should be based on IPM principles and practices including field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop or site systems in your area.

**PRECAUTIONS**

- Do not graze treated fields or feed treated forage or hay (harvested straw may be used for bedding and/or feed).
- Varieties of wheat (including durum), barley and triticale may differ in their response to various herbicides. DuPont recommends that you first consult your state experiment station, university, or extension agent as to crop sensitivity to any herbicide. If no information is available, limit the initial use to a small area.
- Under certain conditions such as heavy rainfall, prolonged cold weather, or wide fluctuations in day/night temperatures prior to or soon after DuPont™ EXPRESS® XP application, temporary discoloration and/or crop injury may occur. To reduce the potential of crop injury, tank mix EXPRESS® XP with 2,4-D (ester formulations perform best— see the "TANK MIXTURES" section of this label) and apply after the crop is in the tillering stage of growth.
- EXPRESS® XP should not be applied to wheat, barley or triticale that is stressed by severe weather conditions, drought, low fertility, water-saturated soil, disease, or insect damage, as crop injury may result. Risk of injury is greatest when crop is in the 2 to 5- leaf stage. Severe winter stress, drought, disease, or insect damage following application also may result in crop injury.
- Do not apply to wheat, barley or triticale underseeded with another crop.
- Dry, dusty field conditions may result in reduced control in wheel track areas.
- Injury to or loss of desirable trees or vegetation may result from failure to observe the following:
  - Do not apply, drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
  - Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants.
- Injury to or loss of adjacent sensitive crops and vegetation may

15/19

result from failure to observe the following:

- Take all necessary precautions to avoid all direct or indirect contact (such as spray drift) with non-target plants or areas.
- Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than wheat or barley.

**PESTICIDE STORAGE AND DISPOSAL**

**PESTICIDE STORAGE:** Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage.

**PRODUCT DISPOSAL:** Do not contaminate water, food or feed by disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL: For Plastic Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **For Fiber Sacks:** Completely empty fiber sack by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into manufacturing or application equipment. Then dispose of sack in a sanitary landfill or by incineration if allowed by State and local authorities. **For Fiber Drums With Liners:** Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner. **For Bags Containing Water Soluble Packets:** Do not reuse the outer box or the resealable plastic bag. When all water-soluble packets are used, the outer packaging should be clean and may be disposed of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by open burning. If burned, stay out of smoke. If the resealable plastic bag contacts the formulated product in any way, the bag must be triple-rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer wrap as described above. **For Metal Containers (non aerosol):** Triple rinse (or equivalent) the container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. **For Paper and Plastic Bags:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

When packaged in returnable and/or refillable containers:

**Container Refilling and Disposal (For Containers up to 250 gal) -** This is a refillable container. If the container is to be refilled, do not rinse with any material or introduce any pesticide other than DuPont™ EXPRESS® XP herbicide. Reseal and return the container to any authorized DuPont refilling facility. If the container is not to be refilled, triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-441-3637 day or night.

**Container Disposal for Bulk Containers:** When this container is empty, replace the cap and seal all openings that have been opened during use; and return the container to the point of purchase or to a designated location named at time of purchase of this product. The container must only be refilled with this pesticide product. **DO NOT REUSE THE CONTAINER FOR ANY OTHER PURPOSE.** Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transporting. Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, contact DuPont at 1-800-441-3637. If not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling. Disposal of this container must be in compliance with state and local regulations.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-441-3637, day or night.

**NOTICE TO BUYER:** Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval Logo, DuPont™, and EXPRESS® are trademarks or registered trademarks of E. I. duPont de Nemours & Company

16/19

"Maverick", "Landmaster II", "Fallow Master", "RT Master" and "Roundup" are registered trademarks of Monsanto

"Pursuit", "Raptor", "Beyond" and "Clarity" are registered trademarks of BASF Corporation.

"Puma", "Buctril", "Bronate", "Bronate Advanced" and "Hoelon 3EC" are registered trademarks of Bayer Corporation.

"Raindrop RA" is a registered trademark of Delavan

"Everest" is a registered trademark of Arvesta Corporation

"Aim" is a registered trademark of FMC Corporation

"Discover" is a registered trademark of Syngenta Corporation

"Stinger", "Curtail", "Curtail M", "Lorsban", "Starane + Salvo", "Starane + Sword" and "Starane", "WideMatch" are registered trademarks of Dow AgroSciences

"Malathion" is registered trademark of UAP Loveland Products, Inc.

"Bison" is a registered trademark of Agriliance, LLC.

"Banvel" and "Assert" are registered trademarks of MicroFlo Company

Net Wt. 10 oz

**D- 1077 120905**

### **LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.**

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants.

DuPont does not agree to be an insurer of these risks. **TO THE FULLEST EXTENT PERMITTED BY LAW, WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

**DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE FULLEST EXTENT PERMITTED BY LAW, IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

**For product information call: 1-888-6-DUPONT**

**Internet address: <http://cropprotection.dupont.com/>**

**© 2000-2005 E. I. du Pont de Nemours and Company, Wilmington, Delaware 19898.**

**All rights reserved.**



17/19

# SUPPLEMENTAL LABELING

**DuPont Crop  
Protection**

**DUPONT™ EXPRESS®  
HERBICIDE  
EXPRESS® HERBICIDE  
FOR USE ONLY ON  
CDC TRIFFID FLAX**

## **DUPONT™ EXPRESS® XP HERBICIDE EXPRESS® HERBICIDE**

EPA REG. NO. 352-509

### **FOR USE ONLY ON CDC TRIFFID FLAX IN THE STATES OF MN, ND AND MT**

EXPRESS® Herbicide or DuPont™ EXPRESS® XP Herbicide (EXPRESS®) can be applied on flax that contains the "Freedom™" trait (CDC Triffid Flax).

• Applications to CDC Triffid Flax that is or has been stressed prior to application by severe weather conditions, frost, low fertility, drought, water saturated soil, disease or insect damage, may result in crop injury. Applications to CDC Triffid Flax if the above stress conditions are expected within 3 days after application may also result in crop injury.

#### **DIRECTIONS FOR USE**

Apply EXPRESS® at 1/10 ounce per acre. Add a DuPont recommended, nonionic surfactant having at least 80% active ingredient strength at 0.25% v/v (1 qt per 100 gal of spray solution).

Apply EXPRESS® from the time the CDC Triffid Flax is 1/2 inch in crop height to the pre-bud stage. Temporary crop yellowing may be observed shortly after application of EXPRESS® Herbicide, especially when applied to crops growing under environmentally stressful conditions (See PRECAUTIONS).

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### **PRECAUTIONS**

- Application of EXPRESS® prior to CDC Triffid Flax crop height reaching 1/2 inch tall could result in crop injury.
- Avoid application to CDC Triffid Flax fields in which germination is uneven (some plants outside the recommended leaf stage for application), as crop injury may result.
- Under certain environmentally stressful conditions (such as heavy rainfall, prolonged cool or hot weather, frost conditions, wide fluctuations in day/night temperatures), temporary lightening in crop color and occasionally a slight reduction in crop height may occur.

#### **RESTRICTIONS**

- Only apply to flax that contains the "Freedom™" trait. EXPRESS® will severely damage flax varieties that do not contain the "Freedom™" trait.

#### **IMPORTANT**

#### **BEFORE USING EXPRESS®, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.**

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

R-405 081004 11-18-04

For product information call 1-888-6-DUPONT  
© 2004 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898

18/19

# SUPPLEMENTAL LABELING

DUPONT™ EXPRESS® XP

HERBICIDE

EXPRESS® HERBICIDE

FOR USE ONLY ON IMAZETHAPYR  
TOLERANT CANOLA

**DuPont Crop  
Protection**

## DUPONT™ EXPRESS® XP HERBICIDE EXPRESS® HERBICIDE

EPA REG. NO. 352-509

**FOR USE ONLY ON IMAZETHAPYR TOLERANT CANOLA  
(CANOLA VARIETIES WITH THE SMART™ TRAIT)  
FOR USE IN THE STATES OF MN, ND AND MT**

EXPRESS® Herbicide or DuPont™ EXPRESS® XP Herbicide (EXPRESS®) can be applied on Imazethapyr Tolerant Canola (eg. canola varieties with the SMART™ trait).

### DIRECTIONS FOR USE

Apply EXPRESS® at 1/10 ounce per acre. Add a DuPont recommended, nonionic surfactant having at least 80% active ingredient strength at 0.25% v/v (1 qt per 100 gal of spray solution).

Apply EXPRESS® when the Imazethapyr Tolerant Canola is in the 3 to 6 leaf stage of growth but prior to beginning of bolting. Temporary crop yellowing may be observed shortly after application of EXPRESS®, especially when applied to crops growing under environmentally stressful conditions (See PRECAUTIONS).

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### TANK MIXTURES

For the control of annual grasses as well as broadleaf weeds, apply a tank mixture of EXPRESS® at 1/10 ounce per acre plus ASSURE® II Herbicide (See the EPA approved ASSURE II label for use rates, weed size, precautions, and restrictions). Add a DuPont recommended, nonionic surfactant having at least 80% active ingredient strength at 0.25% v/v (1 qt per 100 gal of spray solution).

### PRECAUTIONS

- Application of EXPRESS® prior to the 3 leaf stage of Imazethapyr Tolerant Canola could result in crop injury.
- Avoid application to Imazethapyr Tolerant Canola fields in which germination is uneven (some plants outside the recommended leaf stage for application), as crop injury may result.
- Under certain environmentally stressful conditions (such as heavy rainfall, prolonged cool or hot weather, frost conditions, wide fluctuations in day/night temperatures), temporary lightening in crop color and occasionally a slight reduction in crop height may occur.
- Applications to Imazethapyr Tolerant Canola that are or have been stressed prior to application by severe weather conditions, frost, low fertility, drought, water saturated soil, disease or insect damage, may result in crop injury. Applications to Imazethapyr Tolerant Canola if the above stress conditions are expected within 3 days after application, may also result in crop injury.
- Application of the EXPRESS® or EXPRESS® plus ASSURE II Herbicide tank mixture on Imazethapyr Tolerant Canola may delay flowering.

### RESTRICTIONS

- DO NOT apply to NON-Imazethapyr Tolerant Canola (i.e. Canola varieties that DO NOT contain the SMART™ trait) as severe crop injury or death of the plant may occur.

For product information call 1-888-6-DUPONT

© 2004 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898

17/19

**IMPORTANT**

**BEFORE USING EXPRESS® OR ASSURE II HERBICIDE, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.**

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

R-406 081004 11-18-04

For product information call 1-888-6-DUPONT

© 2004 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898