PM 25 350-500 PRUPUSED

11-3-98

SUPPLEMENTAL LABELING

GLEAN® FC HERBICIDE

ROTATION INTERVAL TO STS™ SOYBEAN

"..... A Growing Partnership With Nature"

Products

DuPont Agricultural

GLEAN® FC HERBICIDE

EPA REG. NO. 352-522

ROTATION INTERVAL TO STS® SOYBEAN®

ACCEPTED

NOV - 3 1998

Under the Federal Insections, Fungloide, and Roderticide Act as amended, for the posteride registered under BPA Reg. No 75252

DIRECTIONS FOR USE

This product is a water dispersible granule containing 75% active ingredient by weight.

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

CROP ROTATION

Minimum Rotation Intervals

Minimum rotation intervals* are determined by the rate of breakdown of GLEAN FC applied. GLEAN FC breakdown in the soil is affected by soil pH, soil temperature, soil microorganisms, and soil moisture. Low soil pH, high soil temperature, and high soil moisture increase GLEAN FC breakdown in soil, while high soil pH, low soil temperature, and low soil moisture slow GLEAN FC breakdown.

Of these three factors, only soil pH remains relatively constant. Soil temperature, and to a greater extent, soil moisture, can vary significantly from year to year and from area to area. For this reason, soil temperatures and soil moisture should be monitored regularly when considering rotating to other crops.

Soil pH Limitations

GLEAN FC should not be used on fields having a soil pH above 7.9, as extended soil residual activity could extend crop rotation intervals beyond those specified in the rotation table, and under certain conditions, could injure wheat or barley. In addition, other crops planted in high-pH soils can be extremely sensitive to low concentrations of GLEAN FC.

Before using GLEAN FC, determine the soil pH of the field. To obtain a representative pH value, take several samples from different areas of the field between 0" and 4" deep and analyze them separately. Consult local extension publications for addi-

tional information on recommended soil sampling procedures. Before using GLEAN FC, carefully consider your crop rotation plans and options. For rotational flexibility, do not treat all of your wheat, barley, or fallow acres at the same time.

Rotation Intervals for STS Soybeans - Irrigated/Non Irrigated land following wheat, barley or fallow land at the Maximum Use Rates listed in the following table.

			Maximum Sammum	
Areas	Crop	Soil pH	Use Rate	Retation lawarest
			(02/3CIC)	t=onths?
All Areas of	STS	7.5 or	1/2	6
AR, MD, VA, NC	Soybeans	, lower		
SC and GA				

*The minimum rotation interval represents the period of time from the last application to the anticipated date of the next planting. Rotation intervals are based on normal precipitation/irrigation amounts. See EPA approved GLEAN FC label for additional details on crop rotation recommendations and restrictions.

IMPORTANT PRECAUTIONS

- Injury to or loss of desirable trees or vegetation may result from failure to observe the following:
 - Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
 - Do not use on lawns, walks, driveways, tennis courts or similar areas.
 - Prevent drift of spray to desirable plants.
- Do not contaminate any body of water. Thoroughly clean application equipment immediately after use
- GLEAN FC is non-corrosive, non-flammable, non-volatile, and does not freeze in storage.

© 1998 E. I. du Pont de Nemours and Company, DuPont Agricultural Products, Wilmington, Delaware 19898

Safa

- Under certain conditions (such as drought, prolonged cold weather, pH variability in the fields) temporary discoloration and/or crop injury may occur to STS soybeans planted after GLEAN FC applications.
- This supplemental tabel does not apply to crops grown for seed.

IMPORTANT

BEFORE USING GLEAN FC, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

¹ Sulfonylurea Tolerant Soybeans

090398