

**SUPPLEMENTAL LABELING****DUPONT™ GLEAN® FERTILIZER
COMPATIBLE HERBICIDE****TANK MIX WITH EVEREST HERBICIDE
FOR WEED CONTROL IN WHEAT****DuPont Crop Protection****DUPONT™ GLEAN® FERTILIZER COMPATIBLE HERBICIDE
EPA Reg. No. 352-522****TANK MIX WITH EVEREST HERBICIDE
FOR WEED CONTROL IN WHEAT****GENERAL INFORMATION**

DUPONT™ GLEAN® FERTILIZER COMPATIBLE HERBICIDE (GLEAN® FC) is a dry flowable formulation that selectively controls certain broadleaf weeds in wheat. The degree and duration of weed control depends on the weed spectrum and infestation intensity, the weed size at the time of application, and/or the environmental conditions at and following treatment.

GLEAN® FC is noncorrosive to equipment, nonflammable, nonvolatile, and does not freeze.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GLEAN® FC may be tank mixed with EVEREST Herbicide for improved control of weeds in wheat. Read and follow all label instructions regarding timing of application, precautions, and warnings for these herbicides before using the tank mixture. Follow the most restrictive labeling.

POSTEMERGENCE APPLICATION

Winter Wheat: Apply in the fall or spring anytime after the crop has 2 total leaves on main stem and any number of tillers. Apply before jointing begins.

Treat late-seeded winter wheat after the crop has started to tiller since the combined effect of herbicide stress and stress from cold weather and/or moisture could cause crop injury.

Spring Wheat: Apply anytime after emergence, but before the majority of plants have 4 total leaves on the main stem plus 2 tillers. Do not apply after jointing begins. Do not apply to durum wheat.

The addition of 2,4-D or Dicamba to the GLEAN® FC + EVEREST tank-mix is required when applying to spring wheat. The 2,4-D and Dicamba rates are as follows:

- 2,4-D Amine or LV Ester (4 lbs./gallon): 0.25 – 0.75 pt./acre
- 2,4-D LV Ester (6 lbs./gallon): 0.17 – 0.5 pt./acre
- Dicamba (4 lbs./gallon): 2 – 4 fl. oz. per acre

Note: When GLEAN® FC + EVEREST is applied in a tank-mix with a dicamba-containing herbicide, wild oat control may be reduced.

ADDITIONAL TANK MIX INFORMATION AND PRECAUTIONS

Tank Mixes with Insecticides: Under certain conditions (stress from drought, cold weather or warm days and cold nights following application, or crops in the 2-4 leaf stage), tank mixtures or sequential treatments of GLEAN® FC + EVEREST and organophosphate insecticides (such as methyl or ethyl parathion, "Di-Syston", etc.) may produce temporary crop yellowing or, in severe cases, crop injury. Test these mixtures in a small area first. If no symptoms of crop injury occur 14 days after treatment, treat the rest of the acreage.

Do not use GLEAN® FC + EVEREST plus Malathion, as crop injury may result.

Do not apply GLEAN® FC + EVEREST within 60 days of crop emergence where an organophosphate insecticide (such as "Di-Syston") has been applied as an in-furrow treatment, as crop injury may result.

SURFACTANTS

Winter Wheat

Non-ionic Surfactant (NIS): Unless otherwise specified, add a non-ionic surfactant having at least 80% active ingredient at 0.5% v/v (2 quarts per 100 gallons of spray solution). Antifoaming agents may be used if needed.

Ammonium Nitrogen Fertilizer: In addition to a non-ionic surfactant, use 2 qt./acre of a high-quality urea ammonium nitrate (UAN), such as 28%N or 32%N, or 2 lbs./acre of a spray-grade ammonium sulfate (AMS). Use 4 qt/acre UAN or 4 lbs./acre AMS under arid conditions.

Do not use low rates of liquid fertilizer as a substitute for surfactant.

Carrier Solutions (With Liquid Nitrogen Solution Fertilizer): Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing GLEAN® FC and EVEREST in fertilizer solution. GLEAN® FC and EVEREST must first be slurried with water, then added to liquid nitrogen solutions (e.g., 28-0-0, 32-0-0). Ensure that the agitator is running while GLEAN® FC and EVEREST are added. Note that use of fertilizer as the carrier may result in temporary crop yellowing and stunting.

If using rates of liquid nitrogen fertilizer in the spray solution that are less than 50% of the spray solution volume, the addition of non-ionic surfactant is necessary. Add surfactant at 0.25% v/v (1 quart per 100 gallons of spray solution) based on local recommendations.

When using rates of liquid nitrogen fertilizer in the spray solution that are greater than 50% of the spray solution, adding surfactant increases the risk of crop injury. Consult your agricultural dealer, consultant, fieldsman, or DuPont representative for a specific recommendation before adding surfactant to these tank mixtures.

Spring Wheat

Non-ionic Surfactant (NIS): Unless otherwise specified, add a non-ionic surfactant having at least 80% active ingredient at 0.125% v/v (1 pint per 100 gallons of spray solution). Antifoaming agents may be used if needed.

DO NOT add NIS if also tank-mixing with 2,4-D LV Ester.

RATE OF APPLICATION AND WEEDS CONTROLLED OR PARTIALLY CONTROLLED

A tank mixture of GLEAN® FC + EVEREST will provide control, partial control, and/or suppression of the following weeds when applied at the specified rates and application timing.

GRASS WEED RECOMMENDATIONS

GRASS WEEDS	EARLY* APPLICATION of 0.6 oz EVEREST + 0.3 oz. GLEAN® FC per acre	EARLY APPLICATION of 0.4 oz. EVEREST + 0.2 oz. GLEAN® FC per acre	LATE* APPLICATION of 0.6 oz EVEREST + 0.3 oz. GLEAN® FC per acre	LATE APPLICATION of 0.4 oz. EVEREST + 0.2 oz. GLEAN® FC per acre
Annual Ryegrass ***	C **	PC	C	PC
Downy Brome (<i>Bromus tectorum</i>)	PC	NR	PC	NR
Cheat (<i>Bromus secalinus</i>)	C	C #	C	C #
Japanese Brome (<i>Bromus japonicus</i>)	C	C #	C	C #
Wild Oat	C	C	C	C

*EARLY APPLICATION: Timing - 2 leaf stage to January 1
LATE APPLICATION: Timing - after January 1 but prior to joint

** C = Control
PC = Partial Control (partially controlled weeds exhibit a visual reduction in numbers and/or a significant loss of vigor)
NR = Not recommended

*** If GLEAN® FC + EVEREST is applied in a tank-mix combination with a 2,4-D containing broadleaf herbicide, Ryegrass control may be reduced

For control of these weed species at the reduced rate, liquid nitrogen fertilizer as a carrier solution of 50% or greater must be added to the tank mix

BROADLEAF WEED RECOMMENDATIONS

BROADLEAF WEEDS	EARLY* APPLICATION of 0.6 oz EVEREST + 0.3 oz. GLEAN® FC per acre	EARLY APPLICATION of 0.4 oz. EVEREST + 0.2 oz. GLEAN® FC per acre	LATE* APPLICATION of 0.6 oz EVEREST + 0.3 oz. GLEAN® FC per acre	LATE APPLICATION of 0.4 oz. EVEREST + 0.2 oz. GLEAN® FC per acre
Bedstraw	PC**	NR	PC	NR
Black Mustard	C	C	C	C
Blue Mustard	C	C	C	C
Bur beakchervil	C	NR	C	NR
Bushy Wallflower / Treacle Mustard	C	C	C	C
Buttercup	C	NR	C	NR
Canada thistle	PC	NR	PC	NR
Coast Fiddleneck (tarweed)	C	NR	C	NR
Common chickweed	C	NR	C	NR
Common groundsel	C	NR	C	NR
Conical catchfly	C	C	C	C
Corn gromwell	PC	NR	PC	NR
Curly dock	C	C	C	C
Corn spurry	C	NR	C	NR
Cow cockle	C	NR	C	NR
Cutleaf evening primrose	C	C	C	C
False chamomile	C	NR	C	NR
Falseflax	C	NR	C	NR
Field pennycress	C	C	C	C
Flixweed	C	C	C	C
Hempnettle	C	C	C	C
Henbit	C	C	C	C
Kochia *	PC	NR	PC	NR
Ladysthumb	C	NR	C	NR
Lambsquarter	C	NR	C	NR
Mayweed	C	C	C	C
Miners lettuce	C	C	C	C
Mouseear chickweed	C	NR	C	NR
Pennsylvania smartweed	PC	NR	PC	NR
Pineappleweed	C	C	C	C
Pigweeds (redroot, smooth, prostrate, tumble)	C	C	C	C
Prickly lettuce *	PC	NR	PC	NR
Prostrate knotweed	PC	NR	PC	NR
Purslane (common)	C	NR	C	NR

6/7

BROADLEAF WEEDS	EARLY* APPLICATION of 0.6 oz EVEREST + 0.3 oz. GLEAN® FC per acre	EARLY APPLICATION of 0.4 oz. EVEREST + 0.2 oz. GLEAN® FC per acre	LATE* APPLICATION of 0.6 oz EVEREST + 0.3 oz. GLEAN® FC per acre	LATE APPLICATION of 0.4 oz. EVEREST + 0.2 oz. GLEAN® FC per acre
Redstem filaree	C	NR	C	NR
Russian thistle +	PC	NR	PC	NR
Shepherd's-purse	C**	C	C	C
Speedwell	PC	NR	PC	NR
Sunflower	PC	NR	PC	NR
Tansymustard	C	C	C	C
Volunteer canola	C	NR	C	PC
Waterpod	C	C	C	C
White cockle	C	NR	C	NR
Wild buckwheat	PC	NR	PC	NR
Wild carrot	C	NR	C	NR
Wild garlic/wild onion	PC	NR	PC	NR
Wild mustard	C	C	C	C
Wild radish	PC	NR	PC	NR
Wild turnip	C	NR	C	NR

***EARLY APPLICATION:** Timing - 2 leaf stage to January 1
LATE APPLICATION: Timing - after January 1 but prior to joint

** C = Control
 PC = Partial Control (partially controlled weeds exhibit a visual reduction in numbers and/or a significant loss of vigor)
 NR = Not recommended

+ Naturally occurring resistant biotypes of these weeds are known to occur. See the Tank Mixtures and Resistance sections on the GLEAN® FC label for additional information.

Refer to the complete GLEAN® FC and EVEREST labels for additional information regarding use restrictions, tank-mix partners, rotational cropping recommendations, sprayer cleanup, use precautions, and other information. The most restrictive provisions on either label will apply. Do not use this tank mix if any restrictions on the EVEREST label conflict with recommendations on the GLEAN® FC label.

IMPORTANT

**BEFORE USING GLEAN® FC HERBICIDE AND/OR EVEREST HERBICIDE
READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS,
AND PRECAUTIONS ON THE EPA-REGISTERED LABELS.**

This labeling must be in the possession of the user at the time of pesticide application.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

EVEREST is a registered trademark of Arvesta Corporation.