

APPLICATION INFORMATION

"Assure" tank mixes may be applied by ground or by air. Use a minimum of 10 gallons of water per acre by ground or a minimum of 5 gallons of water per acre by air.

Application of "Assure" tank mixes must include an adjuvant. The following adjuvants are recommended:

For tank mixes of "Assure" plus "Classic" use a non-ionic surfactant at a rate of 0.25% v/v (1 qt/100 gal) by ground or air. OR Use a crop oil concentrate at a rate of 1% v/v (1 gal/100 gal) by ground or 0.5% v/v (2 qt/100 gal) by air.

For tank mixes of "Assure" plus "Pinnacle" or "Assure" plus "Pinnacle" plus "Classic" use a non-ionic surfactant at a rate of 0.125 % v/v (1 pt/100 gal) by ground or air. Under dry conditions a non-ionic surfactant at a rate of 0.25% v/v (1 qt/100 gal) or a crop oil concentrate at a rate of 0.5% v/v (2 qt/100 gal) may be used. Use of the higher rate of non-ionic surfactant or crop oil concentrate in a tank mix containing "Pinnacle" may increase crop injury.

Do not use vegetable (crop origin) oils as a spray adjuvant or carrier.

Note: Tank mixes of "Assure" with post broadleaf herbicides have typically shown some reduction in grass control over applications of "Assure" alone. Grasses other than the ones listed above may not be adequately controlled by the rates of "Assure" listed. Refer to the "Assure" label for specific use directions for other grasses.

IMPORTANT

BEFORE USING "ASSURE", "CLASSIC", OR "PINNACLE", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABELS. WHEN USING A TANK MIX THE MOST RESTRICTIVE LABELING WILL APPLY.

This bulletin contains new or supplemental instructions for use of these products which may not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of application.

[1] Registered trademark of E. I. du Pont de Nemours and Company

E. I. du Pont de Nemours and Company
Agricultural Products Department
Wilmington, DE 19880-0038

032390

212

**"PINNACLE" (1)
HERBICIDE TANK MIX FOR
CONTROL OF CERTAIN
WEEDS IN SOYBEANS**

**"PINNACLE" HERBICIDE
EPA REG. NO. 352-525**

**SUPPLEMENTAL LABELING
"PINNACLE" HERBICIDE
TANK MIX WITH "BASAGRAN" (2) HERBICIDE
FOR CONTROL OF CERTAIN WEEDS IN SOYBEANS**

DIRECTIONS FOR USE

It is a violation of federal law to use this product
in a manner inconsistent with its labeling.


Du Pont "Pinnacle" Herbicide is recommended for control
of the weeds listed below when applied to soybeans
in a tank mix with "Basagran" Herbicide.

Do not apply this tank mix through any type of
irrigation system.

HOW TO USE

A tank mix of "Pinnacle" Herbicide at 1/4 oz /A plus "Basagran" Herbicide
at 1 1/2 pt./A is recommended for control of the listed weeds.
Application should be made when the weeds are young, actively growing,
and prior to exceeding the maximum size listed in the table below.

<u>WEED</u>	<u>MAXIMUM SIZE</u>
Annual Smartweeds	up to 6"
Cocklebur	up to 6"
Common Lambsquarters	up to 4"
Jimsonweed	up to 6"
Pigweed spp.	up to 8"
Prickly Sida	up to 3"
Velvetleaf	up to 5"
Venice Mallow	up to 2"
Wild Buckwheat	up to 3"
Wild Mustard	up to 4"
Wild Sunflower	up to 5"



Include a nonionic surfactant of at least 80% active ingredient at a rate
(concentration) of 0.125 % v/v (1 pt./100 gal. spray solution) and a
liquid nitrogen fertilizer (such as 28-0-0) at a rate (concentration) of
4% v/v (4 gal./100 gal. spray solution).

Under dry conditions a nonionic surfactant at a rate of 0.25% v/v (2 pt./100 gal.) or a crop oil concentrate at a rate of 0.5% v/v (4 pt./100 gal.) may be used to increase weed control. Use of crop oil concentrate or the higher rate of surfactant (0.25% v/v) may increase injury to soybeans.

For ground application use a minimum of 20 gallons of water per acre. When using flat fan nozzles use 40 psi. When using hollow cone nozzles use 40-60 psi. Do not use flood, whirl chamber, or controlled droplet application (CDA) nozzles.

For aerial application use a minimum of 5 gallons of water per acre. Consult the respective labels for special directions for aerial applications.

Apply after the first trifoliolate of the soybean plant has fully expanded and up to 60 days before soybean harvest.

Refer to the "Pinnacle" Herbicide and "Basagran" Herbicide labels for additional use directions, use restrictions, and precautions. The most restrictive provision on either label will apply.

IMPORTANT

BEFORE USING "PINNACLE" HERBICIDE AND "BASAGRAN" HERBICIDE READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABELS.

This bulletin contains new or supplemental use for these products which may not appear on the package labels. Follow the instructions carefully.

This labeling must be in possession of the user at the time of application.

- [1] Registered trademark of E. I. du Pont de Nemours & Co. (Inc.)
- [2] Registered trademark of BASF AG

E. I. du Pont de Nemours & Co. (Inc.)
Agricultural Products Department
Wilmington, DE 19880-0038

022690