

PINNACLE® HERBICIDE

EPA Reg. No. 352-525

CLASSIC® HERBICIDE

EPA Reg. No. 352-436

**"PINNACLE" HERBICIDE PLUS "CLASSIC" HERBICIDE FOR WEED CONTROL IN SOYBEANS IN THE
STATES OF KANSAS, MINNESOTA (SOUTH OF HIGHWAY 27), MISSOURI (EXCEPT THE BOOTHEEL AREA),
NEBRASKA AND SOUTH DAKOTA**

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this tank mix through any type of irrigation system.

HOW TO USE

A tank mix of "Pinnacle" Herbicide at a rate of 1/4 ounce per acre plus "Classic" Herbicide at a rate of 1/3 ounce per acre is recommended for control of the weeds listed below. Application should be made when weeds are young, actively growing and prior to exceeding the maximum size listed in the table. Applications made to weeds in the cotyledon stage or to weeds exceeding the maximum size listed below may result in unsatisfactory control.

WEEDS CONTROLLED**SIZE RANGE**

Annual Smartweeds	2" - 5"
Cocklebur	2" - 6"
Jimsonweed	2" - 5"
Lambsquarters	2" - 4"
Ragweed Species	2" - 12"
Velvetleaf	2" - 8"
Wild Mustard	2" - 4" (diameter)
Wild Sunflower	2" - 8"

WEEDS SUPPRESSED**SIZE RANGE**

Morningglory Species	
Entire leaf	1" - 2"
Ivy leaf	1" - 2"
Pinna	1" - 2"
Small flower	1" - 2"
Tall	1" - 2"
Common Ragweed	1" - 3"

*Requires the addition of ammonium nitrogen fertilizer.

Applications must be made with a surfactant at a rate not in excess of 0.125 ounce per acre per application. Do not use crop oil concentrate.

The addition of ammonium nitrogen fertilizer is required for control of velvetleaf. Use a high quality liquid nitrogen fertilizer such as 28-0-0 at a rate of 2 quarts per acre (use the lower rate for spray volumes less than 15 gallons per acre) or 10-34-0 at a rate of 1 quart per acre. Alternatively, a high quality, sprayable grade of ammonium sulfate (21-0-0) may be used at a rate 2 pounds per acre. The addition of ammonium nitrogen fertilizer does not replace the need for a surfactant.

For broadcast applications by ground equipment, use flat fan nozzles at 25 - 40 PSI. Use 10 - 25 gallons per acre. Do not use hollow cone, flood, rain drop, whirl chamber nozzles. For proper spray coverage, adjust the boom and nozzle height according to the specifications listed by the manufacturer.

For aerial applications, use nozzle types and arrangements that will provide optimum spray distribution and maximum coverage at 5 to 10 gallons per acre.

Applications of "Classic" at 1/3 oz/A are not limited by soil pH. However, the following crop rotation intervals must be observed:

**ROTATION INTERVAL (in months)
FROM "CLASSIC" APPLICATION**

CROP	MI, NE, SD	KS, MO
Soybeans	Anytime	Anytime
Cereal Grains	3	3
Field Corn*	9	9
Clover	9	9
Dry Beans	9	9
Tomato (Transplant)	9	9
Alfalfa	15	9
Sorghum	15	9
Tobacco (Transplant)	15	9

All other crops not listed above require completion of a successful field bioassay. A field bioassay will detect small quantities of "Classic" which can remain in the soil and injure rotational crops. A successful field bioassay means growing to maturity a test strip of crop(s) intended for production the following year. The test strip should cross the whole field including knolls and low areas. Suggested interval before planting bioassay crop is 9 months.

*When using "Classic" Herbicide the term "FIELD CORN" is defined to include only that corn grown for grain or silage, popcorn, and seed corn relative to rotational crop intervals.

© 1991 E.I. DU PONT DE NEMOURS AND COMPANY, AGRICULTURAL PRODUCTS, WILMINGTON, DELAWARE

Do not tank mix with any other pesticide, such as post grass herbicides, as excessive injury may occur.

"Pinnacle" plus "Classic" may be applied anytime after the first trifoliate leaf has fully expanded until 60 days prior to harvest.

Applications during excessively hot and humid weather should be avoided as excessive crop injury may occur.

Refer to the "Pinnacle" and "Classic" labels for additional use directions, use restrictions, and precautions.

may not appear on the package label. Follow the instructions carefully.
This labeling must be in the possession of the user at the time of application.

D-013191

