



PM 25

352-525

3/27/98
SUPPLEMENTAL LABELING 1/2**DuPont Agricultural
Products**

"..... A Growing Partnership With Nature"

**PINNACLE® HERBICIDE
FOR USE IN FIELD CORN****PINNACLE® HERBICIDE**

EPA Reg. No. 352-525

**FOR POST EMERGENCE CONTROL OF CERTAIN BROADLEAF WEEDS IN FIELD CORN FOR
USE ONLY IN THE STATES OF CT, DE, MA, MD, ME, NH, NC, NJ, PA, RI, VA, VT, WV****DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont "Pinnacle" Herbicide is recommended for postemergence control of certain broadleaf weeds in field corn.

Do not apply this product through any kind of irrigation system.

Do not graze or feed forage or grain from treated field corn to livestock within 30 days of application.

Do not apply to fields treated with Counter¹ insecticide applied at planting or over-the-row at cultivation as severe crop injury may result.

APPLICATION INFORMATION

"Pinnacle" can be applied to 2-6 leaf field corn (1-3 collars, up to approximately 8 inches tall) at a rate of 1/4 ounce per acre. Do not apply to field corn taller than 8 inches or 3 collars, whichever is more restrictive.

Apply "Pinnacle" to field corn hybrids with a Relative Maturity (RM) of 88 days or more, including "food grade" (hollow dent, hard endosperm), waxy and DuPont Optimum High-Oil corn. Not all field corn hybrids of less than 88 days RM, not all white corn hybrids nor Hi-Lysine hybrids have been tested for crop safety, not does DuPont have access to all seed company data. Consequently, injury arising from the use of "Pinnacle" on these types of corn is the responsibility of the user. Consult with your seed supplier before applying "Pinnacle" to any of these corn types.

Apply with ground equipment set to deliver 10-40 GPA. Use only flat fan nozzles operating at 20-40 PSI. Do not make more than one application per season.

TIMING TO WEEDS

Apply to weeds whose first true leaves are expanded but before weeds exceed the sizes listed below.

When applied as directed, "Pinnacle" will control the following weeds:

WEED**SIZE (Inches)**

Velvetleaf	2-6
Pigweed species	2-12
Lambsquarters	2-4
Annual smartweeds	2-6
Wild mustard	up to 4(*)

(*) indicates diameter

Apply in 10-40 gallons of water per acre. Always add either nonionic surfactant at 0.25% v/v (1 qt/100 gal) or crop oil concentrate at 1% v/v (1 gal/100 gal) plus either ammonium nitrogen solution such as 28% UAN (2-4 qt/acre) of ammonium sulfate (2-4 lb/acre).

SOIL INSECTICIDE INTERACTIONS

"Pinnacle" may interact with certain insecticides previously applied to the crop. Crop response varies with field corn type, insecticide used, insecticide application method, and soil type. Corn types may be classified as conventional field corn, imazethapyr tolerant (IT), or imidazilinone-resistant (IR) such as Pioneer 3180IR, etc.

CORN TYPE**For Conventional and "IT"**

"Pinnacle" may be applied to corn previously treated with DuPont Fortress[®] Insecticide or Force² soil insecticide regardless of soil type.

• DO NOT APPLY "Pinnacle" to corn previously treated with Counter 15G¹.

• APPLICATIONS of "Pinnacle" TO CORN PREVIOUSLY TREATED WITH "COUNTER 20 CR" OR "THIMET" MAY CAUSE UNACCEPTABLE CROP INJURY, ESPECIALLY ON SOILS OF LESS THAN 4% ORGANIC MATTER.

Applications of "Pinnacle" to corn previously treated with Dyfonate², Lorsdan³, or other organophosphate insecticides not listed above, may result in temporary crop injury.

For product information call 1-888-6-DUPONT

© 1998 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898

2/2

For "IR" Corn

There are no use precautions for applications of "Pinnacle" to "IR" type field corn previously treated with any insecticide.

**IMPORTANT
BEFORE USING "PINNACLE", READ AND
FOLLOW ALL APPLICABLE DIRECTIONS,
RESTRICTIONS AND PRECAUTIONS ON THE
EPA-REGISTERED LABEL.**

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

D-021198

