




352-525

05/03/2000

1/2

DuPont Agricultural Products



SUPPLEMENTAL LABELING

PINNACLE® HERBICIDE FOR USE IN FIELD CORN

"..... A Growing Partnership With Nature"

PINNACLE® HERBICIDE

EPA Reg. No. 352-525

FOR POST EMERGENCE CONTROL OF CERTAIN BROADLEAF WEEDS IN FIELD CORN FOR USE ONLY IN THE STATES OF CT, DE, MA, MD, ME, NH, NC, NJ, NY, PA, RI, VA, VT, WV

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont PINNACLE® Herbicide is recommended for postemergence control of certain broadleaf weeds in field corn.

Do not apply this product through any kind of irrigation system.

Do not graze or feed forage or grain from treated field corn to livestock within 30 days of application.

Do not apply to fields treated with "Counter"™ Insecticide applied at planting or over-the-row at cultivation as severe crop injury may result.

RESTRICTION

This product is limited to ground application only in the State of New York. Do not apply by air.

APPLICATION INFORMATION

PINNACLE® can be applied to 2-6 leaf field corn (1-4 collars, up to 12 inches tall) at a rate of 1/4 ounce per acre. Do not apply to field corn taller than 12 inches or 4 collars, whichever is more restrictive.

Apply PINNACLE® to field corn hybrids with a Relative Maturity (RM) of 88 days or more, including "food grade" (hollow dent, hard endosperm), waxy and DuPont Optimum High-Oil corn. Not all field corn hybrids of less than 88 days RM, not all white corn hybrids nor Hi-Lysine hybrids have been tested for crop safety, nor does DuPont have access to all seed company data. Consequently, injury arising from the use of PINNACLE® on these types of corn is the responsibility of the user. Consult with your seed supplier before applying PINNACLE® to any of these corn types.

Apply with ground equipment set to deliver 10-40 GPA. Use only flat fan nozzles operating at 20-40 PSI. Do not make more than one application per season.

TIMING TO WEEDS

Apply to weeds whose first true leaves are expanded but before weeds exceed the sizes listed below.

When applied as directed, PINNACLE® will control the following weeds:

<u>WEED</u>	<u>SIZE (Inches)</u>
Velvetleaf	2-6
Pigweed species	2-12
Lambsquarters	2-4
Annual smartweeds	2-6
Wild mustard	up to 4(')

(') indicates diameter

Apply in 10-40 gallons of water per acre. Always add either nonionic surfactant at 0.25% v/v (1 qt/100 gal) or crop oil concentrate at 1% v/v (1 gal/100 gal) plus either ammonium nitrogen solution such as 28% UAN (2-4 qt/acre) of ammonium sulfate (2-4 lb/acre).

SOIL INSECTICIDE INTERACTIONS

PINNACLE® may interact with certain insecticides previously applied to the crop. Crop response varies with field corn type, insecticide used, insecticide application method, and soil type. Corn types may be classified as conventional field corn, imazethapyr tolerant (IT), or imidazilinone-resistant (IR) such as Pioneer 3180IR, etc.

CORN TYPE

For Conventional and "IT"

PINNACLE® may be applied to corn previously treated with DuPont FORTRESS® Insecticide or "Force"™ soil insecticide regardless of soil type.

• DO NOT APPLY PINNACLE® to corn previously treated with Counter 15G¹.

For product information call 1-888-6-DUPONT

© 2000 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898

• APPLICATIONS of PINNACLE® TO CORN PREVIOUSLY TREATED WITH "COUNTER 20 CR" OR "THIMET" MAY CAUSE UNACCEPTABLE CROP INJURY, ESPECIALLY ON SOILS OF LESS THAN 4% ORGANIC MATTER.

Applications of PINNACLE® to corn previously treated with "Dyfonate", "Lorsban", or other organophosphate insecticides not listed above, may result in temporary crop injury.

For "IR" Corn

There are no use precautions for applications of PINNACLE® to "IR" type field corn previously treated with any insecticide.

**IMPORTANT
BEFORE USING PINNACLE® , READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.**

This label contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

¹ Trademark of American Cyanamid Company
² Trademark of Zeneca, Inc.
³ Trademark of Dow Elanco

DR-010 042600
(Replaces: H-63473)

