

PAU ... 134

MAR 29 1991

Dr. Tony E. Catka
 E.I. duPont de Nemours & Co., Inc.
 Agricultural Products Department
 Walker's Mill Bldg./Barley Mill Plaza
 Wilmington, Delaware 19880-0038

Dear Dr. Catka:

Subject: DuPont Accent Herbicide
 EPA Reg. No. 352-534
 Re: Label Revision - Use with Counter
 20 CR for Corn
 Your letter dated March 26, 1991

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable subject to the comments listed below.

Please note that this Agency is developing a policy statement that, if adopted, will require registrant statements similar to the subject amendment for all limitations and warnings added to the label because of data or information not submitted to the Agency as a condition of registration.

If you have any questions, please call Thomas L. Adamczyk (703 557-1650).

Sincerely,
 /s/ **T. E. Adamczyk**

Robert J. Taylor, Product Manager 25
 Fungicide-Herbicide Branch
 Registration Division (H7505C)

BEST AVAILABLE COPY

CONCURRENCES							
BOL							
E							

ACCENT® HERBICIDE

For Use On Field Corn, Popcorn and Field Corn Grown for Seed

This product is a water dispersible granule containing 75 percent active ingredient by weight.

ACTIVE INGREDIENT:


2-((((4,6-Dimethoxypyrimidin- 2-yl)aminocarbonyl) aminosulfonyl))-N,N-dimethyl-3-pyridinecarboxamide

.....75.0%

BY WEIGHT

INERT INGREDIENTS:

.....25.0%



TOTAL.....100.0%

EPA Reg. No. 352 - 534

U.S. Patent No. 4,780,383

CAUTION

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION! Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If on skin: Wash with plenty of soap and water.

If swallowed: No specific intervention is indicated as the compound is not likely to be hazardous by ingestion. However consult a physician if necessary.

For medical emergencies involving this product, call toll free 1-800-441-3637.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment washwaters. Do not apply where/when conditions could favor runoff. Do not apply if rainfall or storm is expected within 24 hours.

IMPORTANT

Injury to or loss of desirable trees or vegetation may result from failure to observe the following: Do not apply, or drain or flush equipment, on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water. Thoroughly clean all mixing and spray equipment immediately after application of ACCENT. (Refer to SPRAYER CLEANUP section of this label.) Failure to follow these procedures may result in injury to subsequently sprayed plants.

DIRECTIONS FOR USE

BEST AVAILABLE COPY

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

ACCENT Herbicide should be used only in accordance with recommendations on this label or in separately published Du Pont

recommendations.

Du Pont will not be responsible for losses or damage resulting from the use of this product in any manner not specifically recommended by Du Pont.

5 Single or sequential applications of ACCENT may not exceed a total of 1-1/3 oz/acre in any one crop year.

Do not graze or feed forage or grain from the treated areas to livestock within 30 days after applications of ACCENT.

10 Do not apply after 10-collar stage of corn, or corn taller than 36 inches, whichever occurs first. Refer to APPLICATION RECOMMENDATION section for specific information.

15 Application of ACCENT to corn that is stressed or damaged by cold weather, hail, drought, water-saturated soil, disease or insects may cause further crop injury. Applications made under certain conditions such as prolonged cool weather (daily high temperatures less than 50 degrees F), widely fluctuating day/night temperatures just prior to or soon after treatment, or during periods of high relative humidity and temperature may cause temporary yellowing and/or crop stunt.

20 Do not apply to corn that has been treated within seven days before with foliar-applied organophosphate insecticides such as Lorsban1, malathion, parathion, etc., or with the herbicides Basagran2, Ladonk2, or Tandem1 because severe crop injury may result. Do not apply to corn that has been treated within seven days before with 2,4-D because severe antagonism may result. Do not apply any of these materials within three days after applying ACCENT or severe crop injury or antagonism may result.

25 Do not apply if crop was previously treated with Counter3 15G (all application methods) as severe crop injury may result.

Do not apply if crop was previously treated with "Counter" 20CR applied in furrow at planting or over-the-row at cultivation.

30 Applications of ACCENT to crops previously treated with "Counter" 20CF applied in a band at planting may cause crop injury. DuPont has not conducted a sufficient number of field trials with this combination to be sure of crop safety. Therefore, DuPont will not be held responsible for losses or damage resulting from such use.

Applications to field corn treated with organophosphate soil insecticides such as Thimet3, Dyfote4, "Lorsban", etc., applied at planting or over-the-row at cultivation, may result in temporary crop injury.

35 Growers are directed to contact the seed supplier for full information and recommendations concerning the use of ACCENT on popcorn or field corn grown for seed and also the effect of prior use of a soil-applied organophosphate insecticide. In all cases, the use of ACCENT on popcorn or field corn grown for seed which has been treated with "Counter" insecticide is prohibited. See the APPLICATION RECOMMENDATIONS section for additional information.

40 Do not apply ACCENT to corn that exhibits injury from herbicide applications made to the current crop or previous crop.

In flood or furrow-irrigated fields, if no rain has occurred after the ACCENT treatment (at least 1/2 inch), do not use tailwater from the first irrigation after ACCENT application to irrigate any crop other than field corn, popcorn or field corn grown for seed.

45 For ground application to weeds when dry, dusty field conditions exist, control of weeds in wheel track areas may be reduced.

Do not allow spray from application equipment to drift onto adjacent crops or land as even small amounts may injure other plants. When spraying near adjacent crops or plants, do everything possible to reduce spray drift.

50 GENERAL INFORMATION

BEST AVAILABLE COPY

ACCENT Herbicide is a water dispersible granule containing 75 percent active ingredient by weight.

55 ACCENT Herbicide is a selective herbicide for the control of annual and perennial grass weeds and selected broadleaf weeds when applied post-emergence to field corn, popcorn or field corn grown for seed. Do not use on sweet corn.

NOTE: Prior to using ACCENT, consideration should be given to crop rotational plans. Rotational crops other than corn may be very sensitive to low concentrations of ACCENT remaining in the soil. See ROTATIONAL CROP GUIDELINES tables for

recropping guidelines.

APPLICATION RECOMMENDATIONS

5 Application should be made only by ground equipment to young, actively growing weeds. Weed control may be reduced if application is made to plants under stress due to drought and/or cold temperatures (less than 50 degrees F). Because ACCENT provides weed control mainly through foliar absorption, ACCENT should be applied after weeds have emerged.

10 Foliar absorption is the primary means of ACCENT uptake by plants. Therefore, select a spray volume and delivery system that will insure thorough coverage and a uniform spray pattern. Increase spray volume and pressure as weed density and size increases. Avoid overlapping, and shut off spray booms while starting, turning, slowing or stopping, or injury to the crop might result.

Corn Growth Stage

15 Field Corn: ACCENT may be applied broadcast or with drop nozzles on corn up to 24 inches tall (free-standing). Method of application is dependent upon the weed:crop height differential. For corn 24-36 inches tall (free-standing), apply ACCENT with drop nozzles. Do not apply to field corn taller than 36 inches or exhibiting 10 collars, whichever is the most restrictive.

20 Field Corn Grown for Seed: Various seed companies have tested a wide range of inbreds for sensitivity to ACCENT and have reported excellent crop safety. However, because of the proprietary nature of seed corn production, Du Pont has not been given access to seed company data. Consequently, Du Pont cannot be held responsible for any crop injury arising from the use of ACCENT on seed corn. The grower is directed to contact the seed company for full information and recommendations regarding the use of ACCENT on inbreds as well as the effect on selectivity from the prior use of a soil-applied organophosphate insecticide.

25 In all cases, do not apply ACCENT to seed corn which has been treated with "Counter" insecticide.

Do not apply to seed corn taller than 20 inches.

30 Popcorn: Various popcorn companies have tested hybrids for sensitivity to ACCENT and have reported excellent crop safety. However, not all popcorn hybrids have been tested. Consequently, Du Pont cannot be held responsible for any crop injury arising from the use of ACCENT on popcorn. The grower is directed to contact the contracting popcorn company for full information and recommendations regarding the use of ACCENT on popcorn hybrids as well as the effect on selectivity from the prior use of a soil-applied organophosphate insecticide.

35 In all cases, do not apply ACCENT to popcorn which has been treated with "Counter" insecticide.

Do not apply to popcorn taller than 20 inches.

40 Weed stage: For best results applications should be made to actively growing weeds at growth stages/weed heights no greater than those shown in the ACCENT USE RATE GUIDE. Results of applications at later growth stages may vary from complete control to suppression, depending on the species, stage of growth and environmental conditions.

45 Heavy infestations of grass and/or broadleaf weeds should be treated before they become too competitive with the crop, especially where soil moisture and/or fertility are in marginal supply. Where an early postemergence treatment is deemed necessary, a second application or a timely cultivation may be required to control later emerging weeds.

Weather: Conditions which promote healthy, actively growing plants result in the best weed control performance with ACCENT. Best activity occurs during warm, moist conditions (70 degrees F or more), and adequate soil moisture before, during and immediately after application. Weed control may be reduced when application is made during cool (less than 50 degrees F) dry weather or to weeds hardened off by drought or cold. ACCENT is rainfast 4 to 6 hours after application.

55 Ground application: Spray Equipment: Refer to specific equipment manufacturer's recommendations for additional information on spray volume, pressure, speed, nozzle type and arrangement, nozzle height above the target area, etc., for respective application equipment. Use a nozzle spacing and boom height arrangement that avoids applying an excessive rate of ACCENT directly over the corn rows.

For optimum spray distribution and thorough coverage, use flat fan nozzles at 25-40 psi, or hollow cone nozzles at 40-60 psi.

BEST AVAILABLE COPY

Use a minimum of 10 gallons carrier per acre. Under heavy weed pressure, dense crop foliage, or moisture stress conditions, increase carrier volume to at least 15 gallons per acre and use maximum spray pressure recommended for the nozzle type.

5 Raindrop5 flood and other nozzles which produce very coarse sprays are not recommended for ACCENT applications because weed control performance may be reduced.

Use 50-mesh or larger strainer screens.

Continuous agitation is required to keep ACCENT in suspension.

10 Chemigation: Do not apply this product through any type of irrigation system.

15 SPRAY PREPARATION

It is important that spray equipment is clean and free of existing pesticide deposits before using ACCENT. Follow the cleanup procedures specified on the label of the product(s) previously sprayed. If no cleanout procedure is provided, follow this cleanout procedure for all application equipment before using ACCENT:

20 1. Thoroughly rinse sprayer, tanks, boom and hoses with clean water.

25 2. Partially fill tank with water and add ammonia (1 gallon of household ammonia per 100 gal. of tank volume) or a tank cleaner. Follow individual label instructions for the amount of tank cleaner to use. Complete filling the tank and flush the cleaning solution through the boom hoses. Let stand for 15 minutes while agitating/recirculating and then drain the tank after flushing the hoses booms and nozzles.

3. Thoroughly rinse the sprayer, tanks, boom and hoses with clean water.

30 4. Follow label directions on product(s) previously sprayed for rinsate disposal.

AT THE END OF THE DAY:

To prevent the buildup of dried pesticide deposits, which can accumulate on application equipment during periods when multiple loads of ACCENT will be applied, rinse the interior of the tank at the end of each day of spraying with fresh water.

35 Then, partially fill the tank with water and flush the boom and hoses. Rinsate may be disposed of on-site, if allowed, or at an approved waste disposal facility. Allow the remainder of the water to stand in the tank overnight.

Mix the proper amount of ACCENT into the necessary volume of water in the spray tank with the agitator running, then add the adjuvant(s) and/or any companion herbicide to the tank after all the ACCENT is in suspension.

40 Apply ACCENT spray preparation within 24 hours after mixing or product degradation may occur. If spray preparation is left standing, agitate thoroughly before continuing to spray.

45 Adjuvants: Always add a non-ionic surfactant OR crop oil concentrate (COC) to the spray mixture. Non-ionic surfactants should contain at least 80% active ingredient (of which at least 50% is actual nonionic surfactant). Add surfactant at 0.25% vol./vol. (1 quart per 100 gallons of spray solution). Use only EPA approved surfactants authorized for use on food crops. AVOID PRODUCTS THAT DO NOT ACCURATELY DEFINE THEIR INGREDIENTS.

50 If crop oil concentrate is substituted for non-ionic surfactant, use a good quality petroleum-based COC containing at least 15% emulsifiers/surfactants. Add COC to the spray mixture at 1% vol./vol. (1 gallon per 100 gallons of spray mixture). COC is recommended for certain grasses and for applications made during hot, dry conditions.

55 Adding liquid nitrogen fertilizer solution (e.g., 28-0-0 or 32-0-0) to the spray solution may improve the control of certain weed species. However, a nonionic surfactant or crop oil concentrate will still be necessary. Add liquid nitrogen fertilizer solution at 4% vol./vol. (4 gal. per 100 gal.). Do not use liquid fertilizer solutions or suspensions as the total carrier because excessive crop injury may occur.

Corn tolerance to ACCENT is very high, so adjuvant selection can be based solely on maximizing weed control. If another herbicide is tank mixed with ACCENT to increase broadleaf weed spectrum, adjuvant selection must be determined according

BEST AVAILABLE COPY

to selectivity limitations of the other herbicide. Refer to TANK MIXES section of the label for additional information.

APPLICATION RATE

5

ACCENT may be applied postemergence to control many grass weeds and selected annual broadleaf weeds in field corn, popcorn and field corn grown for seed. Do not apply to sweet corn.

10

For band application, apply proportionately less product than for broadcast applications. To avoid possible crop injury due to improper application, CALIBRATE ROWBANDER UNITS BY FOLLOWING THE DIRECTIONS CONTAINED IN THE DU PONT BULLETIN "APPLICATION ACCURACY - ROWBANDERS"

Apply ACCENT at the rates listed in ACCENT USE RATE GUIDE. Apply when majority of weeds have emerged but before weeds exceed size listed in table.

15

Sequential Applications

Annual weeds sometimes will have more than one flush of seedling emergence and treated perennials may sometimes regrow from underground stems or roots, depending upon rainfall and other environmental conditions. To maximize control of such weeds, it may be necessary to use a sequential application of ACCENT in which the first application goes on early before the weeds become competitive, followed by a second application 14-28 days later, depending upon the timing of a new flush or regrowth. The combined dosage of the sequential applications cannot exceed 1 1/3 oz/acre of ACCENT.

20

Annual weeds: For shattercane, seedling Johnsongrass, and annual grasses — 2/3 oz/acre followed by 2/3 oz/acre 14-28 days later, depending upon the emergence of the new flush.

25

Perennial weeds: For sequential applications to rhizome Johnsongrass apply 2/3 oz. ACCENT when johnsongrass is 8-12" tall, followed by 2/3 oz. ACCENT when new growth is 8-10".

30

"ACCENT" USE RATE GUIDE

Use rate - 2/3 oz. per acre

	Size Range*	
	Height (inches)	
35	Grasses **	
	Shattercane	4-12
	Johnsongrass - seedling	4-12
	- rhizome	8-18
40	Sorghum alatum	4-12
	Foxtails	
	giant	2-4
	yellow	2-4
	bristly	2-4
45	green	2-4
	Fall panicum	2-4
	Texas panicum	1-3
	Itchgrass	2-8
	Barnyardgrass	2-8
50	Broadleaf signalgrass	1-2
	Sandbur	
	field	2-4
	longspine	2-4
	Wild oats	2-4
55	Wild proso millet	1-3
	Woolly Cupgrass	2-4
	Quackgrass	4-10

BEST AVAILABLE COPY

Tank mix ACCENT with 1 1/2 - 3 pints/acre "Buctril+atrazine" and apply before corn is 12 inches tall. Always add a nonionic surfactant at 0.25% vol/vol (1 quart/100 gal.). See SPRAY PREPARATION section of this label for nonionic surfactant guidelines.

- 5 DO NOT ADD CROP OIL CONCENTRATE OR NITROGEN FERTILIZER SOLUTION TO THIS TANK MIXTURE, or excessive crop injury may occur. DO NOT APPLY THIS TANK MIX TO CORN TALLER THAN 12 INCHES.

ACCENT + Banvel⁷: ACCENT may be tank mixed with "Banvel" for improved control of broadleaf weeds. See "Banvel" label for additional weeds controlled, weed sizes and application information. For weeds listed on both ACCENT and "Banvel" labels, follow the weed size guidelines which are least restrictive.

Tank mix ACCENT with 1/2 - 1 pints/acre "Banvel" and always add a nonionic surfactant to this tank mixture at 0.25% vol/vol (1 quart/100 gal.). See SPRAY PREPARATION section of this label for nonionic surfactant guidelines.

- 15 Do not apply after corn reaches eight inches in height. DO NOT ADD CROP OIL CONCENTRATE OR NITROGEN FERTILIZER SOLUTION TO THIS TANK MIXTURE, or excessive crop injury may occur.

ACCENT + Marksman⁷: ACCENT may be tank mixed with "Marksman" for improved control of broadleaf weeds. See "Marksman" label for additional weeds controlled, weed sizes and application conditions. For weeds listed on both ACCENT and "Marksman" labels, follow the weed size guidelines which are least restrictive.

Tank mix ACCENT with 2 - 3 1/2 pints/acre "Marksman" and apply before corn is eight inches tall. Always add a nonionic surfactant at 0.25% vol/vol (1 quart/100 gal.). See SPRAY PREPARATION section of this label for nonionic surfactant guidelines.

Apply this tank mixture before corn reaches eight inches in height. DO NOT ADD CROP OIL CONCENTRATE OR NITROGEN FERTILIZER SOLUTION TO THIS TANK MIXTURE, or excessive crop injury may occur. DO NOT APPLY THIS TANK MIX TO CORN TALLER THAN 12 INCHES.

- 30 NOTE: DO NOT TANKMIX with "Basagran", "Laddok", or "Tandem" as severe crop injury may occur. Do not tank mix with 2,4-D as severe antagonism may occur.

CROP ROTATIONAL GUIDELINES

Rotational crops vary in their response to low concentrations of ACCENT remaining in the soil after corn harvest. The amount of ACCENT which may remain in the soil is dependent on the rate used, soil pH and organic matter content, time elapsed since ACCENT application, and climatic and weather factors.

In order to avoid injury to rotational crops, observe the minimum recrop intervals from date of ACCENT application, as listed in the table ACCENT ROTATIONAL CROP GUIDELINES.

ACCENT ROTATIONAL CROP GUIDELINE - 1 No Soil pH Restrictions

Minimum Time in Months Before
Planting the Rotational Crop

Crop	Minimum Time in Months Before Planting the Rotational Crop
Corn (Field, Seed)	Anytime
Corn (Pop, Sweet) (*)	10
Soybeans	10
Wheat (Winter)	4
Wheat (Spring)	8
Barley (Spring)	8
Rye (Winter)	4
Dry Beans	10
Oats	8
Alfalfa (**)	12

BEST AVAILABLE COPY

Other Crops See ROTATIONAL CROP GUIDELINE - 2

- * Except the sweetcorn varieties "Merit", "Carnival", and "Sweet Success", for which the minimum time interval is 15 months.
- ** Except for the state of Kansas east of Highway 75 and for states east of a line formed by the western borders of Minnesota, Iowa, Missouri, Arkansas and Louisiana, where the minimum time interval is 10 months.

ROTATIONAL CROP GUIDELINE - 2
With Soil pH Restrictions

Crop	Minimum Time in Months Before Planting the Rotational Crop *	
	pH <= 6.5	pH > 6.5
Sorghum	10	18
Cotton	10	18
All other crops not listed in ROTATIONAL GUIDELINE - 1 or 2	10	18

Soil pH is to be determined by laboratory analysis using the 1:1 soil:water suspension method on representative soil samples taken at 0-4" depth. Soil pH varies within fields; therefore, recropping plans should be based on the highest soil pH in each field. Consult local extension publications for recommended soil sampling procedures.

SPRAYER CLEANUP

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately after application of ACCENT:

1. Drain tank; thoroughly rinse spray tank, boom and hoses with clean water. Loosen and physically remove any visible deposits.
2. Fill the tank with clean water, while adding 1 gallon household ammonia (contains 3% active) (*) for every 100 gal. of water. Flush the boom, hoses and nozzles with cleaning solution, then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 minutes. Flush the tank, boom and hoses with the cleaning solution and drain tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat Step 2.
5. Thoroughly rinse the tank, boom and hoses with clean water.
6. Dispose of rinsate on-site or at an approved waste disposal facility.

(*) Equivalent amounts of an alternate strength ammonia solution or a Du Pont-approved cleaner can be used in the cleanout procedure. Du Pont-approved cleaners include Pro-Tank Cleaner[®], Chem-Tank Cleaner and Neutralizer[®], Loveland Tank and Equipment Cleaner 10, Incide-Out 11, and Tank-Aid[®]12. Carefully read and follow the individual cleaner label directions.

STORAGE AND DISPOSAL

Storage: Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed.

Product Disposal: Do not contaminate water, food or feed by disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Emptied containers should be triple-rinsed (or equivalent) and offered for recycling or reconditioning, or punctured and disposed of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If

BEST AVAILABLE COPY

burned, stay out of smoke.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

5

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Du Pont will not be responsible for losses or damage resulting from application of ACCENT when applied to crops previously treated with "Counter" 20GR applied in a band at planting. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESSED OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

- [1] Registered trademark of Dow Elanco
- [2] Registered trademark of BASF Corporation
- [3] Registered trademark of American Cyanamid Co.
- [4] Registered trademark of ICI Americas, Inc.
- [5] Registered trademark of Delavan Corporation
- [6] Registered trademark of Rhone-Poulenc Ag Company
- [7] Registered trademark of Sandoz Ltd
- [8] Manufactured for Cenex/Land O' Lakes Agronomy Company
- [9] Manufactured by Farnbelt Chemicals, Inc.
- [10] Manufactured by Loveland Industries, Inc.
- [11] Manufactured by Precision Laboratories, Inc.
- [12] Manufactured for Combek Chemical Co.

35

D-030004

40

D-030891/032591

BEST AVAILABLE COPY



ACCENT® HERBICIDE

REGISTRATION CODE 0000-000
PACKAGE SIZE 4 - 10 OZ. JUGS/CARTON 12 - CARTONS/BOX

For Use On Field Corn, Popcorn and Field Corn Grown for Seed

This product is a water dispersible granule containing 75 percent active ingredient by weight.

ACTIVE INGREDIENT:

2-((((4,6-Dimethoxypyrimidin-2-yl)aminocarbonyl) amino)sulfonyl)-N,N-dimethyl-3-pyridinecarboxamide75.0%

BY WEIGHT

INERT INGREDIENTS:

.....25.0%

TOTAL.....100.0%

EPA Reg. No. 352 - 534

U.S. Patent No. 4,700,303

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION! Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If on skin: Wash with plenty of soap and water.

If swallowed: No specific intervention is indicated as the compound is not likely to be hazardous by ingestion. However consult a physician if necessary.

For medical emergencies involving this product, call toll free 1-800-441-3637.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment washwaters. Do not apply where/when conditions could favor runoff. Do not apply if rainfall or storm is expected within 24 hours.

IMPORTANT

Injury to or loss of desirable trees or vegetation may result from failure to observe the following: Do not apply, or drain or flush equipment, on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water. Thoroughly clean all mixing and spray equipment immediately after application of ACCENT. (Refer to SPRAYER CLEANUP section of this label.) Failure to follow these procedures may result in injury to subsequently sprayed plants.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

ACCENT Herbicide should be used only in accordance with recommendations on this label or in separately published Du Pont recommendations.

Du Pont will not be responsible for losses or damage resulting from the use of this product in any manner not specifically recommended by Du Pont.

Single or sequential applications of ACCENT may not exceed a total of 1-1/3 oz./acre in any one crop year.

Do not graze or feed forage or grain from the treated areas to livestock within 30 days after applications of ACCENT.

Do not apply after 10-collar stage of corn, or corn taller than 36 inches, whichever occurs first. Refer to APPLICATION RECOMMENDATION section for specific information.

BEST AVAILABLE COPY

Application of ACCENT to corn that is stressed or damaged by cold weather, hail, drought, water-saturated soil, disease or insects may cause further crop injury. Applications made under certain conditions such as prolonged cool weather (daily high temperatures less than 50 degrees F), widely fluctuating day/night temperatures just prior to or soon after treatment, or during periods of high relative humidity and temperature may cause temporary yellowing and/or crop stunting.

Do not apply to corn that has been treated within seven days before with foliar-applied organophosphate insecticides such as Lorsban¹, malathion, parathion, etc., or with the herbicides Basagran², Laddok², or Tandem¹ because severe crop injury may result. Do not apply to corn that has been treated within seven days before with 2,4-D because severe antagonism may result. Do not apply any of these materials within three days after applying ACCENT or severe crop injury or antagonism may result.

Do not apply if crop was previously treated with Counter³ 15G (all application methods) as severe crop injury may result.

Do not apply if crop was previously treated with "Counter" 20CR applied in furrow at planting or over-the-row at cultivation.

Applications of ACCENT to crops previously treated with "Counter" 20CR applied in a band at planting may cause crop injury. DuPont has not conducted a sufficient number of field trials with this combination to be sure of crop safety. Therefore, DuPont will not be held responsible for losses or damage resulting from such use.

Applications to field corn treated with organophosphate soil insecticides such as Thimet³, Dylonate⁴, "Lorsban", etc., applied at planting or over-the-row at cultivation, may result in temporary crop injury.

Growers are directed to contact the seed supplier for full information and recommendations concerning the use of ACCENT on popcorn or field corn grown for seed and also the effect of prior use of a soil-applied organophosphate insecticide. In all cases, the use of ACCENT on popcorn or field corn grown for seed which has been treated with "Counter" insecticide is prohibited. See the APPLICATION RECOMMENDATIONS section for additional information.

Do not apply ACCENT to corn that exhibits injury from herbicide applications made to the current crop or previous crop.

In flood or furrow-irrigated fields, if no rain has occurred after the ACCENT treatment (at least 1/2 inch), do not use tailwater from the first irrigation after ACCENT application to irrigate any crop other than field corn, popcorn or field corn grown for seed⁴.

For ground application to weeds when dry, dusty field conditions exist, control of weeds in wheel track areas may be reduced.

Do not allow spray from application equipment to drift onto adjacent crops or land as even small amounts may injure other plants. When spraying near adjacent crops or plants, do everything possible to reduce spray drift.

GENERAL INFORMATION

ACCENT Herbicide is a water dispersible granule containing 75 percent active ingredient by weight.

ACCENT Herbicide is a selective herbicide for the control of annual and perennial grass weeds and selected broadleaf weeds when applied post-emergence to field corn, popcorn or field corn grown for seed. Do not use on sweet corn.

NOTE: Prior to using ACCENT, consideration should be given to crop rotational plans. Rotational crops other than corn may be very sensitive to low concentrations of ACCENT remaining in the soil. See ROTATIONAL CROP GUIDELINES tables for recropping guidelines.

APPLICATION RECOMMENDATIONS

Application should be made only by ground equipment to young, actively growing weeds. Weed control may be reduced if application is made to plants under stress due to drought and/or cold temperatures (less than 50 degrees F). Because ACCENT provides weed control mainly through foliar absorption, ACCENT should be applied after weeds have emerged.

Foliar absorption is the primary means of ACCENT uptake by plants. Therefore, select a spray volume and delivery system that will insure thorough coverage and a uniform spray pattern. Increase spray volume and pressure as weed density and size increases. Avoid overlapping, and shut off spray booms while starting, turning, slowing or stopping, or injury to the crop might result.

Corn Growth Stage

Field Corn: ACCENT may be applied broadcast or with drop nozzles on corn up to 24 inches tall (free-standing). Method of application is dependent upon the weed:crop height differential. For corn 24-36 inches tall (free-standing), apply ACCENT with drop nozzles. Do not apply to field corn taller than 36 inches or exhibiting 10 collars, whichever is the most restrictive.

Field Corn Grown for Seed: Various seed companies have tested a wide range of inbreds for sensitivity to ACCENT and have reported excellent crop safety. However, because of the proprietary nature of seed corn production, Du Pont has not been given access to seed company data. Consequently, Du Pont cannot be held responsible for any crop injury arising from the use of ACCENT on seed corn. The grower is directed to contact the seed company for full information and recommendations regarding the use of ACCENT on inbreds as well as the effect on selectivity from the prior use of a soil-applied organophosphate insecticide.

In all cases, do not apply ACCENT to seed corn which has been treated with "Counter" insecticide.

Do not apply to seed corn taller than 20 inches.

Popcorn: Various popcorn companies have tested hybrids for sensitivity to ACCENT and have reported excellent crop safety. However, not all popcorn hybrids have been tested. Consequently, Du Pont cannot be held responsible for any crop injury arising from the use of ACCENT on popcorn. The grower is directed to contact the contracting popcorn company for full information and recommendations regarding the use of ACCENT on popcorn hybrids as well as the effect on selectivity from the prior use of a soil-applied organophosphate insecticide.

In all cases, do not apply ACCENT to popcorn which has been treated with "Counter" insecticide.

Do not apply to popcorn taller than 20 inches.

Weed stage: For best results applications should be made to actively growing weeds at growth stages/seed heights no greater than those shown in the ACCENT USE RATE GUIDE. Results of applications at later growth stages may vary from complete control to suppression, depending on the species, stage of growth and environmental conditions.

Heavy infestations of grass and/or broadleaf weeds should be treated before they become too competitive with the crop, especially where soil moisture and/or fertility are in marginal supply. Where an early post-emergence treatment is deemed necessary, a second application or a timely cultivation may be required to control later emerging weeds.

BEST AVAILABLE COPY

Weather: Conditions which promote healthy, actively growing plants result in the best weed control performance with ACCENT. Best activity occurs during warm, moist conditions (70 degrees F or more), and adequate soil moisture before, during and immediately after application. Weed control may be reduced when application is made during cool (less than 50 degrees F) dry weather or to weeds hardened off by drought or cold. ACCENT is rainfast 4 to 6 hours after application.

Ground application: Spray Equipment: Refer to specific equipment manufacturer's recommendations for additional information on spray volume, pressure, speed, nozzle type and arrangement, nozzle height above the target area, etc., for respective application equipment. Use a nozzle spacing and boom height arrangement that avoids applying an excessive rate of ACCENT directly over the corn rows.

For optimum spray distribution and thorough coverage, use flat fan nozzles at 25-40 psi, or hollow cone nozzles at 40-60 psi. Use a minimum of 10 gallons carrier per acre. Under heavy weed pressure, dense crop foliage, or moisture stress conditions, increase carrier volume to at least 15 gallons per acre and use maximum spray pressure recommended for the nozzle type.

Raindrop⁵ flood and other nozzles which produce very coarse sprays are not recommended for ACCENT applications because weed control performance may be reduced.

Use 50-mesh or larger strainer screens.

Continuous agitation is required to keep ACCENT in suspension.

Chemigation: Do not apply this product through any type of irrigation system.

SPRAY PREPARATION

It is important that spray equipment is clean and free of existing pesticide deposits before using ACCENT. Follow the cleanup procedures specified on the label of the product(s) previously sprayed. If no cleanup procedure is provided, follow the cleanup procedure for all application equipment before using ACCENT:

1. Thoroughly rinse sprayer, tanks, boom and hoses with clean water.
2. Partially fill tank with water and add ammonia (1 gallon of household ammonia per 100 gal. of tank volume) or a tank cleaner. Follow individual label instructions for the amount of tank cleaner to use. Complete filling the tank and flush the cleaning solution through the boom hoses. Let stand for 15 minutes while agitating/recirculating and then drain the tank after flushing the hoses booms and nozzles.
3. Thoroughly rinse the sprayer, tanks, boom and hoses with clean water.
4. Follow label directions on product(s) previously sprayed for rinse disposal.

AT THE END OF THE DAY:

To prevent the buildup of dried pesticide deposits, which can accumulate on application equipment during periods when multiple loads of ACCENT will be applied, rinse the interior of the tank at the end of each day of spraying with fresh water. Then, partially fill the tank with water and flush the boom and hoses. Rinseate may be disposed of on-site, if allowed, or at an approved waste disposal facility. Allow the remainder of the water to stand in the tank overnight.

Mix the proper amount of ACCENT into the necessary volume of water in the spray tank with the agitator running, then add the adjuvant(s) and/or any companion herbicide to the tank after all the ACCENT is in suspension.

Apply ACCENT spray preparation within 24 hours after mixing or product degradation may occur. If spray preparation is left standing, agitate thoroughly before continuing to spray.

Adjuvants: Always add a non-ionic surfactant OR crop oil concentrate (COC) to the spray mixture. Non-ionic surfactants should contain at least 80% active ingredient (of which at least 50% is actual anionic surfactant). Add surfactant at 0.25% vol./vol. (1 quart per 100 gallons of spray solution). Use only EPA approved surfactants authorized for use on food crops. AVOID PRODUCTS THAT DO NOT ACCURATELY DEFINE THEIR INGREDIENTS.

If crop oil concentrate is substituted for non-ionic surfactant, use a good quality petroleum-based COC containing at least 15% emulsifiers/surfactants. Add COC to the spray mixture at 1% vol./vol. (1 gallon per 100 gallons of spray mixture). COC is recommended for certain grasses and for applications made during hot, dry conditions.

Adding liquid nitrogen fertilizer solution (e.g., 28-0-0 or 32-0-0) to the spray solution may improve the control of certain weed species. However, a non-ionic surfactant or crop oil concentrate will still be necessary. Add liquid nitrogen fertilizer solution at 4% vol./vol. (4 gal. per 100 gal.). Do not use liquid fertilizer solutions or suspensions as the total carrier because excessive crop injury may occur.

Corn tolerance to ACCENT is very high, so adjuvant selection can be based solely on maximizing weed control. If another herbicide is tank mixed with ACCENT to increase broadleaf weed spectrum, adjuvant selection must be determined according to selectivity limitations of the other herbicide. Refer to TANK MIXES section of the label for additional information.

APPLICATION RATE

ACCENT may be applied postemergence to control many grass weeds and selected annual broadleaf weeds in field corn, popcorn and field corn grown for seed. Do not apply to sweet corn.

For band application, apply proportionately less product than for broadcast applications. To avoid possible crop injury due to improper application, CALIBRATE ROWBANDER UNITS BY FOLLOWING THE DIRECTIONS CONTAINED IN THE DU PONT BULLETIN "APPLICATION ACCURACY - ROW-BANDERS"

Apply ACCENT at the rates listed in ACCENT USE RATE GUIDE. Apply when majority of weeds have emerged but before weeds exceed size listed in table.

Sequential Applications

Annual weeds sometimes will have more than one flush of seedling emergence and treated perennials may sometimes regrow from underground stems or roots, depending upon rainfall and other environmental conditions. To maximize control of such weeds, it may be necessary to use a sequential application of ACCENT in which the first application goes on early before the weeds become competitive, followed by a second application 14-28 days later, depending upon the timing of a new flush or regrowth. The combined dosage of the sequential applications cannot exceed 1 1/3 oz/acre of ACCENT.

BEST AVAILABLE COPY

Annual weeds: For shattercane, seedling Johnsongrass, and annual grasses — 2/3 oz/acre followed by 2/3 oz/acre 14-28 days later, depending upon the emergence of the new flush.

Perennial weeds: For sequential applications to rhizome Johnsongrass apply 2/3 oz. ACCENT when johnsongrass is 9-12" tall, followed by 2/3 oz. ACCENT when new growth is 8-10".

"ACCENT" USE RATE GUIDE

Use rate - 2/3 oz. per acre

Grasses**	Size Range* Height (inches)
Shattercane	4-12
Johnsongrass - seedling	4-12
- rhizome	8-18
Sorghum alnum	4-12
Foxtails	
giant	2-4
yellow	2-4
bristly	2-4
green	2-4
Fall panicum	2-4
Texas panicum	1-3
Itchgrass	2-6
Barnyardgrass	2-6
Broadleaf signalgrass	1-2
Sandbur	
field	2-4
longspine	2-4
Wild oats	2-4
Wild proso millet	1-3
Woolly Cupgrass	2-4
Quackgrass	4-10
Broadleaves	Height (inches)
Morningglory	
lyleaf	2-3
Tall	1-2
Pitted	1-2
Pigweed	
Smooth	1-4
Redroot	1-4
Smartweed, annual	1-4
Jimsonweed	1-3
Bur cucumber	1-3

* Optimum activity for this size range can be achieved with 2/3 oz. product. Weeds which exceed this height range by up to 50% may be partially controlled with rates between 2/3 and 1 1/3 oz product per acre.

** The use of a second application of 2/3 oz. per acre or a timely cultivation may be necessary due to the effects of adverse weather conditions following the initial Accent application.

Cultivation

Cultivation may be used as an additional weed control measure to remove weed escapes following ACCENT treatment and/or as an alternative to a sequential ACCENT treatment. Ideal timing for cultivation is 7-14 days after ACCENT application. Do not cultivate within 10 days prior to ACCENT application, or control may be reduced.

TANK MIXES

ACCENT + atrazine: ACCENT may be tank mixed with atrazine for improved control of broadleaf weeds. See atrazine label for additional weeds controlled, weed sizes and application conditions. For weeds listed on both ACCENT and atrazine labels, follow the weed size guidelines which are least restrictive.

Tank mix ACCENT with 3/4 - 1 1/2 quarts/acre atrazine 4L or 0.83 - 1.66 lb atrazine 90% dry flowable and apply before corn is 12 inches tall. Always add a crop oil concentrate at 1% vol/vol (1 gal/100 gal.). See SPRAY PREPARATION section of this label for crop oil concentrate guidelines. A nitrogen fertilizer solution (20%, 32% N) may be added at 4% vol/vol (4 gal/100 gal) to enhance ACCENT performance on certain weeds, but crop oil concentrate must always be added.

DO NOT APPLY THIS TANK MIX TO CORN TALLER THAN 12 INCHES.

ACCENT + Buctril®: ACCENT may be tank mixed with "Buctril" for improved control of broadleaf weeds. See "Buctril" label for additional weeds controlled, weed sizes and application information. For weeds listed on both ACCENT and "Buctril" labels, follow the weed size guidelines which are least restrictive.

Tank mix ACCENT with 1 - 1 1/2 quarts/acre "Buctril" and always add a nonionic surfactant to this tank mixture at 0.25% vol/vol (1 quart/100 gal). See SPRAY PREPARATION section of this label for nonionic surfactant guidelines. Apply either broadcast or with drop nozzles to corn up to 24 inches tall. Apply only with drop nozzles on corn 24-36 inches tall. Do not apply to corn taller than 36 inches.

DO NOT ADD CROP OIL CONCENTRATE OR NITROGEN FERTILIZER SOLUTION TO THIS TANK MIXTURE, or excessive crop injury may occur.

ACCENT + Buctril+atrazine®: ACCENT may be tank mixed with "Buctril+atrazine" for improved control of broadleaf weeds. See "Buctril+atrazine" label for additional weeds controlled, weed sizes and application conditions. For weeds listed on both ACCENT and "Buctril+atrazine" labels, follow the weed size guidelines which are least restrictive.

Tank mix ACCENT with 1 1/2 - 3 quarts/acre "Buctril+atrazine" and apply before corn is 12 inches tall. Always add a nonionic surfactant at 0.25% vol/vol (1 quart/100 gal.). See SPRAY PREPARATION section of this label for nonionic surfactant guidelines.

DO NOT ADD CROP OIL CONCENTRATE OR NITROGEN FERTILIZER SOLUTION TO THIS TANK MIXTURE, or excessive crop injury may occur. DO NOT APPLY THIS TANK MIX TO CORN TALLER THAN 12 INCHES.

BEST AVAILABLE COPY

ACCENT + Banvel⁷: ACCENT may be tank mixed with "Banvel" for improved control of broadleaf weeds. See "Banvel" label for additional weeds controlled, weed sizes and application information. For weeds listed on both ACCENT and "Banvel" labels, follow the weed size guidelines which are least restrictive.

Tank mix ACCENT with 1/2 - 1 pints/acre "Banvel" and always add a non-ionic surfactant to this tank mixture at 0.25% vol/vol (1 quart/100 gal). See SPRAY PREPARATION section of this label for nonionic surfactant guidelines.

Do not apply after corn reaches eight inches in height. **DO NOT ADD CROP OIL CONCENTRATE OR NITROGEN FERTILIZER SOLUTION TO THIS TANK MIXTURE**, or excessive crop injury may occur.

ACCENT + Marksman⁷: ACCENT may be tank mixed with "Marksman" for improved control of broadleaf weeds. See "Marksman" label for additional weeds controlled, weed sizes and application conditions. For weeds listed on both ACCENT and "Marksman" labels, follow the weed size guidelines which are least restrictive.

Tank mix ACCENT with 2 - 3 1/2 pints/acre "Marksman" and apply before corn is eight inches tall. Always add a nonionic surfactant at 0.25% vol/vol (1 quart/100 gal). See SPRAY PREPARATION section of this label for non-ionic surfactant guidelines.

Apply this tank mixture before corn reaches eight inches in height. **DO NOT ADD CROP OIL CONCENTRATE OR NITROGEN FERTILIZER SOLUTION TO THIS TANK MIXTURE**, or excessive crop injury may occur. **DO NOT APPLY THIS TANK MIX TO CORN TALLER THAN 12 INCHES.**

NOTE: DO NOT TANKMIX with "Basagran", "Laddok", or "Tandem" as severe crop injury may occur. Do not tank mix with 2,4-D as severe antagonism may occur.

CROP ROTATIONAL GUIDELINES

Rotational crops vary in their response to low concentrations of ACCENT remaining in the soil after corn harvest. The amount of ACCENT which may remain in the soil is dependent on the rate used, soil pH and organic matter content, time elapsed since ACCENT application, and climatic and weather factors.

In order to avoid injury to rotational crops, observe the minimum recrop intervals from date of ACCENT application, as listed in the table ACCENT ROTATIONAL CROP GUIDELINES.

ACCENT ROTATIONAL CROP GUIDELINE - 1 No Soil pH Restrictions	
Crop	Minimum Time in Months Before Planting the Rotational Crop
Corn (Field, Seed)	Anytime
Corn (Pop, Sweet) (*)	10
Soybeans	10
Wheat (Winter)	4
Wheat (Spring)	8
Barley (Spring)	8
Rye (Winter)	4
Dry Beans	10
Oats	8
Alfalfa (**)	12
Other Crops	See ROTATIONAL CROP GUIDELINE - 2

* Except the sweetcorn varieties "Merf", "Carnival", and "Sweet Success", for which the minimum time interval is 15 months.
 ** Except for the state of Kansas east of Highway 76 and for states east of a line formed by the western borders of Minnesota, Iowa, Missouri, Arkansas and Louisiana, where the minimum time interval is 10 months.

ROTATIONAL CROP GUIDELINE - 2 With Soil pH Restrictions		
Crop	Minimum Time in Months Before Planting the Rotational Crop *	
	pH < 6.5	pH > 6.5
Sorghum	10	18
Cotton	10	18
All other crops not listed in ROTATIONAL GUIDELINE - 1 or 2	10	18

Soil pH is to be determined by laboratory analysis using the 1:1 soil/water suspension method on representative soil samples taken at 0-4" depth. Soil pH varies within fields; therefore, recropping plans should be based on the highest soil pH in each field. Consult local extension publications for recommended soil sampling procedures.

SPRAYER CLEANUP

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately after application of ACCENT:

1. Drain tank; thoroughly rinse spray tank, boom and hoses with clean water. Loosen and physically remove any visible deposits.
2. Fill the tank with clean water, while adding 1 gallon household ammonia (contains 3% active) (*) for every 100 gal. of water. Flush the boom, hoses and nozzles with cleaning solution, then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 minutes. Flush the tank, boom and hoses with the cleaning solution and drain tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat Step 2.
5. Thoroughly rinse the tank*, boom and hoses with clean water.
6. Dispose of rinsate on-site or at an approved waste disposal facility.

(*) Equivalent amounts of an alternate strength ammonia solution or a Du Pont-approved cleaner can be used in the cleanout procedure. Du Pont-approved cleaners include Pro-Tank Cleaner[®], Chain-Tank Cleaner and Neutralizer[®], Loveland Tank and Equipment Cleaner[™], Insite-Out[™], and Tank-Aid[™]. Carefully read and follow the individual cleaner label directions.

STORAGE AND DISPOSAL

Storage: Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed.

Product Disposal: Do not contaminate water, food or feed by disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Emptied containers should be triple-rinsed (or equivalent) and offered for recycling or reconditioning, or punctured and disposed of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Du Pont will not be responsible for losses or damage resulting from application of ACCENT when applied to crops previously treated with "Counter" 20CR applied in a band at planting. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESSED OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

- {1} Registered trademark of Dow Elanco
- {2} Registered trademark of BASF Corporation
- {3} Registered trademark of American Cyanamid Co.
- {4} Registered trademark of ICI America, Inc.
- {5} Registered trademark of Dalsven Corporation
- {6} Registered trademark of Rhone-Poulenc Ag Company
- {7} Registered trademark of Sandoz Ltd
- {8} Manufactured for Conax/Land O Lakes Agronomy Company
- {9} Manufactured by Farmbelt Chemicals, Inc.
- {10} Manufactured by Loveland Industries, Inc.
- {11} Manufactured by Precision Laboratories, Inc.
- {12} Manufactured for Corbett Chemical Co

D-030891/032591

BEST AVAILABLE COPY