



000352-00538-020899


Printed: 13:18:52 Friday, 21 May, 1999 # 93 / 2959

Systems Integration Group, Inc.



Harmony[®] Extra

herbicide



"..... A Growing Partnership With Nature"

"HARMONY EXTRA" HIGHLIGHTS

- *For selective postemergence broadleaf weed control in Wheat, Barley, Oat, and Fallow*
- *Apply at the rate of 0.3 to 0.6 ounce per acre on Wheat, Barley and Fallow, 0.3 to 0.4 ounce per acre on Oat (see Application Information).*
- *Apply after the crop is in the 2-leaf stage, but before the flag leaf is visible on Wheat, Barley and Winter Oat. On Spring Oat, apply after the crop is in the 3 leaf stage, but before jointing.*
- *May be applied by ground or by air.*
- *Use in tank mixtures with other registered herbicides for broader spectrum weed control (see Tank Mixtures).*
- *Rotate to Wheat, Barley, and Oat anytime. Rotate to Sugarbeets, Winter Rape and Canola after 60 days. Rotate to any other crop after 45 days.*
- *Consult label text for complete instructions. Always read and follow label "Directions For Use".*

TABLE OF CONTENTS

PRECAUTIONARY STATEMENTS	1
DIRECTIONS FOR USE	2
GENERAL INFORMATION	2
Environmental Conditions and Biological Activity	2
APPLICATION INFORMATION	2
Use Rate	2
Application Timing	3
Crop Rotation	3
Weeds Controlled	3
Weeds Partially Controlled	4
Specific Weed Problems.....	4
Surfactants	4
Ground Application	4
Aerial Application	5
Product Measurement	5
TANK MIXTURES	5
MIXING INSTRUCTIONS	6
GRAZING	6
SPRAY EQUIPMENT	7
SPRAYER CLEANUP	7
At the End of the Day	7
Before Spraying Other Crops	7
SPRAY DRIFT MANAGEMENT	7
Importance of Droplet Size	7
Controlling Droplet Size - General Techniques	7
Controlling Droplet Size - Aircraft	8
Boom Height	8
Wind	8
Temperature and Humidity	8
Temperature Inversions	8
Shielded Sprayers	8
Air-Assisted (Air Blast) Field Crop Sprayers	8
RESISTANCE	8
INTEGRATED PEST MANAGEMENT	8
PRECAUTIONS	9
STORAGE AND DISPOSAL	9
NOTICE OF WARRANTY	10



Harmony[®] Extra

herbicide

Dry flowable

For Use on Wheat, Barley, Oat and Fallow

Active Ingredients	By Weight
Thifensulfuron-methyl Methyl 3-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl) amino]carbonyl]amino]sulfonyl]-2-thiophenecarboxylate	50%
Tribenuron-methyl Methyl 2-[[[N-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)methylamino]carbonyl]amino]sulfonyl]benzoate	25%
Inert Ingredients	25%
TOTAL	100%

EPA Reg. No. 352-538

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! Avoid contact with skin, eyes and clothing. In case of contact with eyes, immediately flush with plenty of water. Get medical attention if irritation persists. Wash thoroughly after handling.

For medical emergencies involving this product, call toll free 1-800-441-3637.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Waterproof gloves.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Ensure that all operation employees accurately measure pesticides.
- Mix only enough product for the job at hand.
- Avoid overfilling of spray tank.
- Do not discharge excess material on the soil at a single spot in the field, grove, or mixing/loading station.
- Dilute and agitate excess solution and apply at labeled rates or uses.
- Avoid storage of pesticides near well sites.
- When triple-rinsing the pesticide container, be sure to add the rinsate to the spray mix.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Waterproof gloves.
- Shoes plus socks.

HARMONY EXTRA is recommended for use on wheat, barley, oat, and fallow in most states, check with your state extension or Dept. of Agriculture before use, to be certain HARMONY EXTRA is registered in your state. HARMONY EXTRA is not registered for use in Alamosa, Conejos, Costilla, Rio Grande, and Saguache counties of Colorado - unless directed otherwise by supplemental labeling.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by DuPont.

GENERAL INFORMATION

DuPont Harmony® Extra Herbicide is a dry flowable granule that is used for selective postemergence weed control in wheat (including durum), barley, oat and fallow. The best control is obtained when HARMONY EXTRA is applied to young, actively growing weeds. The use rate will depend on weed spectrum and size of weed at time of application. The degree and duration of control may depend on the following:

- weed spectrum and infestation intensity
- weed size at application
- environmental conditions at and following treatment

HARMONY EXTRA is noncorrosive, nonflammable, nonvolatile, and does not freeze. HARMONY EXTRA should be mixed in water and applied as a uniform broadcast spray.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

HARMONY EXTRA is absorbed primarily through the foliage of plants, rapidly inhibiting the growth of susceptible weeds. One to 3 weeks after application to weeds (2 to 5 weeks for wild garlic), leaves of susceptible plants appear chlorotic, and the growing point subsequently dies.

HARMONY EXTRA provides the best control in vigorously growing crops that shade competitive weeds. Weed control in areas of thin crop stand or seeding skips may not be as satisfactory. However, a crop canopy that is too dense at application can intercept spray and reduce weed control.

The herbicidal action of HARMONY EXTRA may be affected in crops stressed from adverse environmental conditions (such as extreme temperatures or moisture), abnormal soil conditions, cultural practices, or variations in crop variety. In warm, moist conditions, the expression of herbicide symptoms is accelerated; in cold, dry conditions, expression of herbicide symptoms is delayed. In addition, weeds hardened-off by drought stress are less susceptible to HARMONY EXTRA.

APPLICATION INFORMATION

USE RATE

Do not use less than 0.3 ounce HARMONY EXTRA per acre.

Wheat (including Durum) and Barley

Apply 0.3 to 0.6 oz HARMONY EXTRA per acre to wheat (including durum) or barley. Two applications of HARMONY EXTRA may be made provided the total amount applied does not exceed 1.0 oz per acre per crop season.

Use 0.3 to 0.4 oz HARMONY EXTRA per acre for light infestation of the weeds listed under Weeds Controlled. Conditions at application should be optimum for effective treatment of these weeds.

Use 0.5 oz HARMONY EXTRA per acre for heavy infestation of the weeds listed under Weeds Partially Controlled.

Use 0.6 oz HARMONY EXTRA per acre for heavy infestation of the weeds listed under Weeds Partially Controlled when application timing and environmental conditions are marginal (refer to Environmental Conditions and Biological Activity for best performance).

Oat (Spring and Winter)

Apply 0.3 to 0.4 ounce HARMONY EXTRA per acre for control of the weeds listed in WEEDS CONTROLLED table. Do not make more than one application of HARMONY EXTRA per crop season on oat.

Fallow

Apply 0.3 to 0.6 oz HARMONY EXTRA per acre to fallow. Two applications of HARMONY EXTRA may be made provided the total amount applied does not exceed 1.0 oz per acre per crop season.

HARMONY EXTRA should be applied in combination with other suitable registered fallow herbicides such as Landmaster¹ II, Fallow Master¹, Roundup¹ plus 2,4-D (ester formulations work best), "Roundup" plus Banvel² "Banvel" SGF³ "Clarity"³, 2,4-D, "Banvel"² "Banvel" SGF³ "Clarity"³.

Pre-plant Burndown

Apply 0.3 to 0.6 oz HARMONY EXTRA per acre as a burndown treatment prior to, or shortly after planting (prior to emergence).

Sequential treatments of HARMONY EXTRA may also be made provided the total amount of HARMONY EXTRA applied during one fallow/pre plant cropland season does not exceed 1.0 ounce per acre; for example, 0.5 ounce in the fall followed by 0.5 ounce in the spring.

Use 0.6 ounce per acre rate when weed infestation is heavy and predominantly consists of those weeds listed under PARTIAL CONTROL, or when application timing and environmental conditions are marginal.

(See APPLICATION TIMING Section for restriction on planting intervals)

APPLICATION TIMING

Wheat (Including Durum), Barley and Winter Oat

Make applications after the crop is in the 2-leaf stage, but before the flag leaf is visible.

Spring Oat

Make applications after the crop is in the 3 leaf stage, but before jointing.

Do not use on "Ogle", "Porter" or "Premier" varieties as crop injury can occur.

Fallow

HARMONY EXTRA may be used as a fallow treatment, in the spring or fall when the majority of weeds have emerged and are actively growing.

Pre-plant Burndown

Apply HARMONY EXTRA as a burndown treatment to wheat (including durum), barley, and oat to control emerged weeds prior to, or shortly after planting (prior to emergence). Make applications when the majority of weeds have emerged and are actively growing.

Applications to sugarbeets, winter rape and canola can be made at least 60 days prior to planting. Applications to any other crop can be made at least 45 days prior to planting (crops such as corn, cotton, rice, grain sorghum or soybeans).

Since HARMONY EXTRA has very little or no soil activity, it controls only those weeds that have germinated; therefore, apply HARMONY EXTRA when all or most of the weeds have germinated. Annual broadleaf weeds should be past the cotyledon stage, actively growing, and less than 4" tall or wide. Wild garlic plants should be less than 12" tall with 2" to 4" of new growth. See Specific Weed Problems for more information.

Rainfall immediately after treatment can wash HARMONY EXTRA off of weed foliage, resulting in reduced weed control. Several hours of dry weather are needed to allow HARMONY EXTRA to be sufficiently absorbed by weed foliage.

CROP ROTATION

Wheat, Barley, and Oat may be replanted anytime after the application of HARMONY EXTRA.

Sugarbeets, Winter Rape, and Canola can be planted 60 days after the application of HARMONY EXTRA. Any other crop may be planted 45 days after the application of HARMONY EXTRA.

WEEDS CONTROLLED

HARMONY EXTRA effectively controls the following weeds when used according to label directions:

- | | |
|--------------------------------------|--------------------------------|
| Annual knawel | London rocket |
| Annual sowthistle | Marshelder |
| Black mustard | Mayweed chamomile |
| Blue/Purple mustard | Miners lettuce |
| Broadleaf dock | Narrowleaf lambsquarters |
| Bur buttercup | Nightflowering catchfly |
| Bushy wallflower/
Treacle mustard | Pennsylvania smartweed |
| Clasping pepperweed | Pineappleweed |
| Coast fiddleneck | Prickly lettuce * |
| Common buckwheat | Prostrate knotweed |
| Common chickweed | Prostrate pigweed |
| Common cocklebur * | Redmaids |
| Common groundsel | Redroot pigweed |
| Common lambsquarters | Russian thistle * |
| Common radish | Scentless chamomile
mayweed |
| Common ragweed * | Shepherd's-purse |
| Common sunflower | Slimleaf lambsquarters |
| Corn chamomile | Smallflower buttercup |
| Corn gromwell* | Smallseed falseflax |
| Corn spurry | Stinking chickweed |
| Cowcockle | Stinking mayweed/
dogfennel |
| Cress (mouse-ear) | Swinecress |
| Curly dock | Tansymustard |
| False chamomile | Tarweed fiddleneck |
| Field chickweed | Tumble/ Jim Hill mustard |
| Field pennycress | Volunteer lentils |
| Filaree (redstem, Texas) | Volunteer peas |
| Flixweed | Volunteer sunflower |
| Green smartweed | Wild buckwheat* |
| Henbit | Wild chamomile |
| Kochia * | Wild garlic* |
| Ladysthumb | Wild mustard |
| Lanceleaf sage * | Wild radish* |

WEEDS PARTIALLY CONTROLLED**

HARMONY EXTRA partially controls the following weeds when used according to label directions:

Canada thistle*	Mallow (common, little)
Carolina geranium	Nightshade (cutleaf, hairy)
Catchweed bedstraw	Vetch* (common, hairy)
Cutleaf eveningprimrose	

* See SPECIFIC WEED PROBLEMS for more information.

**Partial control: A visual reduction of weed population as well as a significant loss of vigor. For better results, use the highest recommended rate of HARMONY EXTRA per acre and include a tank mix partner such as 2,4-D, MCPA, Buctril[®] or "Banvel"/"Banvel" SGF/ "Clarity" (refer to TANK MIXTURES).

SPECIFIC WEED PROBLEMS

Canada thistle: For control in wheat and barley, use 0.6 oz per acre plus surfactant when all thistles are 4" to 8" with 2" to 6" of new growth. Make the application in the spring. Control will be improved by using HARMONY EXTRA in combination with 2,4-D (refer to TANK MIXTURES).

For control in oat, use 0.4 ounce HARMONY EXTRA per acre plus 2,4-D (refer to TANK MIXTURES)

Common cocklebur, Common ragweed, Lanceleaf sage: In wheat and barley, apply HARMONY EXTRA at 0.4 to 0.5 ounce per acre in combination with 2, 4-D at rates from 1/4 to 3/8 lb active ingredient (ester formulations work best) when weeds are small and actively growing. When using 1/4 lb active ingredient of 2, 4-D, be sure to add surfactant at the rate of 1/4 to 1/2 quart per 100 gallons of spray solution (0.06 to 0.125% v/v--use the higher rate under stress conditions).

For control in oat, use 0.4 ounce HARMONY EXTRA per acre plus 2,4-D. Refer to the Tank Mixtures sections of this label for additional details.

Corn growwell, Wild buckwheat: For control in wheat and barley, use 0.5 to 0.6 ounce HARMONY EXTRA per acre plus surfactant.

For control in oat, use 0.4 ounce HARMONY EXTRA per acre plus 2,4-D, MCPA or "Buctril" (refer to TANK MIXTURES).

Kochia, Russian thistle, Prickly lettuce: Naturally occurring resistant biotypes of these weeds are known to occur. For best results, use HARMONY EXTRA in a tank mix with dicamba (such as "Banvel"/"Banvel" SGF/"Clarity") and 2, 4-D; or Bromoxynil (such as "Buctril") and 2,4-D (3/4 - 1 pt "Buctril" + 1/4 - 3/8 lb active ingredient 2, 4-D ester). HARMONY EXTRA should be applied in the spring when weeds are less than 2" tall or 2" across and are actively growing. Refer to the Tank Mixtures section of this label for additional details.

Vetch (common and hairy): For control in wheat and barley, use 0.5 to 0.6 oz of HARMONY EXTRA per acre plus surfactant when vetch is less than 6" in length. For severe infestations of vetch, or when vetch is greater than 6" in length, use HARMONY EXTRA in combination with 2,4-D or MCPA (refer to the Tank Mixtures section of this label).

For control in oat, use 0.4 ounce HARMONY EXTRA per acre plus 2,4-D or MCPA (refer to TANK MIXTURES).

6 7 12
Wild garlic: For control in wheat and barley, use 0.5 to 0.6 oz HARMONY EXTRA per acre plus surfactant when wild garlic plants are less than 12" tall with 2" to 4" of new growth. For severe infestations, use the 0.6 ounce per acre rate of HARMONY EXTRA. Plants hardened-off by cold weather and/or drought stress may be more difficult to control. Thorough spray coverage of all garlic plants is essential. Typical symptoms of dying garlic plants may not be noticeable for 2 to 5 weeks.

For control in oat, use 0.4 ounce HARMONY EXTRA per acre plus 2,4-D or MCPA (refer to TANK MIXTURES).

Wild radish: For best results in wheat and barley, apply 0.4 to 0.6 ounce HARMONY EXTRA per acre plus surfactant either in the fall or spring to wild radish rosettes less than 6 inches in diameter. Applications made later than 30 days after weed emergence will result in partial control. For increased control of severe wild radish infestations, or wild radish emerged greater than 30 days, apply HARMONY EXTRA at 0.3 ounce per acre in combination with MCPA at 1/4 lb active ingredient per acre. Surfactant is required when tank mixing with MCPA, add 1 quart per 100 gallons of spray solution (0.25% vol/vol). Fall applications should be made prior to hardening off of plants.

For control in oat, use 0.4 ounce HARMONY EXTRA per acre plus 2,4-D or MCPA (refer to TANK MIXTURES).

SURFACTANTS

Unless otherwise specified, add a DuPont recommended nonionic surfactant having at least 80% active ingredient at 1 to 2 qt per 100 gal of spray solution (0.25 to 0.5% v/v - refer to TANK MIXTURES for specific adjuvant recommendations when HARMONY EXTRA is used in a tank mix).

Consult your agricultural dealer, applicator, or DuPont representative for a listing of recommended surfactants. Antifoaming agents may be used if needed.

Do not use low rates of liquid nitrogen fertilizer solution as a substitute for surfactant.

GROUND APPLICATION

For optimum spray distribution and thorough coverage, use flat-fan or low-volume flood nozzles.

For flat-fan nozzles, use a spray volume of at least 5 gal per acre (GPA).

For flood nozzles on 30" spacings, use at least 10 GPA, flood nozzles no larger than TK10 (or the equivalent), and a pressure of at least 30 psi. For 40" nozzle spacings, use at least 13 GPA; for 60" spacings use at least 20 GPA. It is essential to overlap the nozzles 100% for all spacings.

Raindrop "RA" nozzles are not recommended for HARMONY EXTRA applications, as weed control performance may be reduced.

Use screens that are 50-mesh or larger.

7 4 12

Chemigation - Refer to specific supplemental labeling for use directions for HARMONY EXTRA herbicide in chemigation systems. Do not apply this product through any irrigation system unless the supplemental labeling on chemigation is followed.

AERIAL APPLICATION

Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage at 2 to 5 GPA. Use at least 3 GPA in Idaho, Oregon, or Utah.

Do not apply HARMONY EXTRA by air in the state of New York.

See the **Spray Drift Management** section of this label.

PRODUCT MEASUREMENT

HARMONY EXTRA is measured using the HARMONY EXTRA volumetric measuring cylinder. The degree of accuracy of this cylinder varies by $\pm 7.5\%$. For more precise measurement, use scales calibrated in ounces.

TANK MIXTURES

HARMONY EXTRA may be tank mixed with other suitable registered herbicides to control weeds listed as suppressed, weeds resistant to HARMONY EXTRA or weeds not listed under Weeds Controlled. Read and follow all manufacturer's label recommendations for the companion herbicide. If those recommendations conflict with this label, do not tank mix the herbicide with HARMONY EXTRA.

HARMONY EXTRA can also be mixed with registered fungicides, insecticides, or liquid fertilizer for use on wheat, barley, oat, or fallow.

With 2,4-D (amine or ester) or MCPA (amine or ester)

HARMONY EXTRA may be tank mixed with the amine and ester formulations 2, 4-D and MCPA herbicides for use on wheat, barley, and oat.

For best results in the Red River Valley and adjacent areas of North Dakota and Minnesota, add the ester formulations of 2,4-D or MCPA herbicides to the tank at 3/8 lb active ingredient (such as 3/4 pt of a 4 lb/gal product, 1/2 pt of a 6 lb/gal product). No additional surfactant is needed with this mixture.

For best results in other areas, add the ester formulations of 2, 4-D or MCPA herbicides to the tank at 1/4 to 3/8 lb active ingredient (such as 1/2 - 3/4 pt of a 4 lb/gal product, 1/3 - 1/2 pt of a 6 lb/gal product). Surfactant may be added to the mixture at 1/2 to 1 qt per 100 gal of spray solution (0.125 to 0.25% v/v); however, adding surfactant may increase the potential for crop injury, especially at the higher phenoxy rates.

Higher rates of 2, 4-D or MCPA may be used, but do not exceed the highest rate allowed by those respective labels. Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using these tank mixtures.

With dicamba (such as "Banvel" / "Banvel" SGF / "Clarity")

HARMONY EXTRA may be tank mixed with 1/16 to 1/8 lb active ingredient dicamba (such as 2-4 fluid oz "Banvel", 4-8 fluid oz "Banvel" SGF, 2-4 fluid oz "Clarity"). Use higher rates when weed infestation is heavy. Surfactant may be added to the mixture at 1/2 to 1 qt per 100 gal of spray solution (0.125 to 0.25% v/v); however, adding surfactant may increase the potential for crop injury. Refer to the specific dicamba label for application timing and restrictions. Tank mixes of HARMONY EXTRA plus dicamba may result in reduced control of some broadleaf weeds.

With 2,4-D (amine or ester) and "Banvel" / "Clarity"

HARMONY EXTRA may be applied in a 3-way tank mix with formulations of dicamba and 2,4-D. Make application of HARMONY EXTRA + 1/16 to 1/8 lb active ingredient dicamba (such as 2 - 4 fluid oz "Banvel", 4 - 8 fluid oz "Banvel" SGF, 2 - 4 fluid oz "Clarity") + 1/4 - 3/8 lb active ingredient 2,4-D Ester or Amine per acre. Use higher rates when weed infestation is heavy. Surfactant may be added to the mixture at 1/2 to 1 qt per 100 gal of spray solution (0.125 to 0.25% v/v); however, adding surfactant may increase the potential for crop injury. Consult the specific 2,4-D label, dicamba label, or local recommendations for more information and restrictions.

Apply this 3-way combination to winter wheat and winter oat after the crop is tillering and prior to jointing (first node). In Spring Wheat (including Durum) and Spring Oat, apply after the crop is tillering and before it exceeds the 5-leaf stage.

In Spring Barley, apply after the crop is tillering and before it exceeds the 4-leaf stage.

With bromoxynil (such as BUCTRIL, BRONATE[®])

HARMONY EXTRA may be tank mixed with bromoxynil containing herbicides registered for use on wheat, barley, or fallow. For best results, add bromoxynil containing herbicides to the tank at 3/16 to 3/8 lb active ingredient per acre (such as "Bronate" or "Buctril" at 3/4 - 1 1/2 pt per acre).

Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using these tank mixtures. Follow the most restrictive labeling. Tank mixes of HARMONY EXTRA plus "Buctril" may result in reduced control of Canada thistle.

With DuPont Express[®] Herbicide

HARMONY EXTRA may be tank mixed with EXPRESS based on local recommendations. Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using this tank mixture.

With DuPont Ally[®] Herbicide

HARMONY EXTRA may be tank mixed with ALLY based on local recommendations. Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using this tank mixture.

With Other Herbicides

Tank mixes of HARMONY EXTRA plus metribuzin may result in reduced control of wild garlic.

With Hoelon[®] Herbicide

HARMONY EXTRA may be used in combination with "Hoelon" 3EC and "Buctril" herbicides in accordance with the "Hoelon" 3EC label. For best results, use the three-way tank mix of HARMONY EXTRA at 0.4 oz per acre plus "Hoelon" 3EC at 2 2/3 pt per acre plus "Buctril" at 1 1/2 pt per acre. Apply only to winter wheat. This tank mix should only be used under good soil conditions when wild oat is in the 1-4 leaf stage. If conditions are not ideal for the performance of "Hoelon" 3EC, wild oat control may be reduced. Be sure to follow all warnings and cautions on the "Hoelon" 3EC and "Buctril" labels

With Assert[®] Herbicide or Avenge[®] Herbicide

HARMONY EXTRA can be tank mixed with "Avenge" or "Assert". When tank mixing HARMONY EXTRA with "Assert", always include another broadleaf weed herbicide with a different mode of action (for example: 2,4-D ester, MCPA ester, "Buctril," or "Bronate"). Tank-mixed applications of HARMONY EXTRA plus "Assert" may cause temporary crop discoloration, stunting, or injury when heavy rainfall occurs shortly after application.

With other grass control products

Tank mixtures of HARMONY EXTRA and grass control products may result in poor grass control. DuPont recommends that you first consult your state experiment station, university, or extension agent, Agricultural dealer, or DuPont representative as to the potential for antagonism before using the mixture. If no information is available, limit the initial use of HARMONY EXTRA and the grass product to a small area.

With Insecticides

HARMONY EXTRA may be tank mixed or used sequentially with insecticides (or fungicides) registered for use on cereal grains. However, under certain conditions (drought stress, or if the crop is in the 2-4 leaf stage), tank mixes or sequential applications of HARMONY EXTRA with organophosphate insecticides (such as parathion) may produce temporary crop yellowing or, in severe cases, crop injury. Test these mixtures in a small area before treating large areas.

Do not use HARMONY EXTRA plus Malathion, as crop injury will result.

With Liquid Nitrogen Solution Fertilizer

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing HARMONY EXTRA in fertilizer solution. HARMONY EXTRA must first be slurried with water and then added to liquid nitrogen solutions (e.g., 28-0-0, 32-0-0). Ensure that the agitator is running while the HARMONY EXTRA is added. Use of this mixture may result in temporary crop yellowing and stunting.

If using low rates of liquid nitrogen fertilizer in the spray solution (less than 50% of the spray solution volume), the addition of surfactant is necessary. Add surfactant at 1/4 qt - 1 qt per 100 gal of spray solution (0.06 -0.25% v/v) based on local recommendations.

When using high rates of liquid nitrogen fertilizer solution in the spray solution, adding surfactant increases the risk of crop injury. Consult your agricultural dealer, consultant,

fieldman, or DuPont representative for a specific recommendation before adding an adjuvant to these tank mixtures.

If 2,4-D or MCPA is included with HARMONY EXTRA and fertilizer mixture, ester formulations tend to be more compatible (See manufacturer's label). Additional surfactant is not needed when using HARMONY EXTRA in tank mix with 2,4-D ester or MCPA ester and liquid nitrogen fertilizer solutions.

Do not use low rates of liquid nitrogen fertilizer solution as a substitute for a surfactant.

Do not use with liquid fertilizer solutions with a pH less than 3.0.

TANK MIXTURES IN FALLOW

HARMONY EXTRA may be used as a fallow treatment, and should be tank mixed with other herbicides that are registered for use in fallow.

Read and follow all manufacturer's label recommendations for the companion herbicide. If those recommendations conflict with this label, do not tank mix the herbicide with HARMONY EXTRA.

TANK MIXTURES IN PRE-PLANT BURNDOWN

HARMONY EXTRA may be used as a pre-plant burndown treatment alone or tank mixed with other herbicides that are registered for use as a pre-plant burndown product, such as "Landmaster" II, "Fallow Master", "Roundup" plus "Banvel" /"Banvel" SGF, "Clarity", or "Banvel"/ "Banvel" SGF/ "Clarity" alone.

Read and follow all manufacturer's label recommendations for the companion herbicide. If those recommendations conflict with this label, follow the most restrictive labeling (such as planting interval after application), or do not tank mix the herbicide with HARMONY EXTRA.

MIXING INSTRUCTIONS

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of HARMONY EXTRA
3. Continue agitation until the HARMONY EXTRA is fully dispersed, at least 5 minutes.
4. Once the HARMONY EXTRA is fully dispersed, maintain agitation and continue filling tank with water. HARMONY EXTRA should be thoroughly mixed with water before adding any other material.
5. As the tank is filling, add tank mix partners (if desired) then add the required volume of nonionic surfactant. Always add surfactant last. Do not use with spray additives that alter the pH of the spray solution below pH 5.0 or above pH 9.0, as rapid product degradation can occur. Spray solutions of pH 6.0-8.0 allow for optimum stability of HARMONY EXTRA.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Apply HARMONY EXTRA spray mixture within 24 hours of mixing to avoid product degradation.
8. If HARMONY EXTRA and a tank mix partner are to be applied in multiple loads, pre-slurry the HARMONY

EXTRA in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the HARMONY EXTRA.

GRAZING

Do not graze livestock in treated areas. In addition, do not feed forage or hay from treated areas to livestock (harvested straw may be used for bedding and/or feed).

SPRAY EQUIPMENT

For specific application equipment, refer to the manufacturer's recommendations for additional information on GPA, pressure, speed, nozzle types and arrangements, nozzle heights above the target canopy, etc.

Be sure to calibrate air or ground equipment properly before application. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern with minimum drift. Use higher spray volumes to obtain better coverage when crop canopy is dense. Avoid swath overlapping, and shut off spray booms while starting, turning, slowing, or stopping, to avoid injury to the crop.

Do not make applications using equipment and/or spray volumes or during weather conditions that might cause spray to drift onto nontarget sites. For additional information on spray drift refer to Spray Drift Management section of label.

Continuous agitation is required to keep HARMONY EXTRA in suspension.

SPRAYER CLEANUP

The spray equipment must be cleaned before HARMONY EXTRA is sprayed. Follow the cleanup procedures specified on the labels of the previously applied products. If no directions are provided, follow the six steps outlined in After Spraying HARMONY EXTRA.

AT THE END OF THE DAY

It is recommended that during periods when multiple loads of HARMONY EXTRA herbicide are applied, at the end of each day of spraying the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits which can accumulate in the application equipment.

AFTER SPRAYING HARMONY EXTRA AND BEFORE SPRAYING CROPS OTHER THAN WHEAT, BARLEY AND OAT

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of HARMONY EXTRA as follows:

1. Drain tank; thoroughly rinse spray tanks, boom, and hoses with clean water. Loosen and physically remove any visible deposits.
2. Fill the tank with clean water and 1 gal of household ammonia* (contains 3% active) for every 100 gal of water. Flush the hoses, boom, and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 min. Flush the hoses, boom, and nozzles again with the cleaning solution, and then drain the tank.

3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.

4. Repeat step 2.

5. Rinse the tank, boom, and hoses with clean water.

6. If only Ammonia is used as a cleaner, the rinsate solution may be applied back to the crop(s) recommended on this label. Do not exceed the maximum labeled use rate. If other cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

* Equivalent amounts of an alternate-strength ammonia solution or a DuPont-approved cleaner can be used in the cleanout procedure. Carefully read and follow the individual cleaner instructions. Consult your Ag dealer, applicator, or DuPont representative for a listing of approved cleaners.

Notes:

1. **CAUTION:** Do not use chlorine bleach with ammonia as dangerous gases will form. Do not clean equipment in an enclosed area.
2. Steam-cleaning aerial spray tanks is recommended prior to performing the above cleanout procedure to facilitate the removal of any caked deposits.
3. When HARMONY EXTRA is tank mixed with other pesticides, all cleanout procedures should be examined and the most rigorous procedure should be followed.
4. In addition to this cleanout procedure, all pre-cleanout guidelines on subsequently applied products should be followed as per the individual labels.
5. Where routine spraying practices include shared equipment frequently being switched between applications of HARMONY EXTRA and applications of other pesticides to HARMONY EXTRA-sensitive crops during the same spray season, it is recommended that a sprayer be dedicated to HARMONY EXTRA to further reduce the chance of crop injury.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150 - 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **Wind, Temperature and Humidity, and Temperature Inversions** sections of this label.

Controlling Droplet Size - General Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - The boom length should not exceed 3/4 of the wing or rotor length - longer booms increase drift potential.
- **Application Height** - Application more than 10 ft above the canopy increases the potential for spray drift.

BOOM HEIGHT

Setting the boom at the lowest referenced height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID GUSTY AND WINDLESS CONDITIONS.**

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly

dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring.

Note: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the spray equipment section of this label to determine if use of an air assist sprayer is recommended.

RESISTANCE

Biotypes of certain weeds listed on this label are resistant to HARMONY EXTRA and other herbicides with the same mode of action,* even at exaggerated application rates. Biotypes are naturally occurring individuals of a species identical in appearance but with slightly different genetic compositions: the mode of action of a herbicide is the chemical interaction that interrupts a biological process necessary for plant growth and development.

If weed control is unsatisfactory, it may be necessary to respray problem areas using a product with a different mode of action. If resistant weed biotypes such as kochia and Russian thistle are suspected or known to be present, consider using another herbicide treatment or adjust the use rate of the HARMONY EXTRA tank-mix partner to help control these biotypes.

To better manage weed resistance when using HARMONY EXTRA use a combination of tillage, and tank-mix partners or sequential herbicide applications that have a different mode of action than HARMONY EXTRA to control escaped weeds. Do not let weed escapes go to seed.

Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative herbicide recommendations available in your area.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes.

* Naturally occurring weed biotypes that are resistant to DuPont ALLY* Herbicide, DuPont EXPRESS* Herbicide, DuPont FINESSE* Herbicide or DuPont GLEAN* FC Herbicide will also be resistant to HARMONY EXTRA.

INTEGRATED PEST MANAGEMENT

DuPont recommends the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an Integrated Pest Management (IPM) program which can include biological, cultural, and genetic practices aimed at preventing economic pest damage. Application of this product should be based on IPM principles and practices

including field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

PRECAUTIONS

Injury to or loss of adjacent sensitive crops, desirable trees or vegetation may result from failure to observe the following:

- Do not apply, drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants.
- Take all necessary precautions to avoid all direct or indirect contact (such as spray drift) with non-target plants or areas.
- Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than wheat, barley or oat.

HARMONY EXTRA is only registered on wheat, barley, oat and fallow. Do not use on any other crop.

The total rate of HARMONY EXTRA for wheat (including durum) and barley cannot exceed 1.0 ounce product per acre applied to any one crop during one growing season.

The total rate of HARMONY EXTRA for oat (spring and winter) cannot exceed 0.4 ounces product per acre applied to any one crop during one growing season.

Varieties of wheat (including durum) and barley may differ in their response to various herbicides. DuPont recommends that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use to a small area.

Under certain conditions such as heavy rainfall, prolonged cold weather, or wide fluctuations in day/night temperatures prior to or soon after HARMONY EXTRA application, temporary discoloration and/or crop injury may occur. To reduce the potential of crop injury, tank mix HARMONY EXTRA with 2,4-D (ester formulations perform best—see Tank Mixtures) and apply after the crop is in the tillering stage of growth.

HARMONY EXTRA should not be applied to wheat, barley or oat that is stressed by severe weather conditions, drought, low fertility, water-saturated soil, disease, or insect damage, as crop injury may result. Risk of injury is greatest when crop is in the 2 to 5- leaf stage. Severe winter stress, drought, disease, or insect damage following application also may result in crop injury.

Do not apply to wheat, barley or oat crops underseeded with another crop.

Dry, dusty field conditions may result in reduced control in wheel track areas.

Do not harvest sooner than 45 days after the last application of HARMONY EXTRA.

STORAGE AND DISPOSAL

Storage: Store the product in original container only. Do not contaminate water, other pesticides, fertilizer, food, or feed in storage.

Product Disposal: Do not contaminate water, food, or feed by disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent) the container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

- 1 Registered trademark of Monsanto
- 2 Registered trademark of BASF Corp.
- 3 Registered trademark of Rhone-Poulenc Ag Company.
- 4 Registered trademark of AgrEvo USA
- 5 Registered trademark of American Cyanamid Company.

SL - 643 9019 1/15/99

For product information call 1-888-6-DUPONT

© 1999 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898

10 2 12

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants.

DuPont does not agree to be an insurer of these risks. WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.